

Nuclear factor kappa B role in inflammation associated gastrointestinal malignancies

Sahil Gambhir, Dinesh Vyas, Michael Hollis, Apporva Aekka, Arpita Vyas

Sahil Gambhir, Dinesh Vyas, Michael Hollis, Apporva Aekka, Department of Surgery, College of Human Medicine, Michigan State University, East Lansing, MI 48912, United States

Arpita Vyas, Department of Pediatrics, College of Human Medicine - Michigan State University, East Lansing, MI 48912, United States

Author contributions: Gambhir S and Vyas D contributed equally to this work; Gambhir S and Vyas D designed research; analyzed data and wrote the paper; Hollis M contributed in research and editing; Vyas A contributed in data and conceptualization; Aekka A contributed in editing.

Conflict-of-interest: No potential conflicts of interest relevant to this article were reported.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Correspondence to: Dinesh Vyas, MD, MS, FICS, FACS, Department of Surgery, College of Human Medicine, Michigan State University, 1200 East Michigan Avenue, Suite 655, East Lansing, MI 48912, United States. dinesh.vyas@hc.msu.edu

Telephone: +517-2672491

Fax: +1-517-2672488

Received: June 15, 2014

Peer-review started: June 16, 2014

First decision: August 15, 2014

Revised: August 29, 2014

Accepted: February 12, 2015

Article in press: February 13, 2015

Published online: March 21, 2015

Abstract

Nuclear factor kappa B (NF- κ B) has an established role in the regulation of innate immunity and inflammation. NF- κ B is also involved in critical mechanisms connecting inflammation and cancer development. Recent investigations

suggest that the NF- κ B signaling cascade may be the central mediator of gastrointestinal malignancies including esophageal, gastric and colorectal cancers. This review will explore NF- κ B's function in inflammation-associated gastrointestinal malignancies, highlighting its oncogenic contribution to each step of carcinogenesis. NF- κ B's role in the inflammation-to-carcinoma sequence in gastrointestinal malignancies warrants stronger emphasis upon targeting this pathway in achieving greater therapeutic efficacy.

Key words: Gastrointestinal cancer; Inflammation; Nuclear factor kappa B

© **The Author(s) 2015.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Inflammation has a critical role in cancer, its metastasis and angiogenesis. One of the critical mediator is nuclear transcription factor nuclear factor kappa B. This article will be critically useful for basic scientist and clinicians in understanding the the importance of this transcription factor.

Gambhir S, Vyas D, Hollis M, Aekka A, Vyas A. Nuclear factor kappa B role in inflammation associated gastrointestinal malignancies. *World J Gastroenterol* 2015; 21(11): 3174-3183 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v21/i11/3174.htm> DOI: <http://dx.doi.org/10.3748/wjg.v21.i11.3174>

INTRODUCTION

Gastrointestinal cancers, comprised of esophageal, gastric and colorectal cancers, are some of the most common and fatal cancers worldwide. However, mortality rates have declined due to improved efforts in cancer prevention, earlier detection and a growing number of treatment options, especially in the development of therapies targeted to specific signaling pathways. These

therapies are founded upon the body of basic scientific research that has implicated various signaling pathways in different gastrointestinal (GI) malignancies. Recent literature suggests that the nuclear factor kappa B (NF- κ B) signaling cascade in particular may serve as a central mediator of carcinogenesis. The NF- κ B signaling pathway has numerous physiological roles, including regulation of innate immunity and inflammation, and therefore may be involved in carcinogenesis in multiple ways^[1,2]. The primary function of this signaling cascade is to protect the cell from harm, but if it becomes aberrant, then the transition from inflammation to cancerous growth can be elicited. In our previous study, we explored gastrointestinal epithelial response to injury and by extension intend to scrutinize NF- κ B's effects on the gut in this review^[3].

Recent epidemiological investigations have shown that chronic inflammation, including infection, is associated with 15%-20% of all human malignancies^[4]. Numerous genetic studies have also exposed NF- κ B's potential role in inflammation-related cancers^[4-7]. For example, NF- κ B's constitutive activation has been demonstrated in various human solid tumors, including those of ovarian, lung, breast, melanoma, hepatocellular, thyroid, pancreatic, prostate, colon, esophageal, gastric, laryngeal, parathyroid, bladder, endometrial, retinoblastoma, astrocytoma and squamous cell carcinoma of the head and neck^[8-17]. Moreover, aberrant NF- κ B regulation has been observed in hematopoietic malignancies, such as multiple myeloma, mantle cell lymphoma, MALT lymphoma, diffuse large B-cell lymphoma, Hodgkin's lymphoma, myelodysplastic syndrome, adult T - cell leukemia, chronic lymphocytic leukemia, acute myeloid leukemia, and chronic myeloid leukemia^[8,11,18,19].

NF- κ B is a pleiotropic transcriptional factor that is encoded by a family of five main genes^[5,20]. These genes contain the homologous domain Rel homology domain (RHD), which includes RelA (p65), RelB, cRel (REI), p50 and p52. NF- κ B1 (p100) and NF- κ B2 (p102) are processed to produce p52 and p50 respectively^[21]. The RHD allows these proteins to homo- and heterodimerize, localize to the nucleus, bind to DNA and interact with the inhibitor of NF- κ B (I κ B)^[22]. NF- κ B homodimers or heterodimers remain inhibited in the cytoplasm bound to ankyrin rich regions of I κ B proteins, which prevent NF- κ B translocation to the nucleus and thus transcription^[22]. The I κ B family consists of I κ B α , I κ B β , I κ B γ , I κ B ϵ , BCL-3 and p100 and p105. It is through I κ B kinase dependent (IKK-dependent) phosphorylation, polyubiquitination and proteasomal degradation of I κ B proteins that the NF- κ B proteins are liberated^[22]. The IKK family consists of two catalytic subunits, IKK α and IKK β , complexed to the regulatory subunit IKK γ /NEMO (NF- κ B essential modulator)^[22].

The tight regulation and activation of NF- κ B is controlled by two principal signaling pathways, modulated by divergent but specific stimuli: the classical or canonical pathway and the alternative

pathway^[1,22]. With regards to the former, inflammatory stimuli, including TNF- α , IL-1 β , TLRs and viruses, promote phosphorylation at two serine sites of I κ B through the activation of IKK α , IKK β and NEMO^[21]. This leads to polyubiquitination at adjacent lysine residues and eventually proteolysis, releasing NF- κ B dimers-most commonly p50/p65 heterodimers-and degradation of I κ B^[5]. The alternative pathway is characterized by its independence from IKK β and NEMO and reliance on IKK α ^[22]. Following stimulation by TNF- α receptor superfamily members (lymphotoxin (LT) β R, B-cell activating factor receptor (BAFFR), receptor activator of NF- κ B (RANK), and CD40), IKK mediates the phosphorylation of NF- κ B2/p100:RelB complexed with an I κ B^[22]. The complex is subsequently polyubiquitinated, leading to the proteasome-dependent processing of RelB's inhibitor NF- κ B2 p100, which results in the liberation of the RelB-p50 heterodimer^[1,21]. In both pathways, the dimers proceed to translocate to the nucleus, bind to NF- κ B DNA sites and activate transcription. It should be noted that one of the first proteins transcribed in this process is I κ B α , which then translocates to the nucleus and binds to the dimer to inhibit further transcription^[5]. The classical pathway is implicated in preventing apoptosis and in innate immunity and inflammation^[1,21]. The alternative pathway is involved in secondary lymphoid organogenesis and B cell maturation and survival^[21]. Thus, both are responsible for cell survival and ultimately play a role in carcinogenesis.

NF- κ B AND CANCER

Virchow first postulated and proposed the causal relationship between chronic inflammation and cancer. Chronic inflammation due to irritants is thought to lead to increased cell multiplication, predisposing cells to neoplastic DNA changes. An important mechanistic link may include NF- κ B, as recent studies have suggested that NF- κ B promotes tumor growth by inducing downstream proteins that have oncogenic effects^[22]. Initial evidence of NF- κ B's link to cancer was the identification of the p50 subunit and RelA and its close homology with the oncoprotein v-REL of the avian REL retrovirus^[23]. The onco-virus constitutively activates the v-REL oncoprotein, which is an important catalyst in the progression of lymphomas^[23]. Similarly, the human T-cell leukemia virus's oncoprotein TAX has been shown to constitutively activate the IKK complex, stimulating both pathways and again leading to upregulation of NF- κ B's downstream proteins^[24].

Hanahan and Weinberg's postulation of each step of tumorigenesis includes self-sufficiency in growth signals, insensitivity to growth inhibitor signals, evasion of apoptosis, limitless replicative potential, sustained angiogenesis, and tissue invasion and metastasis^[25-28]. NF- κ B can exert its effects on each of these aspects of tumorigenesis through the induction of downstream protein expression, and thus might be the basis of the transition from inflammation to cancer growth.

Essentially, NF- κ B controls cell proliferation by activating growth factors, including IL-2, granulocyte monocyte colony stimulator factor and CD40L^[25,26]. It also serves as a positive regulator of cell cycle progression as it can activate c-myc and cyclin D1^[26]. NF- κ B inhibition of programmed cell death, through its regulation of the anti-apoptotic proteins cIAPs, c-FLIP and members of the bcl-2 family, is also well documented^[25,26]. These proteins are vital in sustaining genetically altered precancerous cells, and their effects therefore increase the probability of malignant changes. NF- κ B activity can also lead to increased angiogenesis and metastasis through the upregulation of chemokines, such as IL-8 (migration), VEGF (angiogenesis) and MMP (spread). The altered expression of these genes, with NF- κ B as a key mediator, is associated with tumor growth and spread^[25,26]. Finally, the heavily scrutinized pro-inflammatory gene COX-2, which is under direct control of NF- κ B, has been demonstrated to be closely involved in the process of carcinogenesis^[25-30].

This review article will explore NF- κ B's role in cancers of the gastrointestinal tract, including esophageal, gastric and colonic. Discussion of each cancer will begin with a presentation of inflammatory stimuli, including infection and other inciting factors that affect NF- κ B's expression. Evidence will then be provided in support of the fact that NF- κ B activity increases through progression of inflammation to cancer on a continuum that culminates in constitutive expression. The documented upregulation of NF- κ B's downstream proteins in each specific GI malignancy will be discussed next, with emphasis of COX-2 expression. This will be followed by an illustration of the importance of targeting the NF- κ B pathway for greater therapeutic efficacy, including addressing chemoresistance. In essence, the amalgamation of the data presented in this review will uncover NF- κ B as the mechanistic link between inflammation and tumor growth, specifically in gastrointestinal cancers. Finally, common themes will be clarified in emphasizing the connection between the two entities.

NF- κ B'S ROLE IN ESOPHAGEAL CARCINOGENESIS

Esophageal cancer is responsible for roughly 16000 deaths in the United States per year; mortality depends on cell type (adenocarcinoma vs squamous), risk factors and location of pathology^[31]. There is increasing evidence that the NF- κ B pathway mediates the progression of inflamed esophageal epithelium through carcinogenesis^[30,31]. This begins when the upper GI tract is exposed to noxious stimuli that promote inflammation and foster a propensity for cancerous growth^[30,31]. Examples of such stimuli include tobacco smoke, alcohol, reflux, corrosive injury, nitrosamines, HPV and thermal injury^[31]. Moreover, multiple studies link exposure to upregulation of NF- κ B. One study conducted by Abdel-

Latif *et al*^[32] showed that bile acids and low pH induced expression of NF- κ B in esophageal cell lines. Similarly, another study illustrated that physiological levels of bile acid not only activated NF- κ B, but also led to increased expression of the downstream protein IL-8^[33]. It is suggested that after epithelial cells are exposed to inflammatory stimuli, they defensively resort to activating the NF- κ B pathway in order to prevent apoptosis and sustain existence by DNA repair^[34]. This persistence predisposes the cells to malignancy as they continue to proliferate due to genetic alterations^[35].

Involvement of NF- κ B in promoting inflammatory changes to metaplasia and the progression to esophageal cancer has been demonstrated in the literature^[36]. Current evidence supports constitutive NF- κ B activity in cancerous growth of both adenocarcinoma and squamous cell carcinoma in *in vivo* and *in vitro* studies^[11,36-38]. In one study, nuclear immunologic staining revealed increased nuclear expression of p50, p52 and RelI in patient biopsies of esophageal cancer cells compared to normal squamous cells^[25]. Abdel-Latif *et al*^[38] showed that overexpression of NF- κ B in tumor tissues correlated with higher expression of Rel-A and lower expression of I κ B when compared to expression levels in non-tumor cells. This was further supported by the finding that NF- κ B expression was increased in patients of Barrett's epithelium, however there was virtually no expression in patients with normal esophageal epithelium^[38,39]. In another study, 40% NF- κ B expression was noted in Barrett's tissues, but markedly increased to 76% in adenocarcinomas^[36]. In addition to demonstrating NF- κ B's presence in esophageal malignancy, studies have also correlated NF- κ B activity with metastasis^[40].

It is the downstream proteins that are responsible for exerting oncogenic effects, however, and these are vital in inflammation-related tumorigenesis. A number of studies have examined this increase in inflammatory proteins specific to esophageal malignancy. O'Riordan *et al*^[37] demonstrated the elevation of proinflammatory cytokines IL-8 and IL-1B in esophagitis, Barrett's epithelium and adenocarcinoma in human histological specimens. These cytokines contribute to tumor progression by regulating angiogenesis, sustaining cancer cell growth and promoting tumor cell migration. Investigators further found a significant association of NF- κ B activation and cytokine upregulation in adenocarcinoma. Similar results have been found in other studies with regard to increased pro-inflammatory cytokines in esophageal cancerous growth^[11,38]. Recent evidence also supports NF- κ B's role in invasion and metastasis in esophageal carcinomas as it leads to the upregulation of MMP-9 and reduction of E cadherin, two proteins involved in cell migration^[41].

Another example illustrating NF- κ B's involvement in neoplastic progression is the overexpression of COX-2, an enzyme important in regulating prostaglandin synthesis^[42]. In the context of cancer, increased COX-2 activity has been shown to be associated with key pathways that control cell proliferation, migration,

apoptosis, and angiogenesis^[11]. Previous studies have revealed increased simultaneous expression of NF- κ B and COX-2 in cells exposed to inflammation and cells of malignant growth when compared to normal esophageal mucosa^[29]. Studies exploring the effect of COX-2 inhibitors have also shed light upon the relationship between COX-2 and progression to malignancy. Liu *et al.*^[43] studied the effects of aspirin at different concentrations and different times in esophageal squamous cell carcinoma cell lines and found that aspirin significantly reduced COX-2 mRNA and protein expression and prostaglandin synthesis. The study's results demonstrate a dose-dependent response to aspirin in reducing cell proliferation and inducing apoptosis. In a recent meta-analysis, use of aspirin and other COX inhibitors was shown to be associated with reduced risk of esophageal adenocarcinoma among patients. They noted that the use of COX inhibitors decreased the risk of transition from Barrett's esophagus to esophageal cancer. Many *in vivo* and *in vitro* studies have reported similar results and advocate for aspirin use in both preventing and treating abnormal esophageal changes.

Patients with esophageal cancer often have a poor prognosis due to late presentation of the disease. Therefore, it is important to explore new treatments, such as those that target NF- κ B^[44,45]. In esophageal adenocarcinoma, elevated NF- κ B expression is associated with advanced stages and inversely correlated with response to neoadjuvant chemotherapy and radiation^[46,47]. NF- κ B expression in esophageal cancers is indicative of poor prognosis and may be implicated in multidrug resistance and treatment failure. In a recent study, NF- κ B activity in patients with esophageal squamous and adenocarcinoma correlated with TMN staging, increased susceptibility to spread, chemoresistance and overall poor survival rate^[48]. Interestingly, their data also demonstrated patients who were established to have NF- κ B negative cancer prior to treatment became positive after receiving chemotherapy, suggesting chemotherapy as a potential initiator of tumor resistance through upregulation of NF- κ B^[48]. Treatment with chemotherapy may induce resistance to the therapy, and therefore, result in poorer prognosis. In addition, Izzo *et al.*^[48] showed that after constitutive NF- κ B activity was established, exposure to NF- κ B inhibitors, such as Bay11-7082 and sulfasalazine, reduced cancer cell proliferation and induced apoptosis. They also found that treatment with these inhibitors in combination with chemotherapeutic drugs 5-fluorouracil and cisplatin led to a synergistic effect on inhibiting cell growth, decreasing chemoresistance^[11]. This investigation also illustrated that Bay 11 was associated with decreased tumor growth and angiogenesis in animal studies^[11]. Tian *et al.*^[49] found similar results with siRNA molecules of p65. They first found p65 expression was decreased in ESCC cell lines when exposed to siRNA p65 after establishing constitutive activity of the NF- κ B signaling pathway. Moreover, they too found a synergistic effect

of siRNA with 5FU as ESCC cells became more sensitive to 5-FU after exposure to siRNA p65. Finally, utilization of neoadjuvant treatment with NF- κ B inhibitors in the management of esophageal carcinomas was shown to help in a specific cohort of refractory treatment cases^[46,47].

NF- κ B'S ROLE IN GASTRIC CARCINOGENESIS

Gastric cancer further elucidates the role of NF- κ B as a central mediator of carcinogenesis and as an important link between inflammation and cancer^[50,51]. Gastric cancer, especially the intestinal type, entails the progression from chronic gastritis to chronic atrophic gastritis to intestinal metaplasia, then dysplasia and finally adenocarcinoma^[52]. The initiation of chronic inflammation typically begins with the presence of such risk factors as *Helicobacter pylori* (*H. pylori*) infection, pernicious anemia, smoking or high salt diet. This leads to longstanding chronic superficial gastritis, which eventually progresses to intestinal metaplasia followed by adenocarcinoma^[53,54]. *H. pylori*'s detrimental effect on gastric mucosa has been well established in the literature through clinical data, animal studies and cell cultures^[52,53]. However, only certain *H. pylori* strains have been shown to be involved in malignancy and these have the pathogenic island, CagA^[54]. Current studies have established that CagA-positive *H. pylori* significantly contributes to gastric cancer cell tissue invasion^[55,56]. Wu *et al.*^[57] visualized the link between CagA-positive *H. pylori* and NF- κ B by using NF- κ B blockers. Exposing gastric cancer cells infected by CagA-positive strains to NF- κ B blockers significantly reduced tissue invasiveness. This not only supported CagA's effects as a pathogenic factor, but demonstrated the significant role of NF- κ B. These data support the idea that *H. pylori* becomes a potent NF- κ B activator if it carries CagA pathogenicity. The exact molecular pathway between CagA and NF- κ B cannot be addressed due to a paucity of evidence. Some studies speculate the connection lies in the protein transforming growth factor-beta-activated kinase 1 (TAK1)^[54].

Constitutive activation of the NF- κ B pathway and its increased production in gastric cancer cells of human tissue and cell lines has been demonstrated through various laboratory techniques, including IHC, EMSA and Western blotting^[5,57,58]. Constitutive NF- κ B activity has also been documented in *H. pylori* gastritis and diffuse gastric cancer^[11,59]. Although chronic irritants initiate mucosal inflammation, it is the NF- κ B pathway which is vital in sustaining the gastritis-metaplasia-carcinoma sequence. Studies have shown that NF- κ B is notably higher in patients with *H. pylori* gastritis, and this is significantly correlated with histological scores of gastritis^[60]. In intestinal type gastric cancer, increased NF- κ B expression is strongly associated with CagA strain-induced metaplasia, dysplasia and gastric

cancer itself^[56]. This is currently under debate for diffuse type gastric cancer. Yamanaka *et al*^[61] identified a major function of NF- κ B in carcinogenesis when they found that patients with higher NF- κ B activity had a shorter survival rate and concluded that NF- κ B was a prognostic indicator in gastric cancer. Interestingly, NF- κ B expression is also associated with stage, depth of invasion, WHO classification and Lauren's histological classification^[11,51,52,58,62]. Levidou *et al*^[11] further concluded through multivariate survival analysis that NF- κ B1 expression was an independent predictor of gastric cancer prognosis.

For the gastritis-to-carcinoma sequence to be successful, each step requires a pattern of oncoprotein expression to promote carcinogenesis. As in esophageal cancer, NF- κ B regulates these oncoproteins in gastric cancer as well in terms of endorsing tumor growth, preventing apoptosis, and initiating tumor angiogenesis and migration. Keates *et al*^[63] demonstrated an increased activation of NF- κ B and its downstream proteins by infecting gastric epithelial cells with *H. pylori* and then observing increases in p50/p65 heterodimers and p50 dimers, followed by increased IL-8 mRNA and protein synthesis. Yin *et al*^[64] found that IL-6 and VEGF were significantly increased in NF- κ B positive gastric cancer cells compared to their levels in normal mucosa. Like IL-8, IL-6 is also associated with malignancy-both facilitate cancer cell apoptosis and stimulation of angiogenesis. VEGF is a well-known growth factor implicated in angiogenesis and therefore is crucial in carcinogenesis^[65]. Other studies have shown that gastric cell lines exposed to *H. pylori* have displayed upregulation of important tumorigenesis markers including MMP-9, VEGF and COX-2^[50].

NF- κ B is also an important regulator of COX-2 in gastric cancer as it is in esophageal cancer. Expression of NF- κ B and COX-2 in the same neoplastic mucosa has been established in previous studies^[66,67]. Studies exploring COX-2 inhibitors also shed light upon the specific actions of COX-2 and its potential applications^[58,68]. In one study, p50 positive cells treated with COX inhibitors showed a dose-dependent suppression of cell growth in gastric cells^[66]. A large population-based case-control study conducted by Farrow *et al*^[69] revealed that patients who took aspirin or NSAIDs had a decreased risk of gastric and esophageal cancer compared to never users. Evidence provides support that COX-2 inhibitors may prevent gastric carcinogenesis and can aid in treating gastric malignancy^[69]. COX-2 expression is, in part, facilitated by NF- κ B expression; if the central regulator can be inhibited, then all downstream proteins, including COX-2, may be suppressed, thereby resulting in reduced oncogenesis^[50].

Despite advances in current therapeutic approaches, gastric cancer is still one of the most prevalent cancers and the second leading cause of death due to cancer, worldwide. Therefore, new treatment options should be explored by scrutinizing the molecular pathway of the gastritis-adenocarcinoma sequence. The NF- κ B pathway provides a link between inflammation

and cancer and can explain many of the hallmarks of cancer^[50,52]. Moreover, literature has illustrated the dysregulation of the NF- κ B signaling pathway in gastric cancer, with its presence indicating poor prognosis^[61]. Studies have shown in CagA-mediated gastric cancer that cell migration is attenuated by NF- κ B inhibitors. These results were expanded upon in a study conducted by Keates *et al*^[63], which found that pretreatment with PDTC, a potent NF- κ B inhibitor, reduced *H. pylori*-activated p65 and IL-8. In one study, parthenolide, another NF- κ B inhibitor, was used on three gastric cancer cell lines and significantly induced apoptosis in all three^[70]. They also found a synergistic effect when NF- κ B inhibitors were combined with chemotherapeutic drugs^[71]. This finding is analogous to that of combining NF- κ B inhibitors and chemotherapy in addressing esophageal cancers.

NF- κ B'S ROLE IN COLORECTAL CARCINOGENESIS

The development of colorectal cancer (CRC) is a multistep process and there is growing evidence in favor of the connection between inflammation and carcinogenesis. NF- κ B has become the main focus of this neoplastic transformation. CRC can be divided into sporadic CRC, hereditary CRC and colitis-associated carcinoma (CAC). CAC primarily stems from the two major forms of inflammatory bowel disease (IBD): ulcerative colitis and Crohn's disease. NF- κ B's role as the mechanistic link between inflammation and cancer has been intensively investigated regarding constitutive activity of NF- κ B in both IBDs and CAC^[71-74]. CAC is one of the most well-known examples of an inflammation-dysplasia-carcinoma sequence^[73,75]. IBD prompts a chronic inflammatory state leading to the constant production of noxious compounds including reactive oxygen species (ROS) and other cytokines (TNF- α , IL-6, and IL-1). In the long term, ROS are detrimental as they repress regulators of DNA damage and induce mutagenic enzymes, such as activation-induced cytidine deaminase^[74]. Furthermore, studies indicate that both ROS and cytokines activate the NF- κ B pathway^[74]. These cytokines activate signal transducer and activator of transcription 3 (STAT3) in intestinal epithelial cells. This, in turn, contributes to further inflammation, perpetuating noxious stimuli and ultimately creating a positive feedback loop. The combination of DNA damaging agents and inflammatory cytokines results in constant activation of the NF- κ B pathway, setting up the progression to carcinogenesis.

Overexpression of NF- κ B has been demonstrated in colon cancer cell lines and human tumor specimens, including those of sporadic CRC and CAC^[74,76]. This is synonymous with genetic syndromes including familial adenomatous polyposis (FAP) and hereditary non-polyposis cancer (HNPCC)^[72]. The expression of

NF- κ B was determined to be significantly higher in the adenocarcinoma tissue compared to expression levels in tissue specimens earlier in the inflammation-adenocarcinoma sequence^[77,78]. Furthermore, studies have shown that NF- κ B activity increases in association with histological tumor progression^[74,79]. This indicates that NF- κ B activity correspondingly increases from the initiation of inflammation to malignant proliferation. Notably, one study concluded that NF- κ B activation in CRC served as a poor prognostic indicator^[78].

As in esophageal and gastric cancer, NF- κ B regulates expression of various oncoproteins in colorectal cancer as well to sustain inflammation and support cell proliferation, inhibition of apoptosis, angiogenesis, invasion and metastasis. Overexpression of NF- κ B and its downstream effectors are well recognized in CRC^[56,79]. For example, IL-8 upregulation was also described in CRC and was identified in various ways including microarray and protein array analyses^[28]. In addition, another cytokine, IL-6, has been found to be weakly correlated with poor prognosis in CRC. In a well-known study, Greten *et al.*^[24] investigated the role of NF- κ B in inflammation-associated tumor growth using a mouse model and found that IKK β deletion led to a direct decrease in tumor incidence. They attributed these results to decreased expression of anti-apoptotic proteins. CRC also has a significant association between NF- κ B overexpression and VEGF^[24].

Another emerging theme is the relationship between NF- κ B and its downstream protein COX-2. Studies of CRC have shown an upregulation of both NF- κ B and COX-2 and the literature also supports their individual and coexistent overexpression^[23,24,27,75,78-82]. Maihöfner *et al.*^[83] studied surgically resected tissues of CRC patients and found a high expression of p65, IL-6 and COX-2 compared to controls^[81,82]. It is well established that regular aspirin use is associated with a significant reduction in the risk of COX-2-positive CRC^[80]. The relative risk of CRC is reduced by 40%-50% when patients consume aspirin or NSAIDs over a period of 10-15 years^[24]. Moreover, epidemiologic evidence shows that NSAID usage protects against CRC to a greater extent than against other GI malignancies. In one study, aspirin exposure resulted in apoptosis of colorectal cancer cell lines in a concentration-dependent manner^[83]. Specifically, given that CAC is the result of inflammatory processes, it follows that anti-inflammatory medication would reduce the risk of cancer.

Inhibition of the master regulator NF- κ B will shut off the different steps of carcinogenesis. NF- κ B inhibitors, such as sulfasalazine, mesalamine, and glucocorticoids, have already shown promise in treating IBD^[23]. Gan *et al.*^[84] explored the effects of sulfasalazine on the NF- κ B signaling pathway. After establishing that patients with ulcerative colitis (UC) had higher NF- κ B and downstream proteins including IL-6 and IL-8, those who were exposed to sulfasalazine demonstrated a decrease in all three proteins^[84]. Corticosteroids strongly inhibit NF- κ B activation *in vivo* and *in vitro*, and clinical trials have found dexamethasone to be useful in inhibiting CRC

metastasis. One study illustrated this by using NF- κ B inhibitors and displaying that blocking the action of NF- κ B led to inhibition of angiogenesis in CRC^[79]. Another study reported these same inhibitors prompted apoptosis, as NF- κ B could not overcome the anti-apoptotic machinery to exert pro-survival activity^[72]. After establishing constitutive activity of NF- κ B in CRC, Lind *et al.*^[72] further showed that NF- κ B binding was increased in CRC cell lines after exposure to the chemotherapeutic agent. In contrast, pretreating cells with NF- κ B inhibitors prior to administration of gemcitabine showed a decrease in tumor size. The common theme of chemoresistance resurfaces and NF- κ B may be involved in its development. With regards to CRC, NF- κ B inhibitors are already showing clinical applications. Further research should be employed, as this pathway is involved in many innate processes and therefore may have a large range of systemic side effects.

CONCLUSION

This review showed support for NF- κ B's role in inflammation-associated cancers. NF- κ B activation is elicited by various inflammatory stimuli (Figure 1). Chronic irritation induces constitutive NF- κ B activity, promoting carcinogenesis (Figure 1). NF- κ B's mechanistic link to cancer is best displayed through its pleiotropic effects, as it leads to upregulation of important proteins that promote tumor progression (Figure 1). Upregulation of COX-2 by NF- κ B in GI malignancies is consistently supported in the literature and the efficacy of COX-2 inhibitors is still being scrutinized. This review also confirmed the presence of NF- κ B in advanced gastrointestinal malignancies. Additionally, this review noted that NF- κ B may play a role in developing chemo-resistance in gastrointestinal malignancies. Recent investigations by Vyas *et al.*^[85] postulate that chemotherapies may pose therapy-induced resistance by stimulating various signaling pathways including the NF- κ B cascade. The authors clearly delineated literature supporting that first-line chemotherapy agents such as doxorubicin, 5-FU, cisplatin and paclitaxel commonly induce NF- κ B production. They showed that this led to increased expression of downstream proteins that promote proliferation, anti-apoptosis and angiogenesis to sustain tumor growth^[85]. These findings demonstrated the need for further research in discovering common themes among the gastrointestinal malignancies. Interestingly, Kim *et al.*^[86] recently reported a significant increase in the protein caspase-associated recruitment domain 6 (CARD6), a NF- κ B activator in esophageal, gastric and colorectal tissues^[87]. Finally, targeted therapy focusing upon NF- κ B inhibition and its outcomes for the individual cancers was also discussed. The amalgamation of all the literature presented in this review emphasizes the importance of exploring molecular targeted therapy to hone in on NF- κ B. In addition, although NF- κ B is the central mediator,




Figure 1 Summary model representing stimulation of the NF- κ B pathway leading to the classical pathway, followed by expression of NF- κ B dependent proteins. Analogous inflammatory stimuli are common amongst esophageal, gastric and colorectal tissue. The unique stimuli commonly induce inflammation. The classic pathway is displayed and this model would be consistent with the alternate pathway. An array of genes are transcribed and translated contributing to the carcinogenesis. ROS: Reactive oxygen species.

targeting the regulators of the actual pathway itself could also prove fruitful. For instance, the use of histone deacetylases (HDACs) has gained popularity as an emerging strategy in inhibiting the NF- κ B pathway. Histone acetyltransferases and deacetylases modulate NF- κ B activity through their actions on the Re1A/p65 subunit^[87]. Yun *et al*^[88] showed acetylation of Re1A/p65 subsequently increased NF- κ B activation and deacetylation lead to diminished levels of Re1A/p65 and thus NF- κ B. This was further explored when Yeung *et al*^[89] demonstrated that the use of sirtuins, a group of nicotinamide adenosine dinucleotide-dependent HDACs, led to increased apoptosis as these compounds decreased NF- κ B levels; this is yet another potential strategy for intervention in the future.

NF- κ B may induce inflammatory-associated gastrointestinal carcinomas which are often refractory to current treatments. We, therefore, urge greater exploration and development of therapies specifically targeting the NF- κ B pathway as these may prove more successful for patients than existing therapeutic options.

REFERENCES

- 1 **Baud V, Karin M.** Is NF-kappaB a good target for cancer therapy? Hopes and pitfalls. *Nat Rev Drug Discov* 2009; **8**: 33-40 [PMID: 19116625 DOI: 10.1038/nrd2781]
- 2 **Bonizzi G, Karin M.** The two NF-kappaB activation pathways and their role in innate and adaptive immunity. *Trends Immunol* 2004; **25**: 280-288 [PMID: 15145317 DOI: 10.1016/j.it.2004.03.008]
- 3 **Vyas D, Robertson CM, Stromberg PE, Martin JR, Dunne WM, Houchen CW, Barrett TA, Ayala A, Perl M, Buchman TG, Coopersmith CM.** Epithelial apoptosis in mechanistically distinct methods of injury in the murine small intestine. *Histol Histopathol* 2007; **22**: 623-630 [PMID: 17357092]
- 4 **Karin M, Cao Y, Greten FR, Li ZW.** NF-kappaB in cancer: from innocent bystander to major culprit. *Nat Rev Cancer* 2002; **2**: 301-310 [PMID: 12001991 DOI: 10.1038/nrc780]
- 5 **Karin M.** NF-kappaB and cancer: mechanisms and targets. *Mol Carcinog* 2006; **45**: 355-361 [PMID: 16673382 DOI: 10.1002/mc.20217]
- 6 **DiDonato JA, Mercurio F, Karin M.** NF- κ B and the link between inflammation and cancer. *Immunol Rev* 2012; **246**: 379-400 [PMID: 22435567 DOI: 10.1111/j.1600-065X.2012.01099.x]
- 7 **Ditsworth D, Zong WX.** NF-kappaB: key mediator of inflammation-associated cancer. *Cancer Biol Ther* 2004; **3**: 1214-1216 [PMID: 15611628 DOI: 10.4161/cbt.3.12.1391]
- 8 **Bours V, Dejardin E, Goujon-Letawe F, Merville MP, Castronovo V.** The NF-kappa B transcription factor and cancer: high expression of NF-kappa B- and I kappa B-related proteins in tumor cell lines. *Biochem Pharmacol* 1994; **47**: 145-149 [PMID: 8311838 DOI: 10.1016/0006-2952(94)90448-0]
- 9 **Claudio E, Brown K, Park S, Wang H, Siebenlist U.** BAFF-induced NEMO-independent processing of NF-kappa B2 in maturing B cells. *Nat Immunol* 2002; **3**: 958-965 [PMID: 12352969 DOI: 10.1038/ni842]
- 10 **Howe LR.** Inflammation and breast cancer. Cyclooxygenase/prostaglandin signaling and breast cancer. *Breast Cancer Res* 2007; **9**: 210 [PMID: 17640394 DOI: 10.1186/bcr1678]
- 11 **Levidou G, Korkolopoulou P, Nikiteas N, Tzanakis N, Thymara I, Saetta AA, Tsigris C, Rallis G, Vlasik K, Patsouris E.** Expression of Nuclear Factor κ B in Human Gastric Carcinoma: Relationship

- with I κ Ba and Prognostic Significance. *Virchows Archiv* 2007; **450**: 519-527 [PMID: 17429689 DOI: 10.1007/s00428-007-0396-5]
- 12 **Mukhopadhyay T**, Roth JA, Maxwell SA. Altered expression of the p50 subunit of the NF-kappa B transcription factor complex in non-small cell lung carcinoma. *Oncogene* 1995; **11**: 999-1003 [PMID: 7675461]
 - 13 **Nakshatri H**, Bhat-Nakshatri P, Martin DA, Goulet RJ, Sledge GW. Constitutive activation of NF-kappaB during progression of breast cancer to hormone-independent growth. *Mol Cell Biol* 1997; **17**: 3629-3639 [PMID: 9199297]
 - 14 **Shattuck-Brandt RL**, Richmond A. Enhanced degradation of I-kappaB alpha contributes to endogenous activation of NF-kappaB in Hs294T melanoma cells. *Cancer Res* 1997; **57**: 3032-3039 [PMID: 9230219]
 - 15 **Tai DI**, Tsai SL, Chang YH, Huang SN, Chen TC, Chang KS, Liaw YF. Constitutive activation of nuclear factor kappaB in hepatocellular carcinoma. *Cancer* 2000; **89**: 2274-2281 [PMID: 11147598 DOI: 10.1002/1097-0142]
 - 16 **Visconti R**, Cerutti J, Battista S, Fedele M, Trapasso F, Zeki K, Miano MP, de Nigris F, Casalino L, Curcio F, Santoro M, Fusco A. Expression of the neoplastic phenotype by human thyroid carcinoma cell lines requires NFkappaB p65 protein expression. *Oncogene* 1997; **15**: 1987-1994 [PMID: 9365245 DOI: 10.1038/sj.onc.1201373]
 - 17 **Wang W**, Abbruzzese JL, Evans DB, Larry L, Cleary KR, Chiao PJ. The nuclear factor-kappa B RelA transcription factor is constitutively activated in human pancreatic adenocarcinoma cells. *Clin Cancer Res* 1999; **5**: 119-127 [PMID: 9918209]
 - 18 **Bargou RC**, Leng C, Krappmann D, Emmerich F, Mapara MY, Bommert K, Royer HD, Scheidereit C, Dörken B. High-level nuclear NF-kappa B and Oct-2 is a common feature of cultured Hodgkin/Reed-Sternberg cells. *Blood* 1996; **87**: 4340-4347 [PMID: 8639794]
 - 19 **Feinman R**, Koury J, Thames M, Barlogie B, Epstein J, Siegel DS. Role of NF-kappaB in the rescue of multiple myeloma cells from glucocorticoid-induced apoptosis by bcl-2. *Blood* 1999; **93**: 3044-3052 [PMID: 10216101]
 - 20 **Bassères DS**, Baldwin AS. Nuclear factor-kappaB and inhibitor of kappaB kinase pathways in oncogenic initiation and progression. *Oncogene* 2006; **25**: 6817-6830 [PMID: 17072330 DOI: 10.1038/sj.onc.1209942]
 - 21 **Dolcet X**, Llobet D, Pallares J, Matias-Guiu X. NF-kB in development and progression of human cancer. *Virchows Arch* 2005; **446**: 475-482 [PMID: 15856292 DOI: 10.1007/s00428-005-1264-9]
 - 22 **Brown KD**, Estefania Claudio, and Ulrich Siebenlist. The Roles of the Classical and Alternative Nuclear Factor-kappaB Pathways: Potential Implications for Autoimmunity and Rheumatoid Arthritis. *Arthritis Research Therapy* 2008; **10**: 212 [PMID: 18771589 DOI: 10.1186/ar2457]
 - 23 **Greten FR**, Karin M. The IKK/NF-kappaB activation pathway-a target for prevention and treatment of cancer. *Cancer Lett* 2004; **206**: 193-199 [PMID: 15013524 DOI: 10.1016/j.canlet.2003.08.029]
 - 24 **Greten FR**, Eckmann L, Greten TF, Park JM, Li ZW, Egan LJ, Kagnoff MF, Karin M. IKKbeta links inflammation and tumorigenesis in a mouse model of colitis-associated cancer. *Cell* 2004; **118**: 285-296 [PMID: 15294155 DOI: 10.1016/j.cell.2004.07.013]
 - 25 **Kang MR**, Kim MS, Kim SS, Ahn CH, Yoo NJ, Lee SH. NF-kappaB signalling proteins p50/p105, p52/p100, RelA, and IKKepsilon are over-expressed in oesophageal squamous cell carcinomas. *Pathology* 2009; **41**: 622-625 [PMID: 20001340 DOI: 10.3109/00313020903257756]
 - 26 **Karin M**, Lin A. NF-kappaB at the crossroads of life and death. *Nat Immunol* 2002; **3**: 221-227 [PMID: 11875461 DOI: 10.1038/ni0302-221]
 - 27 **Rogler G**, Brand K, Vogl D, Page S, Hofmeister R, Andus T, Knuechel R, Baeuerle PA, Schölmerich J, Gross V. Nuclear Factor κ B Is Activated in Macrophages and Epithelial Cells of Inflamed Intestinal Mucosa. *Gastroenterology* 1998; **115**: 357-369 [PMID: 9679041 DOI: 10.1016/S0016-5085(98)70202-1]
 - 28 **Zubair A**, Frieri M. Role of nuclear factor- κ B in breast and colorectal cancer. *Curr Allergy Asthma Rep* 2013; **13**: 44-49 [PMID: 22956391 DOI: 10.1007/s11882-012-0300-5]
 - 29 **Konturek PC**, Nikiforuk A, Kania J, Raithe M, Hahn EG, Mühlendorfer S. Activation of NFkappaB represents the central event in the neoplastic progression associated with Barrett's esophagus: a possible link to the inflammation and overexpression of COX-2, PPARgamma and growth factors. *Digest Dis Sci* 2004; **49**: 1075-1083 [PMID: 15387324]
 - 30 **Chan AT**, Ogino S, Fuchs CS. Aspirin and the risk of colorectal cancer in relation to the expression of COX-2. *N Engl J Med* 2007; **356**: 2131-2142 [PMID: 17522398 DOI: 10.1056/NEJMoa067208]
 - 31 **Gibson MK**, Dhaliwal AS, Clemons NJ, Phillips WA, Dvorak K, Tong D, Law S, Pirchi ED, Räsänen J, Krasna MJ, Parikh K, Krishnadath KK, Chen Y, Griffiths L, Colleypriest BJ, Farrant JM, Tosh D, Das KM, Bajpai M. Barrett's esophagus: cancer and molecular biology. *Ann N Y Acad Sci* 2013; **1300**: 296-314 [PMID: 24117650 DOI: 10.1111/nyas.12252]
 - 32 **Abdel-Latif MM**, O'Riordan J, Windle HJ, Carton E, Ravi N, Kelleher D, Reynolds JV. NF-kappaB activation in esophageal adenocarcinoma: relationship to Barrett's metaplasia, survival, and response to neoadjuvant chemoradiotherapy. *Ann Surg* 2004; **239**: 491-500 [PMID: 15024310 DOI: 10.1097/01.sla.0000118751.95179.c6]
 - 33 **Jenkins GJ**, Harries K, Doak SH, Wilmes A, Griffiths AP, Baxter JN, Parry JM. The bile acid deoxycholic acid (DCA) at neutral pH activates NF-kappaB and induces IL-8 expression in oesophageal cells in vitro. *Carcinogenesis* 2004; **25**: 317-323 [PMID: 14656946 DOI: 10.1152/ajpgi.00092.2011]
 - 34 **Luo JL**, Kamata H, Karin M. IKK/NF-kappaB signaling: balancing life and death--a new approach to cancer therapy. *J Clin Invest* 2005; **115**: 2625-2632 [PMID: 16200195 DOI: 10.1172/JCI26322]
 - 35 **Hormi-Carver K**, Zhang X, Zhang HY, Whitehead RH, Terada LS, Spechler SJ, Souza RF. Unlike Esophageal Squamous Cells, Barrett's Epithelial Cells Resist Apoptosis by Activating the Nuclear Factor- B Pathway. *Cancer Res* 2009; **69**: 672-677 [PMID: 19147583 DOI: 10.1158/0008-5472.CAN-08-3703]
 - 36 **Jenkins GJ**, Mikhail J, Alhamdani A, Brown TH, Caplin S, Manson JM, Bowden R, Toffazal N, Griffiths AP, Parry JM, Baxter JN. Immunohistochemical Study of Nuclear Factor- B Activity and Interleukin-8 Abundance in Oesophageal Adenocarcinoma; a Useful Strategy for Monitoring These Biomarkers. *J Clin Pathol* 2007; **60**: 1232-237 [PMID: 17220207 DOI: 10.1136/jcp.2006.043976]
 - 37 **O'Riordan JM**, Abdel-latif MM, Ravi N, McNamara D, Byrne PJ, McDonald GS, Keeling PW, Kelleher D, Reynolds JV. Proinflammatory cytokine and nuclear factor kappa-B expression along the inflammation-metaplasia-dysplasia-adenocarcinoma sequence in the esophagus. *Am J Gastroenterol* 2005; **100**: 1257-1264 [PMID: 15929754 DOI: 10.1111/j.1572-0241.2005.41338.x]
 - 38 **Abdel-Latif MMMJ**, O'riordan M, Ravi N, Kelleher D, Reynolds JV. Activated Nuclear Factor-kappa B and Cytokine Profiles in the Esophagus Parallel Tumor Regression following Neoadjuvant Chemoradiotherapy. *Dis Esoph* 2005; **18**: 246-252 [DOI: 10.1111/j.1442-2050.2005.00497.x]
 - 39 **Szachnowicz S**, Ceconello I, Iriya K, Marson AG, Takeda FR, Gama-Rodrigues JJ. Origin of adenocarcinoma in Barrett's esophagus: p53 and Ki67 expression and histopathologic background. *Clinics (Sao Paulo)* 2005; **60**: 103-112 [PMID: 15880245]
 - 40 **Izzo JG**, Correa AM, Wu TT, Malhotra U, Chao CK, Luthra R, Ensor J, Dekovich A, Liao Z, Hittelman WN, Aggarwal BB, Ajani JA. Pretherapy Nuclear Factor- B Status, Chemoradiation Resistance, and Metastatic Progression in Esophageal Carcinoma. *Mole Cancer Therap* 2006; **5**: 2844-850 [DOI: 10.1158/1535-7163.MCT-06-0351]
 - 41 **Wang F**, He W, Fanghui P, Wang L, Fan Q. NF- κ Bp65 promotes invasion and metastasis of oesophageal squamous cell cancer by regulating matrix metalloproteinase-9 and epithelial-to-mesenchymal transition. *Cell Biol Int* 2013; **37**: 780-8 [PMID: 23504993 DOI: 10.1002/cbin.10089]
 - 42 **Wang F**, He W, Fanghui P, Wang L, Fan Q. NF- κ Bp65 promotes invasion and metastasis of oesophageal squamous cell cancer

- by regulating matrix metalloproteinase-9 and epithelial-to-mesenchymal transition. *Cell Biol Int* 2013; **37**: 780-788 [PMID: 23504993 DOI: 10.1136/gut.2004.047100]
- 43 **Liu JF**, Jamieson GG, Drew PA, Zhu GJ, Zhang SW, Zhu TN, Shan BE, Wang QZ. Aspirin Induces Apoptosis in Oesophageal Cancer Cells by Inhibiting the Pathway of NF-kappaB Downstream Regulation of Cyclooxygenase-2. *ANZ J Surg* 2005; **75**: 1011-1016 [PMID: 16336399 DOI: 10.1111/j.1445-2197.2005.03596.x]
- 44 **Lin A**, Karin M. NF-kappaB in cancer: a marked target. *Semin Cancer Biol* 2003; **13**: 107-114 [PMID: 12654254]
- 45 **Monks NR**, Biswas DK, Pardee AB. Blocking anti-apoptosis as a strategy for cancer chemotherapy: NF-kappaB as a target. *J Cell Biochem* 2004; **92**: 646-650 [PMID: 15211562 DOI: 10.1002/jcb.20080]
- 46 **Li B**, Li YY, Tsao SW, Cheung ALM. Targeting NF- B Signaling Pathway Suppresses Tumor Growth, Angiogenesis, and Metastasis of Human Esophageal Cancer. *Mole Cancer Therap* 2009; **8**: 2635-2644 [PMID: 19723887 DOI: 10.1158/1535-7163.MCT-09-0162]
- 47 **Li J**, Wang K, Chen X, Meng H, Song M, Wang Y, Xu X, Bai Y. Transcriptional activation of microRNA-34a by NF-kappa B in human esophageal cancer cells. *BMC Mol Biol* 2012; **13**: 4 [PMID: 22292433 DOI: 10.1186/1471-2199-13-4]
- 48 **Izzo JG**, Malhotra U, Wu TT, Ensor J, Luthra R, Lee JH, Swisher SG, Liao Z, Chao KS, Hittelman WN, Aggarwal BB, Ajani JA. Association of Activated Transcription Factor Nuclear Factor B With Chemoradiation Resistance and Poor Outcome in Esophageal Carcinoma. *J Clin Oncol* 2006; **24**: 748-754 [PMID: 16401681 DOI: 10.1200/JCO.2005.03.8810]
- 49 **Tian F**, Zang WD, Hou WH, Liu HT, Xue LX. Nuclear Factor- κ B Signaling Pathway Constitutively Activated in Esophageal Squamous Cell Carcinoma Cell Lines and Inhibition of Growth of Cells by Small Interfering RNA. *Acta Biochim Biophys Sin* 2006; **38**: 318-326 [PMID: 16680372 DOI: 10.1111/j.1349-7006.2011.02025.x]
- 50 **Han JC**, Kai-Li Zhang, Xiao-Yan Chen, Hai-Feng Jiang, Qing-You Kong, Yuan Sun, Mo-Li Wu, Lei Huang, Hong Li, and Jia Liu. Expression of Seven Gastric Cancer-associated Genes and Its Relevance for Wnt, NF- κ B and Stat3 Signaling. *Apmis* 2007; **115**: 1331-343 [PMID: 18184402 DOI: 10.1111/j.1600-0643.2007.00695.x]
- 51 **Kwon HC**, Kim SH, Oh SY, Lee S, Lee JH, Jang JS, Kim MC, Kim KH, Kim SJ, Kim SG, Kim HJ. Clinicopathologic Significance of Expression of Nuclear Factor- κ B RelA and Its Target Gene Products in Gastric Cancer Patients. *World J Gastroenterol* 2012; **18**: 4744 [PMID: 23002344 DOI: 10.3748/wjg.v18.i34.4744]
- 52 **Peek RM**, Blaser MJ. Helicobacter pylori and gastrointestinal tract adenocarcinomas. *Nat Rev Cancer* 2002; **2**: 28-37 [PMID: 11902583 DOI: 10.1038/nrc703]
- 53 **Isomoto H**, Mizuta Y, Miyazaki M, Takeshima F, Omagari K, Murase K, Nishiyama T, Inoue K, Murata I, Kohno S. Implication of NF-kappaB in Helicobacter pylori-associated gastritis. *Am J Gastroenterol* 2000; **95**: 2768-2776 [PMID: 11051346 DOI: 10.1111/j.1572-0241.2000.02304.x]
- 54 **Lamb A**, Yang XD, Tsang YHN, Li JD, Higashi H, Hatakeyama M, Peek RM, Blanke SR, Chen LF. Helicobacter Pylori CagA Activates NF- κ B by Targeting TAK1 for TRAF6-mediated Lys 63 Ubiquitination. *EMBO Rep* 2009; **10**: 1242-1249 [PMID: 19820695 DOI: 10.1038/embor.2009.210]
- 55 **Kang DW**, Hwang WC, Park MH, Ko GH, Ha WS, Kim KS, Lee YC, Choi KY, Min DS. Rebamipide Abolishes Helicobacter Pylori CagA-induced Phospholipase D1 Expression via Inhibition of NF κ B and Suppresses Invasion of Gastric Cancer Cells. *Oncogene* 2012; **32**: 3531-3542 [PMID: 22890316 DOI: 10.1038/onc.2012.358]
- 56 **Yang GF**, Deng CS, Xiong YY, Gong LL, Wang BC, Luo J. Expression of nuclear factor-kappa B and target genes in gastric precancerous lesions and adenocarcinoma: association with Helicobacter pylori cagA (+) infection. *World J Gastroenterol* 2004; **10**: 491-496 [PMID: 14966904]
- 57 **Wu LF**, Pu Z, Feng J, Li G, Zheng Z, Shen W. The Ubiquitin-proteasome Pathway and Enhanced Activity of NF- κ B in Gastric Carcinoma. *J Surg Oncol* 2008; **97**: 439-444 [PMID: 18163448 DOI: 10.1002/jso.20952]
- 58 **Yu BC**, Jiang XH, Fan XM, Lin MCM, Jiang SH, Lam SK, Kung HF. Suppression of RelA/p65 Nuclear Translocation Independent of IkB- α Degradation by Cyclooxygenase-2 Inhibitor in Gastric Cancer. *Oncogene* 2003; **22**: 1189-197 [PMID: 12606945 DOI: 10.1038/sj.onc.1206234]
- 59 **Sasaki N**, Morisaki T, Hashizume K, Yao T, Tsuneyoshi M, Noshiro H, Nakamura K, Yamanaka T, Uchiyama A, Tanaka M, Katano M. Nuclear factor-kappaB p65 (RelA) transcription factor is constitutively activated in human gastric carcinoma tissue. *Clin Cancer Res* 2001; **7**: 4136-4142 [PMID: 11751513]
- 60 **Ooi CH**, Ivanova T, Wu J, Lee M, Tan IB, Tao J, Ward L, Koo JH, Gopalakrishnan V, Zhu Y, Cheng LL, Lee J, Rha SY, Chung HC, Ganesan K, So J, Soo KC, Lim D, Chan WH, Wong WK, Bowtell D, Yeoh KG, Grabsch H, Boussioutas A, Tan P. Oncogenic Pathway Combinations Predict Clinical Prognosis in Gastric Cancer. *PLoS Genetics* 2009; **5**: e1000676 [PMID: 19798449 DOI: 10.1371/journal.pgen.1000676]
- 61 **Yamanaka N**, Sasaki N, Tasaki A, Nakashima H, Kubo M, Morisaki T, Noshiro H, Yao T, Tsuneyoshi M, Tanaka M, Katano M. Nuclear factor-kappaB p65 is a prognostic indicator in gastric carcinoma. *Anticancer Res* 2004; **24**: 1071-1075 [PMID: 15154625]
- 62 **Long YM**, Ye S, Rong J, Xie WR. Nuclear factor kappa B: a marker of chemotherapy for human stage IV gastric carcinoma. *World J Gastroenterol* 2008; **14**: 4739-4744 [PMID: 18720533 DOI: 10.3748/wjg.14.4739]
- 63 **Keates S**, Hitti YS, Upton M, Kelly CP. Helicobacter pylori infection activates NF-kappa B in gastric epithelial cells. *Gastroenterology* 1997; **113**: 1099-1109 [PMID: 9322504]
- 64 **Yin Y**, Si X, Gao Y, Gao L, Wang J. The nuclear factor- κ B correlates with increased expression of interleukin-6 and promotes progression of gastric carcinoma. *Oncol Rep* 2013; **29**: 34-38 [PMID: 23117246 DOI: 10.3892/or.2012.2089]
- 65 **Guo Y**, Xu F, Lu T, Duan Z, Zhang Z. Interleukin-6 signaling pathway in targeted therapy for cancer. *Cancer Treat Rev* 2012; **38**: 904-910 [PMID: 22651903 DOI: 10.1016/j.ctrv.2012.04.007]
- 66 **Lim JW**, Kim H, Kim KH. Expression of Ku70 and Ku80 mediated by NF-kappa B and cyclooxygenase-2 is related to proliferation of human gastric cancer cells. *J Biol Chem* 2002; **277**: 46093-46100 [PMID: 12324457 DOI: 10.1074/jbc.M206603200]
- 67 **Liu XJ**, Chen ZF, Li HL, Hu ZN, Liu M, Tian AP, Zhao D, Wu J, Zhou YN, Qiao L. Interaction between cyclooxygenase-2, Snail, and E-cadherin in gastric cancer cells. *World J Gastroenterol* 2013; **19**: 6265-6271 [PMID: 24115825 DOI: 10.3748/wjg.v19.i37.6265]
- 68 **Wu CY**, Wang CJ, Tseng CC, Chen HP, Wu MS, Lin JT, Inoue H, Chen GH. Helicobacter pylori promote gastric cancer cells invasion through a NF-kappaB and COX-2-mediated pathway. *World J Gastroenterol* 2005; **11**: 3197-3203 [PMID: 15929167]
- 69 **Farrow DC**, Vaughan TL, Hansten PD, Stanford JL, Risch HA, Gammon MD, Chow WH, Dubrow R, Ahsan H, Mayne ST, Schoenberg JB, West AB, Rotterdam H, Fraumeni JF, Blot WJ. Use of aspirin and other nonsteroidal anti-inflammatory drugs and risk of esophageal and gastric cancer. *Cancer Epidemiol Biomarkers Prev* 1998; **7**: 97-102 [PMID: 9488582]
- 70 **Aranha MM**, Borralho PM, Ravasco P, Moreira da Silva IB, Correia L, Fernandes A, Camilo ME, Rodrigues CM. NF-kappaB and apoptosis in colorectal tumorigenesis. *Eur J Clin Invest* 2007; **37**: 416-424 [PMID: 17461988 DOI: 10.1111/j.1365-2362.2007.01801.x]
- 71 **Sohma I**, Fujiwara Y, Sugita Y, Yoshioka A, Shirakawa M, Moon JH, Takiguchi S, Miyata H, Yamasaki M, Mori M, Doki Y. Parthenolide, an NF- κ B inhibitor, suppresses tumor growth and enhances response to chemotherapy in gastric cancer. *Cancer Genomics Proteomics* 2011; **8**: 39-47 [PMID: 21289336]
- 72 **Lind D**, Hochwald SN, Malaty J, Rekkas S, Hebig P, Mishra G, Moldawer LL, Copeland EM, Mackay S. Nuclear Factor- κ B Is

- Upregulated in Colorectal Cancer. *Surgery* 2001; **130**: 363-369 [PMID: 11490372 DOI: 10.1093/abbs/gmt071]
- 73 **Shanahan F.** Review article: colitis-associated cancer -- time for new strategies. *Aliment Pharmacol Ther* 2003; **18** Suppl 2: 6-9 [PMID: 12950414 DOI: 10.1046/j.1365-2036.18.s2.5.x]
- 74 **Wang S, Liu Z, Wang L, Zhang X.** NF-kappaB signaling pathway, inflammation and colorectal cancer. *Cell Mol Immunol* 2009; **6**: 327-334 [PMID: 19887045 DOI: 10.1038/cmi.2009.43]
- 75 **Grivennikov SI.** Inflammation and colorectal cancer: colitis-associated neoplasia. *Semin Immunopathol* 2013; **35**: 229-244 [PMID: 23161445 DOI: 10.1007/s00281-012-0352-6]
- 76 **Vaiopoulos AG, Athanasoula KCh, Papavassiliou AG.** NF- κ B in colorectal cancer. *J Mol Med (Berl)* 2013; **91**: 1029-1037 [PMID: 23636511 DOI: 10.1007/s00109-013-1045-x]
- 77 **Horst D, Budczies J, Brabletz T, Kirchner T, Hlubek F.** Invasion associated up-regulation of nuclear factor kappaB target genes in colorectal cancer. *Cancer* 2009; **115**: 4946-4958 [PMID: 19658179 DOI: 10.1002/cncr.24564]
- 78 **Kojima M, Morisaki T, Sasaki N, Nakano K, Mibu R, Tanaka M, Katano M.** Increased nuclear factor-kB activation in human colorectal carcinoma and its correlation with tumor progression. *Anticancer Res* 2004; **24**: 675-681 [PMID: 15161011]
- 79 **Sakamoto K, Maeda S, Hikiba Y, Nakagawa H, Hayakawa Y, Shibata W, Yanai A, Ogura K, Omata M.** Constitutive NF- B Activation in Colorectal Carcinoma Plays a Key Role in Angiogenesis, Promoting Tumor Growth. *Clin Cancer Res* 2009; **15**: 2248-258 [PMID: 19276252 DOI: 10.1158/1078-0432.CCR-08-138]
- 80 **Din FV, Dunlop MG, Stark LA.** Evidence for colorectal cancer cell specificity of aspirin effects on NF kappa B signalling and apoptosis. *Br J Cancer* 2004; **91**: 381-388 [PMID: 15188000 DOI: 10.1038/sj.bjc.6601913]
- 81 **Charalambous MP, Maihöfner C, Bhambra U, Lightfoot T, Gooderham NJ.** Upregulation of cyclooxygenase-2 is accompanied by increased expression of nuclear factor-kappa B and I kappa B kinase-alpha in human colorectal cancer epithelial cells. *Br J Cancer* 2003; **88**: 1598-1604 [PMID: 12771929 DOI: 10.1038/sj.bjc.6600927]
- 82 **Shao J, Fujiwara T, Kadowaki Y, Fukazawa T, Waku T, Itoshima T, Yamatsuji T, Nishizaki M, Roth JA, Tanaka N.** Overexpression of the Wild-type P53 Gene Inhibits NF- κ B Activity and Synergizes with Aspirin to Induce Apoptosis in Human Colon Cancer Cells. *Oncogene* 2000; **19**: 726-736 [PMID: 10698490 DOI: 10.1038/sj.onc.1203383]
- 83 **Maihöfner C, Charalambous MP, Bhambra U, Lightfoot T, Geisslinger G, Gooderham NJ.** Expression of cyclooxygenase-2 parallels expression of interleukin-1beta, interleukin-6 and NF-kappaB in human colorectal cancer. *Carcinogenesis* 2003; **24**: 665-671 [PMID: 12727794 DOI: 10.1093/carcin/bgg006]
- 84 **Gan HT, Chen YQ, Ouyang Q.** Sulfasalazine inhibits activation of nuclear factor-kappaB in patients with ulcerative colitis. *J Gastroenterol Hepatol* 2005; **20**: 1016-1024 [PMID: 15955209 DOI: 10.1111/j.1440-1746.2005.03862.x]
- 85 **Vyas D, Laput G, Vyas AK.** Chemotherapy-enhanced inflammation may lead to the failure of therapy and metastasis. *Onco Targets Ther* 2014; **7**: 1015-1023 [PMID: 24959088]
- 86 **Kim SS, Ahn CH, Kang MR, Kim YR, Kim HS, Yoo NJ, Lee SH.** Expression of CARD6, an NF-kappaB activator, in gastric, colorectal and oesophageal cancers. *Pathology* 2010; **42**: 50-53 [PMID: 20025480 DOI: 10.3109/0013020903434421]
- 87 **Kawahara TL, Michishita E, Adler AS, Damian M, Berber E, Lin M, McCord RA, Ongaigui KC, Boxer LD, Chang HY, Chua KF.** SIRT6 links histone H3 lysine 9 deacetylation to NF-kappaB-dependent gene expression and organismal life span. *Cell* 2009; **136**: 62-74 [PMID: 19135889 DOI: 10.1016/j.cell.2008.10.052]
- 88 **Yun D, Rahmani M, Dent P, Grant S.** Blockade of Histone Deacetylase Inhibitor-Induced RelA/p65 Acetylation and NF- κ B Activation Potentiates Apoptosis in Leukemia Cells through a Process Mediated by Oxidative Damage, XIAP Downregulation, and c-Jun N-Terminal Kinase 1 Activation. *Mol Cell Biol* 2005; **25**: 1016-1024 [DOI: 10.1128/MCB.25.13.5429-5444.2005]
- 89 **Yeung F, Hoberg JE, Ramsey CS, Keller MD, Jones DR, Frye RA, Mayo MW.** Modulation of NF-kappaB-dependent transcription and cell survival by the SIRT1 deacetylase. *EMBO J* 2004; **23**: 2369-2380 [PMID: 15152190 DOI: 10.1038/sj.emboj.7600244]

P- Reviewer: Bashashati M, Passi A **S- Editor:** Qi Y
L- Editor: A **E- Editor:** Liu XM





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgooffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>



ISSN 1007-9327

