

World Journal of *Stem Cells*

Monthly Volume 17 Number 2 February 26, 2025



REVIEW

Wang YJ, Chen ZH, Shen YT, Wang KX, Han YM, Zhang C, Yang XM, Chen BQ. Stem cell therapy: A promising therapeutic approach for skeletal muscle atrophy. *World J Stem Cells* 2025; 17(2): 98693 [DOI: [10.4252/wjsc.v17.i2.98693](https://doi.org/10.4252/wjsc.v17.i2.98693)]

MINIREVIEWS

Silva B, Bragança J. Induced pluripotent stem cell-derived mesenchymal stem cells for modeling and treating metabolic associated fatty liver disease and metabolic associated steatohepatitis: Challenges and opportunities. *World J Stem Cells* 2025; 17(2): 99331 [DOI: [10.4252/wjsc.v17.i2.99331](https://doi.org/10.4252/wjsc.v17.i2.99331)]

Chen QH, Zheng JY, Wang DC. Asthma and stem cell therapy. *World J Stem Cells* 2025; 17(2): 103599 [DOI: [10.4252/wjsc.v17.i2.103599](https://doi.org/10.4252/wjsc.v17.i2.103599)]

ORIGINAL ARTICLE**Retrospective Study**

Luo ZY, Fan LQ, Guo WL, Yang JP, Li ZY, Huang YX, Jiang H, Zhang XH. Effect of tandem autologous stem cell transplantation on survival in pediatric patients with high-risk solid tumors in South China. *World J Stem Cells* 2025; 17(2): 100621 [DOI: [10.4252/wjsc.v17.i2.100621](https://doi.org/10.4252/wjsc.v17.i2.100621)]

Basic Study

Xia X, Zhou K, An LY, Zhao M, Tian BL, Zhao JY, Zhou ZG, Tong Y. Nicotinamide adenine dinucleotide rejuvenates septic bone marrow mesenchymal stem cells. *World J Stem Cells* 2025; 17(2): 96893 [DOI: [10.4252/wjsc.v17.i2.96893](https://doi.org/10.4252/wjsc.v17.i2.96893)]

Wang Y, Zhang FQ, Fan ZP, Zhu XL, Yan WH, Zhang XL. WDR36 inhibits the osteogenic differentiation and migration of periodontal ligament stem cells. *World J Stem Cells* 2025; 17(2): 99132 [DOI: [10.4252/wjsc.v17.i2.99132](https://doi.org/10.4252/wjsc.v17.i2.99132)]

Zhang HA, Zhang BY, Tang HB. Effects of macrophages on the osteogenic differentiation of adipose tissue-derived stem cells in two-dimensional and three-dimensional cocultures. *World J Stem Cells* 2025; 17(2): 99326 [DOI: [10.4252/wjsc.v17.i2.99326](https://doi.org/10.4252/wjsc.v17.i2.99326)]

Xia ZY, Wang Y, Shi N, Lu MQ, Deng YX, Qi YJ, Wang XL, Zhao J, Jiang DY. Fetal mice dermal mesenchymal stem cells promote wound healing by inducing M2 type macrophage polarization. *World J Stem Cells* 2025; 17(2): 101030 [DOI: [10.4252/wjsc.v17.i2.101030](https://doi.org/10.4252/wjsc.v17.i2.101030)]

Wang W, Yin J. Exosomal miR-203 from bone marrow stem cells targets the SOCS3/NF- κ B pathway to regulate neuroinflammation in temporal lobe epilepsy. *World J Stem Cells* 2025; 17(2): 101395 [DOI: [10.4252/wjsc.v17.i2.101395](https://doi.org/10.4252/wjsc.v17.i2.101395)]

Xu L, Liu D, Yun HL, Zhang W, Ren L, Li WW, Han C. Effect of adipose-derived stem cells exosomes cross-linked chitosan- α -glycerophosphate thermosensitive hydrogel on deep burn wounds. *World J Stem Cells* 2025; 17(2): 102091 [DOI: [10.4252/wjsc.v17.i2.102091](https://doi.org/10.4252/wjsc.v17.i2.102091)]

LETTER TO THE EDITOR

Zhang JW. Impact of miR-214-5p and miR-21-5p from hypoxic endometrial exosomes on human umbilical cord mesenchymal stem cell function. *World J Stem Cells* 2025; 17(2): 102404 [DOI: [10.4252/wjsc.v17.i2.102404](https://doi.org/10.4252/wjsc.v17.i2.102404)]

Ferreira LVO, Amorim RM. Perspectives on Schwann-like cells derived from bone marrow-mesenchymal stem cells: Advancing peripheral nerve injury therapies. *World J Stem Cells* 2025; 17(2): 102702 [DOI: [10.4252/wjsc.v17.i2.102702](https://doi.org/10.4252/wjsc.v17.i2.102702)]

ABOUT COVER

Peer Review of *World Journal of Stem Cells*, Zahra Jabbarpour, PhD, Postdoctoral Fellow, School of Life Sciences, Keele University, Guy Hilton Research Centre, Staffordshire ST4 7QB, United Kingdom. zjabbarpour@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Stem Cells* (*WJSC*, *World J Stem Cells*) is to provide scholars and readers from various fields of stem cells with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. *WJSC* publishes articles reporting research results obtained in the field of stem cell biology and regenerative medicine, related to the wide range of stem cells including embryonic stem cells, germline stem cells, tissue-specific stem cells, adult stem cells, mesenchymal stromal cells, induced pluripotent stem cells, embryonal carcinoma stem cells, hemangioblasts, lymphoid progenitor cells, *etc.*

INDEXING/ABSTRACTING

The *WJSC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, PubMed, PubMed Central, Scopus, Biological Abstracts, BIOSIS Previews, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for *WJSC* as 3.6; JIF without journal self cites: 3.5; 5-year JIF: 4.2; JIF Rank: 105/205 in cell biology; JIF Quartile: Q3; and 5-year JIF Quartile: Q2. The *WJSC*'s CiteScore for 2023 is 7.8 and Scopus CiteScore rank 2023: Histology is 11/62; Genetics is 78/347; Genetics (clinical) is 19/99; Molecular Biology is 131/410; Cell Biology is 104/285.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Xiang-Di Zhang; Production Department Director: Xiao-Mei Zheng; Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Stem Cells

ISSN

ISSN 1948-0210 (online)

LAUNCH DATE

December 31, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Shengwen Calvin Li, Carlo Ventura

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-0210/editorialboard.htm>

PUBLICATION DATE

February 26, 2025

COPYRIGHT

© 2025 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.fcpublishing.com>

Stem cell therapy: A promising therapeutic approach for skeletal muscle atrophy

Ying-Jie Wang, Ze-Hao Chen, Yun-Tian Shen, Ke-Xin Wang, Yi-Min Han, Chen Zhang, Xiao-Ming Yang, Bing-Qian Chen

Specialty type: Cell and tissue engineering

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade B, Grade B, Grade B, Grade B

Novelty: Grade B, Grade B, Grade B, Grade B

Creativity or Innovation: Grade B, Grade B, Grade B, Grade B

Scientific Significance: Grade A, Grade B, Grade B, Grade C

P-Reviewer: Chen JL; Qiu JY; Zhang S

Received: July 4, 2024

Revised: December 9, 2024

Accepted: January 23, 2025

Published online: February 26, 2025

Processing time: 234 Days and 22.1 Hours



Ying-Jie Wang, Ze-Hao Chen, Yun-Tian Shen, Ke-Xin Wang, Chen Zhang, Key Laboratory of Neuroregeneration of Jiangsu and Ministry of Education, Nantong University, Nantong 226000, Jiangsu Province, China

Yi-Min Han, Medical College, Nantong University, Nantong 226000, Jiangsu Province, China

Xiao-Ming Yang, Co-Innovation Center of Neuroregeneration, Jiangsu Clinical Medicine Center of Tissue Engineering and Nerve Injury Repair, Nantong University, Nantong 226000, Jiangsu Province, China

Xiao-Ming Yang, Research and Development Center for E-Learning, Ministry of Education, Beijing 100816, China

Bing-Qian Chen, Department of Orthopaedics, Changshu Hospital Affiliated to Soochow University, Changshu 215500, Jiangsu Province, China

Co-first authors: Ying-Jie Wang and Ze-Hao Chen.

Corresponding author: Bing-Qian Chen, MD, Professor, Department of Orthopaedics, Changshu Hospital Affiliated to Soochow University, No. 1 Shuyuan Street, Changshu 215500, Jiangsu Province, China. cbq0433@suda.edu.cn

Abstract

Skeletal muscle atrophy results from disruptions in the growth and metabolism of striated muscle, leading to a reduction or loss of muscle fibers. This condition not only significantly impacts patients' quality of life but also imposes substantial socioeconomic burdens. The complex molecular mechanisms driving skeletal muscle atrophy contribute to the absence of effective treatment options. Recent advances in stem cell therapy have positioned it as a promising approach for addressing this condition. This article reviews the molecular mechanisms of muscle atrophy and outlines current therapeutic strategies, focusing on mesenchymal stem cells, induced pluripotent stem cells, and their derivatives. Additionally, the challenges these stem cells face in clinical applications are discussed. A deeper understanding of the regenerative potential of various stem cells could pave the way for breakthroughs in the prevention and treatment of muscle atrophy.

Key Words: Stem cell; Muscle atrophy; Therapy; Mesenchymal stem cells; Induced pluripotent stem cells; Exosome

©The Author(s) 2025. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Muscle atrophy can exert a considerable influence on the quality of life of patients. The intricate molecular mechanisms of muscle atrophy give rise to a scarcity of effective treatment options. Recent advancements in stem cell therapy imply that it is a promising solution to this problem. This review summarizes the molecular mechanisms of muscle atrophy, presents current treatment strategies with a focus on mesenchymal stem cells, induced pluripotent stem cells, and their derivatives. Finally, it discusses the challenges encountered in the clinical application of these stem cells.

Citation: Wang YJ, Chen ZH, Shen YT, Wang KX, Han YM, Zhang C, Yang XM, Chen BQ. Stem cell therapy: A promising therapeutic approach for skeletal muscle atrophy. *World J Stem Cells* 2025; 17(2): 98693

URL: <https://www.wjgnet.com/1948-0210/full/v17/i2/98693.htm>

DOI: <https://dx.doi.org/10.4252/wjsc.v17.i2.98693>

INTRODUCTION

Muscle atrophy occurs in various physiological or pathological circumstances, including aging, disuse, dystrophy, amyotrophic lateral sclerosis (ALS), and systemic diseases such as cancer, sepsis, diabetes, chronic kidney disease and chronic obstructive pulmonary disease (Figure 1)[1-4]. Skeletal muscle atrophy causes a progressive decrease in the body's response to various stresses and chronic diseases, leading to a reduced quality of life and imposing a heavy socioeconomic burden, thus adversely impacting the prognosis of related diseases[5]. As such, the prevention and management of skeletal muscle atrophy has garnered increasing attention from researchers. Despite ongoing efforts, the pathogenesis of skeletal muscle atrophy remains incompletely understood, and aside from rehabilitation, no effective treatment options fully restore muscle function. However, with advancing research into therapeutic strategies for muscle atrophy, stem cell-based approaches have shown promising potential for preventing and treating atrophy-related disorders. The transplantation or injection of various stem cells has demonstrated efficacy in mitigating muscle atrophy and has become an integral part of the treatment strategy[6,7]. Additionally, exosomes derived from stem cells have been shown to modulate the balance between protein synthesis and degradation, further promoting muscle repair. This review explores the potential of stem cell-based therapies and their derivatives in treating muscle atrophy, offering promising prospects for clinical application.

MOLECULAR MECHANISM OF MUSCLE ATROPHY

Skeletal muscle atrophy refers to the reduction or loss of muscle fibers due to nutritional disorders or diseases, leading to decreased muscle mass. Muscle mass is primarily regulated by the balance between protein synthesis and degradation under normal physiological conditions. Disruptions in the body's microenvironment can result in an imbalance in skeletal muscle homeostasis, contributing to atrophy[8,9]. Recent findings suggest that skeletal muscle atrophy is linked to reduced protein synthesis and increased proteolysis, driven by factors such as inflammation, oxidative stress, autophagy, and apoptosis[10-12]. The key mechanisms of proteolysis include the ubiquitin-proteasome system, the autophagy-lysosome pathway, caspase-mediated proteolysis, and calpain-mediated proteolysis (Figure 2)[11,13,14]. Muscle RING-Finger protein 1 (MuRF-1) and Muscle Atrophy F-box protein are strongly associated with muscle protein degradation and serve as reliable biomarkers for this process. Moreover, the insulin-like growth factor/protein kinase B (Akt)/mechanistic target of rapamycin signaling pathway, which is critical for protein synthesis in skeletal muscle, is a major regulator of muscle atrophy[15].

Studies indicate that various triggering factors, including oxidative stress and inflammation either promote proteolytic processes or inhibit protein synthesis pathways, thereby leading to skeletal muscle atrophy[9,16-18]. Our previous research proposed a novel framework for understanding denervation-induced muscle atrophy, which involves four stages: Oxidative stress, inflammation, atrophy, and atrophic fibrosis[19]. Denervated skeletal muscles lose their contractile function, resulting in reduced blood flow to the affected muscles. This, in turn, induces a hypoxic environment that produces excess reactive oxygen species (ROS). The excessive ROS causes oxidative stress damage, triggering the release of inflammatory factors and activating inflammatory pathways, which cascade into a sustained inflammatory response. This response further accelerates muscle atrophy through reduced protein synthesis and enhanced proteolysis, ultimately leading to muscle atrophy and fibrosis[20]. Long-term oxidative stress injury and inflammation alter the microenvironment of skeletal muscle cells[21,22]. In particular, various inflammatory cytokines activate multiple signaling pathways *in vivo*, including nuclear factor- κ B, Janus kinase/signal transducer and activator of transcription (JAK/STAT), and mitogen activated protein kinase (MAPK)/p38 pathway, leading to enhanced protein proteolysis and reduced protein synthesis in skeletal muscle[11,23,24]. In addition, denervation-induced muscular atrophy is likewise




Figure 1 Different causes of muscular atrophy. ALS: Amyotrophic lateral sclerosis.




Figure 2 The mechanism underlying muscle atrophy. ROS: Reactive oxygen species; NF-κB: Nuclear factor-κB; JAK: Janus kinase; STAT: Signal transducer and activator of transcription; MAPK: Mitogen activated protein kinase; IGF: Insulin-like growth factor; Akt: Protein kinase B; mTOR: Mechanistic target of rapamycin; UPS: Ubiquitin-proteasome system; ALP: Autophagy-lysosome pathway.

associated with the diminished availability of neurotrophic factors and lowered levels of activity[25]. In conclusion, the complex molecular mechanisms of muscle atrophy have not been well understood, and still needed to be further investigated to provide potential therapeutic strategies for muscle atrophy.

CURRENT STATUS OF MUSCLE ATROPHY TREATMENT

Skeletal muscle atrophy is characterized by a significant reduction in muscle mass, resulting from the disruption of the dynamic equilibrium between protein synthesis and degradation. Given this imbalance, a series of unfavorable changes occur in muscle function and structure. As muscle tissue plays a crucial role in movement, energy metabolism, storage, circulation, and protection[26,27]. Therefore, muscle atrophy can have a significant impact on quality of life. Consequently, the prevention and treatment of muscle atrophy are critical. At present, rehabilitation therapy is the major clinical treatment for muscle atrophy, and the muscle strength is partially restored with the assistance of various rehabilitation exercises. However, rehabilitation exercises require patients to meet certain basic conditions[28]. For example, patients who have been bedridden for long time or have mobility problems due to fractures or nerve deficits are unable to undergo rehabilitation[7]. In addition, rehabilitation exercises need to be conducted according to strict standards, otherwise secondary injuries are likely to occur. Therefore, massage or acupuncture is mostly used to prevent disuse muscular atrophy in the patients who are unable to undergo rehabilitation[29,30]. Although rehabilitation therapy has minimal side effects and no obvious sequelae, it is not very effective in treating muscle atrophy in certain populations[31-33]. So that pursuing more effective treatment measures may be beneficial in patients with several of muscular disorders.

Current therapies focus on how to promote protein synthesis, inhibit protein degradation and facilitate muscle regeneration to prevent or reduce muscle atrophy[7,34]. To date, numerous experiments have shown that anti-inflam-

matory and antioxidant drugs effectively alleviate the progression of muscle atrophy and significantly improve muscle mass and function[17,35,36]. While the exact molecular mechanisms remain to be fully elucidated, treatment with these drugs has been shown to induce significant changes in blood flow, mitochondrial activity, and apoptosis[17]. Scavenging accumulated ROS, improving autophagy and protecting mitochondria can alleviate skeletal muscle atrophy[37-39]. Transforming growth factor- β 1 (TGF- β 1) is an essential regulator of skeletal muscle several physio-pathologic proceedings, such as myogenesis, regeneration, and muscle atrophy. Injection of TGF- β 1 into mice tibialis anterior induced muscle atrophy, leading to decreased fiber diameter, increased MuRF-1 levels and ROS contents. The antioxidant activity of N-acetyl-L-cysteine is used to inhibit the production of ROS in the target muscle, thereby reducing TGF- β -induced muscle atrophy[37]. Other drugs with potent antioxidant properties, such as astaxanthin, nobiletin, pyrroloquinoline quinone, isoquercitrin, and salidroside, have been shown to significantly reduce ROS levels in muscle[7,40,41]. Consequently, the mitochondrial function is stabilized and enhanced, which further reduces apoptosis caused by oxidative stress and abnormalities in mitochondria, and finally inhibits the progression of muscle atrophy[18,36,39,42]. Overall, inhibiting inflammation and reducing ROS accumulation may influence protein synthesis and degradation through multiple pathways, thereby suppressing skeletal muscle atrophy. However, most of these anti-inflammatory and antioxidant drugs have been primarily tested in pre-clinical studies, and their efficacy and safety in human applications remain to be thoroughly investigated. In recent years, stem cell therapies with superior biological properties have emerged as promising treatments for muscle atrophy. Furthermore, stem cell-based interventions have shown substantial benefits in animal models, indicating their potential for future clinical applications.

STEM CELLS AND ITS DERIVATIVES THERAPY

Stem cell transplantation offers a promising approach for treating various diseases due to the unique self-renewal capability of stem cells and their ability to differentiate into multiple lineages, including osteocytes, neurons, chondrocytes, hepatocytes, and adipocytes. This therapeutic strategy has also been applied to enhance the regenerative capacity of skeletal muscle following injury or degradation due to diseases such as muscular dystrophy[43,44]. In hereditary muscular dystrophies, such as facioscapulohumeral muscular dystrophy (FSHD), stem cell therapy has demonstrated significant improvements over non-transplanted controls, notably by reducing fibrosis and enhancing contractile-specific force[45]. Additionally, hematopoietic stem cells have also been discovered to alleviate muscle atrophy caused by Friedreich's ataxia and Pompe disease[46,47]. Transplantation of bone marrow mesenchymal stem cells (BMSCs) or injection of their secretory vesicles significantly improves symptoms of muscle atrophy with effective control of inflammation[48]. Due to their excellent properties such as proliferation and directed differentiation, stem cells have considerable promise for therapy of muscle atrophy. Some clinical trials have indicated that stem cell therapy is capable of addressing certain intractable hereditary motor neuron (MN) diseases, such as ALS and spinal muscular atrophy muscular atrophy[49-53]. Herein, we summarized the potential role of mesenchymal stem cells (MSCs), induced pluripotent stem cells (iPSCs), and stem cell derivatives as the basis for the treatment of skeletal muscle atrophy.

MSCs in the treatment of muscle atrophy

MSCs is one of the most promising multipotent stem cell types widely used in regenerative medicine. MSCs can be isolated from a variety of tissues, such as adipose tissue, bone marrow, cranial neural crest, umbilical cords, and allogeneic placenta[16,54-56]. Despite cells from different tissues showing similar surface markers, they exhibited substantial differences in cell abundance, proliferation capacity, immune regulation, and differentiation capacity[57]. It has been reported that several MSCs including BMSCs, adipose-derived stem cells (ADSCs) and umbilical cord-derived MSCs (UC-MSCs) are effective in promoting skeletal muscle regeneration in acute and chronic diseases[58-61].

BMSCs in the treatment of muscle atrophy: The application of MSCs has become an established treatment for muscle diseases[62]. MSC injection has been shown to significantly improve muscle mass parameters, including mean fiber diameter, muscle mass, and myonuclear count in the extensor digitorum longus of rats following denervation[63]. ALS, a severe and progressive neurodegenerative disease characterized by the degeneration and loss of MNs, leads to muscle atrophy and loss of motor control[2]. Intrathecal transplantation of BMSCs in ALS mouse models significantly alleviated disease symptoms, delayed disease progression, and improved motor outcomes[64,65]. Studies have demonstrated that BMSCs possess regenerative potential for skeletal muscle[63,66]. Notably, BMSC transplantation increases the number of muscle stem cells and significantly reduces muscle fibrosis. In rats with acute muscle injury, intramuscular BMSC injection downregulated the protein levels of inflammatory cytokines such as interleukin- 1β (IL- 1β), IL-6, and tumor necrosis factor- α (TNF- α), thereby promoting muscle healing. Following BMSC transplantation, damaged muscle fibers were regenerated, and impaired blood vessels were largely restored. Furthermore, fibrosis progression in injured muscles was inhibited through the suppression of TGF- β 1 downstream signaling[67]. Allogeneic BMSCs, when directly injected into acutely injured rat muscle, promoted myofibrogenesis and accelerated functional muscle recovery[68]. MSCs also enhance myotube differentiation, with co-cultures of muscle stem cells and BMSCs showing significant improvements in myotube formation[69]. BMSCs have shown therapeutic potential for dystrophic muscular atrophy as well[70]. In the mdx mouse model of Duchenne muscular dystrophy, peritoneal cavity transplantation of BMSCs significantly alleviated pathological malnutrition in the diaphragm, substantially extending lifespan[66]. Recent studies indicate that hypoxic preconditioning increases the production and secretion of stromal cell-derived factor-1 (SDF-1) and vascular endothelial growth factor in BMSCs. Transplanting hypoxic pretreated BMSCs into injured muscles improves the formation of new blood vessels and enhance muscle repair ability[71]. Thus, BMSCs have demonstrated considerable promise as a

therapeutic approach for preventing muscle atrophy and promoting regeneration (Figure 3).

ADSCs in the treatment of muscle atrophy: A study has highlighted that both allogeneic BMSCs and ADSCs exhibit significant regenerative effects in treating skeletal muscle lacerations in rats, with ADSCs showing superior outcomes [72]. Local injection of ADSCs improves muscle strength and resistance to fatigue, primarily by regulating inflammation and gene products involved in regenerative processes. One study suggested that transplantation of a Matrigel 3D matrix containing TGF β antibody-pretreated ADSCs into the injured gastrocnemius muscle of mice resulted in improved muscle mass and structure, reduced fibrosis, ameliorated the inflammatory microenvironment, and promoted skeletal muscle regeneration [73]. ADSCs can alleviate immune responses and support muscle reconstruction, with cytokine therapy further enhancing these effects. Preconditioning ADSCs with IL-4, SDF-1, or IL-11 overexpression further boosts their regenerative potential in damaged muscle tissue [74,75]. Recently, a novel approach involving the tethering of nanoparticles releasing TNF- α on the surface of ADSCs to stimulate their secretory activity has been shown to enhance perfusion, walking ability, and muscle mass recovery in ischemic hindlimb mice, providing a more efficient method for clinical stem cell therapy compared to previous pre-treatment strategies [76]. ADSCs have also demonstrated significant therapeutic potential in ALS, where they protect neurons and improve motor function by increasing neurotrophic factor expression [77,78]. These findings underscore the potential of ADSCs in treating muscle-wasting diseases (Figure 4).

UC-MSCs in the treatment of muscle atrophy: UC-MSCs are gaining traction as a promising clinical tool due to their noninvasive collection and superior proliferation capacity, alongside lower immunogenicity compared to other MSCs like BMSCs [79]. As early as 2016, Kim *et al* [80] demonstrated that conditioned medium from UC-MSCs protects muscle cells from hindlimb suspension-induced atrophy. The injection of UC-MSCs activated the expression of muscle-specific markers such as desmin and skeletal muscle actin in satellite cells, stimulating myoblast proliferation and regulating the phosphatidylinositol 3-kinase (PI3K)/Akt pathway by downregulating MuRF-1 and Muscle Atrophy F-box protein [80]. In senescence-accelerated mouse prone 10 mice, UC-MSC injection inhibited apoptosis and inflammation while improving mitochondrial biogenesis *via* the AMP-activated protein kinase (AMPK)/peroxisome proliferator-activated receptor gamma coactivator 1-alpha axis, thereby alleviating sarcopenia-related skeletal muscle atrophy [81]. Muscle atrophy is also associated with chronic metabolic and endocrine diseases such as obesity, hyperthyroidism, and type 2 diabetes mellitus [82,83]. A recent study evaluating the effects of human UC-MSCs (hUC-MSCs) on obesity- and diabetes-induced muscle atrophy found that exosomes derived from hUC-MSCs alleviated muscle atrophy symptoms by enhancing AMPK/UNC-51-like kinases 1 (ULK1)-mediated autophagy [84]. Therefore, UC-MSCs represent a promising therapeutic option for muscle atrophy prevention and regeneration (Figure 5).

In summary, MSCs from various sources offer effective strategies for mitigating muscle atrophy symptoms and maintaining muscle homeostasis by regulating local inflammation, promoting muscle regeneration, and protecting nerve innervation. These findings demonstrate the significant therapeutic potential of MSCs in treating muscle atrophy.

iPSCs in the treatment of muscle atrophy

Pluripotent stem cells (PSCs), including embryonic stem cells (ESCs) and iPSCs, are actively explored for applications in regenerative medicine, drug screening, and disease modeling [85]. ESCs, derived from early embryos at the blastocyst stage, are pluripotent and capable of differentiating into all cell types within an organism [86]. However, due to ethical concerns, challenges in generating high-purity lineage-specific cell lines, and the potential risk of tumorigenesis, ESCs are seldom used in direct disease treatment and are primarily employed in the *in vitro* cultivation of tissues and organs [87]. In contrast, iPSCs offer distinct advantages, sharing key properties with ESCs, such as differentiation potential, electrophysiological characteristics, biochemistry, and telomerase activity [88]. Compared to muscle cells or MSCs directly transplanted into patients, iPSCs are more abundant, can be easily isolated and purified using specific markers, and demonstrate robust differentiation capabilities [89-91]. Furthermore, iPSCs mitigate the risk of immune rejection, as they avoid allogeneic transplantation issues [92,93].

Differentiation of human iPSCs into myoblasts presents a promising approach for both autologous and allogeneic therapies in the treatment of muscle atrophy [91,94,95]. Human PSC-derived myogenic progenitor cells (MPCs), generated using both transgene-free and transgene-dependent protocols, have demonstrated potential in skeletal muscle engraftment [95]. Using a fluorescent reporter system, studies have shown that these MPCs promote myofiber regeneration in both local injury models and mdx mice [96]. PSC-derived PAX7⁺ MPCs represent a compelling strategy for muscle regeneration, enabling large-scale production of human myogenic progenitors [97]. Furthermore, under optimized 3D culture conditions, PSC-derived PAX7⁺ MPCs are capable of regenerating functional skeletal muscle [98]. In immunocompromised NSG mice, iSKM bundles derived from these cells were shown to survive, gradually vascularize, and maintain muscle function when implanted into the hind-limb muscles. Recent studies have directed human PSCs toward the differentiation of embryonic muscle progenitors, which subsequently developed into primary and secondary embryonic muscle, eventually forming three-dimensional muscle structures [99]. These tissue-engineered muscles exhibited high levels of maturation and function that closely resemble normal human muscle. After skeletal muscle atrophy, recovery is limited, even with sufficient blood supply and appropriate rest. In most cases, fibroblasts proliferate to repair damaged muscle tissue, leading to muscle mass reduction and impaired mobility. FSHD, a genetically dominant progressive myopathy characterized by muscle fibrosis, atrophy, and fat replacement, is one such condition. In an FSHD mouse model, the transplantation of MPCs derived from PSCs significantly improved muscle contractility compared to non-transplanted controls [45]. Moreover, the transplantation of human iPSC-derived MNs has shown potential in treating neuromuscular atrophy in mice with sciatic nerve injury [100]. Regeneration of neuromuscular junctions is vital for functional recovery in denervated muscles affected by trauma or MN diseases. P75⁺ neural stem cells derived from human iPSCs have been shown to enhance the secretion of soluble factors that promote neuromuscular regeneration.




Figure 3 The potential role of bone marrow mesenchymal stem cells for the treatment of skeletal muscle atrophy. BMSCs: Bone marrow mesenchymal stem cells.




Figure 4 The potential role of adipose-derived stem cells for the treatment of skeletal muscle atrophy. ADSCs: Adipose-derived stem cells; TNF: Tumor necrosis factor.

When injected into denervated gastrocnemius muscle, these NCSC spheres significantly improved neuromuscular function recovery[101]. iPSCs also show promise in treating ALS. In an ALS mouse model, transplantation of neural stem cells derived from iPSCs improved the microenvironment by secreting growth factors and immunomodulatory molecules, which likely contributed to the reconstruction of neural circuits and restoration of motor control in ALS-affected muscles[102,103]. Therefore, iPSCs have the potential to treat muscle atrophy and provide a new technical idea for personalized treatment of dystrophic muscular atrophy in the future (Figure 6). The available results have shown that this method is feasible and has the potential for application and development in clinical practice.

Exosomes derived from stem cells in the treatment of muscle atrophy

The regenerative potential of MSCs stems not only from their direct transplantation into target tissues and subsequent differentiation but also from their secretions, particularly exosomes, which activate the regenerative capacity of resident cells. Exosomes contain various bioactive components, including growth factors, cytokines, mRNAs, signaling lipids, and regulatory microRNAs[104]. Although MSCs from different tissues and donors secrete distinct protein profiles, their actions *in vitro* can vary; they are nonetheless capable of regulating apoptosis in dystrophic myogenic cells and promoting cell migration and proliferation[105]. Skeletal muscle regeneration is closely linked to the biological factors secreted by exosomes, which exert significant bioactive effects on damaged muscle tissue[106,107]. Increasing evidence suggests that exosomes derived from stem cells offer promising therapeutic potential for muscle regeneration (Table 1).

Table 1 The potential role of mesenchymal stem cell-derived exosomes in the treatment of skeletal muscle atrophy

Resource	Mechanism	Function	Ref.
UC-MSCs	Releasing circHIPK3, while circHIPK3 down-regulates miR-421, leading to increased expression of FOXO3a, thereby inhibiting IL-1 β and IL-18	Accelerate repair of ischemic muscle injury in mice	[108]
	hUC-MSCs or hUC-MSC-derived exosomes induce the activation of AMPK/ULK1 signal and enhancement of autophagy	Alleviate muscle atrophy induced by diabetes and obesity in mice	[84]
	Down-regulating IL-6 and IL-1 β , up-regulating IL-10, and regulating inflammation of injured nerves	Promote the recovery of motor function and regeneration of axons, alleviated denervated gastrocnemius atrophy in rats	[109]
	Regulating the miR-132-3p/FoxO3 axis	Improve age-related muscle atrophy in a mouse model	[110]
ADSCs	Promoting peripheral nerve regeneration <i>via</i> optimizing SC function	Alleviate denervated muscle atrophy in rats	[111]
	Inhibiting inflammation, enhancing proliferation of muscle satellite cells, resisting cell senescence and promoting angiogenesis	Maintain muscle homeostasis, promote muscle regeneration in a mouse model of cardiotoxin-induced injury	[112,113]
	Preventing the atrophy, fatty infiltration, inflammation, and vascularization of muscles	Decrease atrophy and degeneration associated with torn rotator cuffs, and improve muscle regeneration in rats	[114]
	Improving motor performance; protecting lumbar motoneurons, neuromuscular junction, and muscle, decreasing glial cells activation	Protect neuromuscular junction and muscle in SOD1 (G93A) murine model	[115]
BMSCs	Promoting M2 macrophages polarization, reducing the production of inflammatory cytokines, upregulating anti-inflammatory factors expression	Attenuate muscle injury and promote muscle healing in mice with hit injury	[116]
	Foxo1 expression was inhibited by up-regulating the miR486-5p/Foxo1 axis	Inhibit dexamethasone-induced muscle atrophy in mice	[117]
	M2 macrophage polarization is regulated by NF- κ B pathway to reduce inflammatory pain, and myeloid axon regeneration is enhanced by MEK/ERK pathway	Improve denervated muscle atrophy and further promote functional recovery in rats	[106,107]

UC-MSCs: Umbilical cord-derived mesenchymal stromal cells; FOXO3a: Forkhead box O3a; IL: Interleukin; hUC-MSCs: Human umbilical cord-derived mesenchymal stromal cells; AMPK: AMP-activated protein kinase; ULK1: UNC-51-like kinases 1; ADSCs: Adipose-derived stem cells; SC: Schwann cell; BMSCs: Bone marrow mesenchymal stem cells; NF- κ B: Nuclear factor- κ B; MEK: Mitogen-activated protein kinase; ERK: Extracellular signal-regulated kinase.

Exosomes secreted by hUC-MSCs (hUC-MSC-Exos) have been confirmed to be effective against muscle injury. In a mouse model of ischemic induced skeletal muscle injury, purified hUC-MSC-Exos accelerates skeletal muscle regeneration by releasing circHIPK3[108]. This led to the downregulation of miR-421, resulting in Forkhead box O3a (FOXO3a) expression, inhibition of pyroptosis, and decreased release of IL-1 β and IL-18[108]. Similarly, Song *et al*[84] reported that hUC-MSC-Exos alleviated diabetes-associated muscle atrophy by enhancing autophagy through the AMPK/ULK1 axis. In a rat sciatic nerve defect model, hUC-MSC-Exos promoted nerve regeneration and attenuated gastrocnemius muscle atrophy by modulating inflammation - downregulating IL-1 β and IL-6, upregulating IL-10[109]. A recent study suggest that hUC-MSC-Exos may improve age-related and dexamethasone-induced muscle atrophy in a mouse model by regulating the miR-132-3p/FoxO3 axis[110]. These studies suggest that the application of exosomes derived from UC-MSCs have great potential for muscle atrophy.

The protective effect of exosomes secreted by human ADSCs on denervated muscle atrophy has been confirmed in rat studies, where exosomes secreted by human ADSCs significantly promoted nerve regeneration and alleviated muscle atrophy following sciatic nerve dissection[111]. Exosomes secreted by ADSCs (ADSC-Exos) enhanced Schwann cell proliferation, migration, and myelination, while also stimulating the release of neurotrophic factors through the upregulation of key genes. Furthermore, ADSC-Exos contributed to muscle homeostasis by inhibiting inflammation, promoting muscle satellite cell proliferation, resisting cellular senescence, and stimulating angiogenesis[112-114]. ADSC-Exos play a critical role in skeletal muscle regeneration at various stages by modulating key signaling pathways, including MAPK, Wnt, PI3K/Akt, and JAK/STAT[113]. For example, catenin derived from ADSC-Exos activates the classical Wnt pathway, supporting myogenic maintenance of satellite cells, while also coordinating their self-renewal and migration *via* the non-classical Wnt pathway[115]. In the early stages of muscle regeneration, ADSC-Exos activate MAPK/extracellular signal-regulated kinase 1/2 and MAPK/JNK pathways, promoting myoblast proliferation and preventing premature differentiation[116]. At later stages, the p38/MK2 and MEF2 pathways drive myogenic differentiation by enhancing MyoD transcriptional activity[117], while the MAPK/extracellular signal-regulated kinase 5 pathway facilitates multinucleated myotube formation. Additionally, ADSC-Exos influence the PI3K/Akt pathway to upregulate protein synthesis and




Figure 5 The potential role of umbilical cord-derived mesenchymal stromal cells for the treatment of skeletal muscle atrophy. UC-MSCs: Umbilical cord-derived mesenchymal stromal cells.




Figure 6 The potential role of induced pluripotent stem cells for the treatment of skeletal muscle atrophy. iPSCs: Induced pluripotent stem cells.

suppress protein degradation, counteracting muscle atrophy. However, the potential inhibitory effect of PI3K/Akt on autophagy *via* mechanistic target of rapamycin and FOXO suggests that its role in muscle regeneration may not be entirely beneficial[118]. Moreover, ADSC-Exos regulate the JAK1/STAT1/3 pathway to prevent premature differentiation of myoblasts, while JAK2/STAT2/3 and JAK3/STAT1/3 pathways promote myogenic differentiation. ADSC-Exos have also been shown to reduce muscle-related protein degeneration and muscle atrophy while enhancing regeneration and biomechanical properties in torn rotator cuff muscles in rats[119]. Additionally, repeated administration of ADSC-Exos reduced glial cell activation, protected muscles, MNs, and neuromuscular junctions, and improved motor performance in SOD1-G93A mice[120]. Proteomic analysis of ADSC-Exos revealed their influence on ALS-related pathways, including oxidative stress response, cell apoptosis, cell adhesion, and PI3K/Akt signaling[121]. In conclusion, extensive evidence highlights the significant potential of ADSC-Exos in the treatment of muscle diseases.

Exosomes secreted by BMSCs (BMSC-Exos) have shown therapeutic potential for muscle injury and atrophy. Intramuscular injection of BMSC-Exos reduces inflammation following muscle contusion in mice, mitigates muscle fibrosis, promotes regeneration, and enhances muscle biomechanical properties[122]. A study also demonstrated that BMSC-Exos counteract dexamethasone-induced muscle atrophy through the miR486-5p/Foxo1 pathway[123]. Recently, a BMSC-Exos-loaded electroconductive nerve dressing has been developed for treating diabetic peripheral nerve injury. Implantation of the BMSC-Exos-loaded electroconductive nerve dressing complex around damaged sciatic nerves enhances myelinated axon regeneration, ameliorates gastrocnemius muscle atrophy, and promotes functional recovery in diabetic rats[107]. Furthermore, supplementation with exosomes secreted by iPSCs in acellular nerve grafts through optimized chemical extraction processes has been shown to repair long-distance peripheral nerve defects, accelerating motor recovery and reversing denervated muscle atrophy[124]. Skin-derived precursors (SKPs), pluripotent cells derived from the dermis, are another promising stem cell source. Artificial nerve grafts incorporating SKP-derived Schwann cells can alleviate denervation-induced muscle atrophy[125]. In our recent study, direct intramuscular injection of SKP-SC-

extracellular vesicles alleviated blood flow reduction, oxidative stress, and inflammation in target muscles, thus reducing skeletal muscle atrophy caused by denervation[126]. Overall, stem cell-derived exosomes mitigate inflammation, reduce fibrosis, accelerate muscle regeneration, and promote functional recovery[122,127]. In conclusion, stem cell-derived exosomes present substantial promise for clinical applications, with exosomes secreted by MSCs offering a potential alternative to whole-cell regenerative therapies.

CHALLENGES OF STEM CELL THERAPY

Despite the considerable promise of stem cells in treating muscle atrophy and related diseases, challenges surrounding their source and biological characteristics, particularly ethical concerns related to the use of human ESCs, remain significant. Furthermore, other types of stem cells present notable limitations that impede the advancement of stem cell therapies. The potential risks of immune rejection and tumor formation after transplantation are significant issues worthy of attention. The approaches to mitigate these risks include strategies such as immunosuppression, immune matching, and stem cell genetic engineering. However, each strategy confronts its own unique challenges and may have varying degrees of impact on patient safety[128].

MSCs are widely utilized in medicine due to their pluripotency, broad tissue origins, and minimal ethical concerns. While MSC-based therapies offer advantages over conventional treatments in clinical trials, their therapeutic outcomes often fall short of expectations. Key challenges, including stem cell homing, immune modulation, and other factors, need to be addressed for successful MSC application. Under various conditions, such as ischemia or injury, MSCs - whether autologous or exogenous - migrate directionally from the vascular endothelium to the affected tissues, where they colonize and survive[129,130]. The activation of MSCs occurs in response to elevated inflammatory factors like IL-1 and TNF- α , which generate a concentration gradient of ligands around the target tissue, promoting MSC migration towards ischemic and hypoxic regions[131,132]. SDF-1 and its receptor C-X-C motif chemokine receptor 4 are the most extensively studied cytokines involved in cell homing[126]. SDF-1 is upregulated at injury sites, acting on MSC migration in a dose-dependent manner, thus enhancing BMSC homing for tissue repair[133,134]. However, maintaining MSCs at specific target sites, such as muscles, remains a significant challenge. MSCs secrete paracrine factors capable of tissue remodeling and repair, act as drug carriers, and modulate immune responses[135-138]. MSCs can detect injury sites and elicit stronger immune responses in low-immune states, while inhibiting immune cell function when inflammation is excessive in the damaged area[139]. They exhibit dual, opposing functions - pro-inflammatory or anti-inflammatory - depending on the inflammatory factors present in the local environment[66,140]. The key to optimizing their therapeutic effects lies in understanding how to modulate or exploit the immune-inducing properties of MSCs. Additionally, factors contributing to suboptimal therapeutic outcomes include the age of MSC donors, variations in cell isolation and culture protocols, the method of administration, and individual differences among recipients[141-143]. Long-term culture impairs the differentiation potential of MSCs, and the same cell batch may exhibit heterogeneous differentiation, which can skew experimental results. Moreover, various biological and experimental factors introduce safety risks in clinical and basic research applications of stem cells, such as tumor formation, infection, necrosis from tissue blockage due to stem cell homing[144], and other exogenous risks[145,146]. Although MSCs have good therapeutic effects in animal models of muscle diseases, biosafety evaluation standards for stem cell experiments and therapy are lacking. The heterogeneity of MSCs is one of the factors contributing to the variation in treatment outcomes, which presents challenges in determining the optimal stem cell therapy protocol, dosage, and timing[57]. These issues necessitate further research and clinical practice for resolution.

There was no unified and effective differentiation method for iPSCs to differentiate into muscle precursor cells, and their purification protocols are also inconsistent[95,147]. For transgene-free protocols, due to discrepancy in purification and transplantation times among the protocols, none of previously reported optimal purification strategies have been reproducibly validated. For transgene-dependent protocols, a risk factor that may not be ignored is the induced expression of proto-oncogenes introduced *via* viral vectors or existing in the host cells. It remains unclear whether other unknown genes are involved in the reprogramming process or if genetic and epigenetic changes occur during this process [148]. Consequently, several challenges remain before iPSCs can be reliably applied in clinical settings.

In the current social, scientific and ethical context, the use of stem cell derivatives as a cell-free therapy is rapidly emerging as an alternative with additional advantages. It can avoid safety issues related to the use of living cells, ethical issues related to the origin of the cells, and immunocompatibility issues. MSC-derived exosomes, which are characterized by low-immunogenicity, high biocompatibility, safety and low toxicity, provide a new approach to deliver therapeutic agents to target cells, which is the best alternative to stem cell therapy[149-151]. Although exosomes are generally considered to have great clinical therapeutic value, there are still many drawbacks such as extremely low concentration of active ingredients, poor specificity, and difficulty in traceability, which restrict the clinical use of exosomes. It is also worthy of note that cell-free therapeutic products might take a considerable amount of time for administration and could result in drug resistance as well as some adverse reactions[152].

It is crucial to prove the safety and efficacy of stem cells and their derived exosomes therapy through clinical trials. Evaluating long-term efficacy means tracking changes in cell survival, integration, and treatment response over time, and identifying any adverse events or delayed complications. This requires establishing standardized methods and guidelines [128]. Also, currently, there's no clear consensus or standard about the administration route, dose, and treatment time [57]. These require the establishment of standardized methods and guidelines. Of course, in addition to the ethical controversy related to the use of ESCs mentioned earlier, issues such as informed consent from patients, the experimental nature of the treatment, and fair access to emerging therapies should also be considered in the consideration of stem cell clinical use. Solving these multifaceted challenges requires not only the efforts of researchers and clinicians, but also

coordinated efforts with regulatory agencies and ethics committees, which will be key to realizing the potential of stem cell therapy for muscle atrophy.

CONCLUSION

The rapid advancements in stem cell research have opened new avenues for treating previously incurable diseases. Numerous clinical trials have demonstrated promising outcomes in addressing cardiovascular, neurological, and immune disorders[151,153-155]. This review systematically analyzes the therapeutic application of stem cells including BMSCs, ADSCs, UCMSCs, and iPSCs in skeletal muscle atrophy due to various causes. In addition, we also summarize the advantages of existing therapeutic strategies and further research development directions in order to mitigate or eliminate therapeutic deficiencies and establish a relatively sound therapeutic evaluation system. Cell-free therapies based on stem cell derivatives have better prospects compared to conventional cell transplantation therapy[156,157]. A full understanding of the capabilities of different types of stem cells, especially their derivatives, in muscle repair and regeneration would be a breakthrough in the treatment of muscle atrophy.

ACKNOWLEDGEMENTS

We would like to thank all the professionals who contributed to the discussion and elaboration of this review.

FOOTNOTES

Author contributions: Wang YJ and Chen ZH contributed equally to write the paper and are co-first authors of this manuscript; Shen YT modified the manuscript; Wang KX, Han YM, Zhang C, and Yang XM provided data; Chen BQ designed the review and were responsible for the final proofreading; and all authors have read and approved the final manuscript.

Supported by Suzhou Science and Technology Development Planning Project, No. SYW2024048; National Natural Science Foundation of China, No. 81901933; and Major Natural Science Research Projects in Universities of Jiangsu Province, No. 24KJA310007.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: China

ORCID number: Yun-Tian Shen 0000-0002-5145-4761; Bing-Qian Chen 0000-0002-4012-3481.

S-Editor: Wang JJ

L-Editor: A

P-Editor: Zhang XD

REFERENCES

- 1 **Sartori R**, Romanello V, Sandri M. Mechanisms of muscle atrophy and hypertrophy: implications in health and disease. *Nat Commun* 2021; **12**: 330 [PMID: 33436614 DOI: 10.1038/s41467-020-20123-1]
- 2 **Wang K**, Liu Q, Tang M, Qi G, Qiu C, Huang Y, Yu W, Wang W, Sun H, Ni X, Shen Y, Fang X. Chronic kidney disease-induced muscle atrophy: Molecular mechanisms and promising therapies. *Biochem Pharmacol* 2023; **208**: 115407 [PMID: 36596414 DOI: 10.1016/j.bcp.2022.115407]
- 3 **Shen Y**, Li M, Wang K, Qi G, Liu H, Wang W, Ji Y, Chang M, Deng C, Xu F, Shen M, Sun H. Diabetic Muscular Atrophy: Molecular Mechanisms and Promising Therapies. *Front Endocrinol (Lausanne)* 2022; **13**: 917113 [PMID: 35846289 DOI: 10.3389/fendo.2022.917113]
- 4 **Chang M**, Cai Y, Gao Z, Chen X, Liu B, Zhang C, Yu W, Cao Q, Shen Y, Yao X, Chen X, Sun H. Duchenne muscular dystrophy: pathogenesis and promising therapies. *J Neurol* 2023; **270**: 3733-3749 [PMID: 37258941 DOI: 10.1007/s00415-023-11796-x]
- 5 **Yin L**, Li N, Jia W, Wang N, Liang M, Yang X, Du G. Skeletal muscle atrophy: From mechanisms to treatments. *Pharmacol Res* 2021; **172**: 105807 [PMID: 34389456 DOI: 10.1016/j.phrs.2021.105807]
- 6 **Qazi TH**, Duda GN, Ort MJ, Perka C, Geissler S, Winkler T. Cell therapy to improve regeneration of skeletal muscle injuries. *J Cachexia Sarcopenia Muscle* 2019; **10**: 501-516 [PMID: 30843380 DOI: 10.1002/jcsm.12416]
- 7 **Huang L**, Li M, Deng C, Qiu J, Wang K, Chang M, Zhou S, Gu Y, Shen Y, Wang W, Huang Z, Sun H. Potential Therapeutic Strategies for Skeletal Muscle Atrophy. *Antioxidants (Basel)* 2022; **12**: 44 [PMID: 36670909 DOI: 10.3390/antiox12010044]
- 8 **You Z**, Huang X, Xiang Y, Dai J, Jiang J, Xu J. Molecular feature of neutrophils in immune microenvironment of muscle atrophy. *J Cell Mol*

- Med* 2022; **26**: 4658-4665 [PMID: 35899367 DOI: 10.1111/jcmm.17495]
- 9 **Sun H**, Sun J, Li M, Qian L, Zhang L, Huang Z, Shen Y, Law BY, Liu L, Gu X. Transcriptome Analysis of Immune Receptor Activation and Energy Metabolism Reduction as the Underlying Mechanisms in Interleukin-6-Induced Skeletal Muscle Atrophy. *Front Immunol* 2021; **12**: 730070 [PMID: 34552592 DOI: 10.3389/fimmu.2021.730070]
 - 10 **Zhang H**, Qi G, Wang K, Yang J, Shen Y, Yang X, Chen X, Yao X, Gu X, Qi L, Zhou C, Sun H. Oxidative stress: Roles in skeletal muscle atrophy. *Biochem Pharmacol* 2023; **214**: 115664 [PMID: 37331636 DOI: 10.1016/j.bcp.2023.115664]
 - 11 **Ji Y**, Li M, Chang M, Liu R, Qiu J, Wang K, Deng C, Shen Y, Zhu J, Wang W, Xu L, Sun H. Inflammation: Roles in Skeletal Muscle Atrophy. *Antioxidants (Basel)* 2022; **11**: 1686 [PMID: 36139760 DOI: 10.3390/antiox11091686]
 - 12 **Liang W**, Xu F, Li L, Peng C, Sun H, Qiu J, Sun J. Epigenetic control of skeletal muscle atrophy. *Cell Mol Biol Lett* 2024; **29**: 99 [PMID: 38978023 DOI: 10.1186/s11658-024-00618-1]
 - 13 **Huang Z**, Zhong L, Zhu J, Xu H, Ma W, Zhang L, Shen Y, Law BY, Ding F, Gu X, Sun H. Inhibition of IL-6/JAK/STAT3 pathway rescues denervation-induced skeletal muscle atrophy. *Ann Transl Med* 2020; **8**: 1681 [PMID: 33490193 DOI: 10.21037/atm-20-7269]
 - 14 **He Q**, Qiu J, Dai M, Fang Q, Sun X, Gong Y, Ding F, Sun H. MicroRNA-351 inhibits denervation-induced muscle atrophy by targeting TRAF6. *Exp Ther Med* 2016; **12**: 4029-4034 [PMID: 28101181 DOI: 10.3892/etm.2016.3856]
 - 15 **Ito N**, Ruegg UT, Takeda S. ATP-Induced Increase in Intracellular Calcium Levels and Subsequent Activation of mTOR as Regulators of Skeletal Muscle Hypertrophy. *Int J Mol Sci* 2018; **19**: 2804 [PMID: 30231482 DOI: 10.3390/ijms19092804]
 - 16 **Yan Y**, Li M, Lin J, Ji Y, Wang K, Yan D, Shen Y, Wang W, Huang Z, Jiang H, Sun H, Qi L. Adenosine monophosphate activated protein kinase contributes to skeletal muscle health through the control of mitochondrial function. *Front Pharmacol* 2022; **13**: 947387 [PMID: 36339617 DOI: 10.3389/fphar.2022.947387]
 - 17 **Zhang L**, Li M, Wang W, Yu W, Liu H, Wang K, Chang M, Deng C, Ji Y, Shen Y, Qi L, Sun H. Celecoxib alleviates denervation-induced muscle atrophy by suppressing inflammation and oxidative stress and improving microcirculation. *Biochem Pharmacol* 2022; **203**: 115186 [PMID: 35882305 DOI: 10.1016/j.bcp.2022.115186]
 - 18 **Shen Y**, Zhang Q, Huang Z, Zhu J, Qiu J, Ma W, Yang X, Ding F, Sun H. Isoquercitrin Delays Denervated Soleus Muscle Atrophy by Inhibiting Oxidative Stress and Inflammation. *Front Physiol* 2020; **11**: 988 [PMID: 32903465 DOI: 10.3389/fphys.2020.00988]
 - 19 **Shen Y**, Zhang R, Xu L, Wan Q, Zhu J, Gu J, Huang Z, Ma W, Shen M, Ding F, Sun H. Microarray Analysis of Gene Expression Provides New Insights Into Denervation-Induced Skeletal Muscle Atrophy. *Front Physiol* 2019; **10**: 1298 [PMID: 31681010 DOI: 10.3389/fphys.2019.01298]
 - 20 **Wan Q**, Zhang L, Huang Z, Zhang H, Gu J, Xu H, Yang X, Shen Y, Law BY, Zhu J, Sun H. Aspirin alleviates denervation-induced muscle atrophy via regulating the Sirt1/PGC-1 α axis and STAT3 signaling. *Ann Transl Med* 2020; **8**: 1524 [PMID: 33313269 DOI: 10.21037/atm-20-5460]
 - 21 **Sun J**, Yang H, Yang X, Chen X, Xu H, Shen Y, Ding F, Gu X, Zhu J, Sun H. Global alternative splicing landscape of skeletal muscle atrophy induced by hindlimb unloading. *Ann Transl Med* 2021; **9**: 643 [PMID: 33987341 DOI: 10.21037/atm-20-5388]
 - 22 **Ma W**, Cai Y, Shen Y, Chen X, Zhang L, Ji Y, Chen Z, Zhu J, Yang X, Sun H. HDAC4 Knockdown Alleviates Denervation-Induced Muscle Atrophy by Inhibiting Myogenin-Dependent AtroGene Activation. *Front Cell Neurosci* 2021; **15**: 663384 [PMID: 34276308 DOI: 10.3389/fncel.2021.663384]
 - 23 **Liang Z**, Zhang T, Liu H, Li Z, Peng L, Wang C, Wang T. Inflammaging: The ground for sarcopenia? *Exp Gerontol* 2022; **168**: 111931 [PMID: 35985553 DOI: 10.1016/j.exger.2022.111931]
 - 24 **Irazoki A**, Martinez-Vicente M, Aparicio P, Aris C, Alibakhshi E, Rubio-Valera M, Castellanos J, Lores L, Palacín M, Gumà A, Zorzano A, Sebastián D. Coordination of mitochondrial and lysosomal homeostasis mitigates inflammation and muscle atrophy during aging. *Aging Cell* 2022; **21**: e13583 [PMID: 35263007 DOI: 10.1111/acel.13583]
 - 25 **Chu XL**, Song XZ, Li Q, Li YR, He F, Gu XS, Ming D. Basic mechanisms of peripheral nerve injury and treatment via electrical stimulation. *Neural Regen Res* 2022; **17**: 2185-2193 [PMID: 35259827 DOI: 10.4103/1673-5374.335823]
 - 26 **Grgic J**, Schoenfeld BJ, Mikulic P. Effects of plyometric vs. resistance training on skeletal muscle hypertrophy: A review. *J Sport Health Sci* 2021; **10**: 530-536 [PMID: 32579911 DOI: 10.1016/j.jshs.2020.06.010]
 - 27 **Chen X**, Ji Y, Liu R, Zhu X, Wang K, Yang X, Liu B, Gao Z, Huang Y, Shen Y, Liu H, Sun H. Mitochondrial dysfunction: roles in skeletal muscle atrophy. *J Transl Med* 2023; **21**: 503 [PMID: 37495991 DOI: 10.1186/s12967-023-04369-z]
 - 28 **Howard EE**, Pasiakos SM, Fussell MA, Rodriguez NR. Skeletal Muscle Disuse Atrophy and the Rehabilitative Role of Protein in Recovery from Musculoskeletal Injury. *Adv Nutr* 2020; **11**: 989-1001 [PMID: 32167129 DOI: 10.1093/advances/nmaa015]
 - 29 **Lawrence MM**, Van Pelt DW, Confides AL, Hunt ER, Hettinger ZR, Laurin JL, Reid JJ, Peeler FF 3rd, Butterfield TA, Dupont-Versteegden EE, Miller BF. Massage as a mechanotherapy promotes skeletal muscle protein and ribosomal turnover but does not mitigate muscle atrophy during disuse in adult rats. *Acta Physiol (Oxf)* 2020; **229**: e13460 [PMID: 32125770 DOI: 10.1111/apha.13460]
 - 30 **Su Z**, Hu L, Cheng J, Klein JD, Hassounah F, Cai H, Li M, Wang H, Wang XH. Acupuncture plus low-frequency electrical stimulation (Acu-LFES) attenuates denervation-induced muscle atrophy. *J Appl Physiol (1985)* 2016; **120**: 426-436 [PMID: 26679610 DOI: 10.1152/jappphysiol.00175.2015]
 - 31 **Lv T**, Mo Y, Yu T, Zhang Y, Shao S, Luo Y, Shen Y, Lu M, Wong SG. An Investigation into the Rehabilitative Mechanism of Tuina in the Treatment of Sciatic Nerve Injury. *Evid Based Complement Alternat Med* 2020; **2020**: 5859298 [PMID: 32724326 DOI: 10.1155/2020/5859298]
 - 32 **Peng S**, Tian Y, Chang W, Yang Y, Li S, Ni J, Zhu W. Current state of research on acupuncture for the treatment of amyotrophic lateral sclerosis: A scoping review. *Front Neurol* 2022; **13**: 1019156 [PMID: 36408510 DOI: 10.3389/fneur.2022.1019156]
 - 33 **Rahiminezhad E**, Sadeghi M, Ahmadinejad M, Mirzadi Gohari SI, Dehghan M. A randomized controlled clinical trial of the effects of range of motion exercises and massage on muscle strength in critically ill patients. *BMC Sports Sci Med Rehabil* 2022; **14**: 96 [PMID: 35619171 DOI: 10.1186/s13102-022-00489-z]
 - 34 **Huang Z**, Zhu J, Ma W, Sun H. Strategies and potential therapeutic agents to counter skeletal muscle atrophy. *Biotarget* 2018; **2**: 8 [DOI: 10.21037/biotarget.2018.05.02]
 - 35 **Qiu J**, Fang Q, Xu T, Wu C, Xu L, Wang L, Yang X, Yu S, Zhang Q, Ding F, Sun H. Mechanistic Role of Reactive Oxygen Species and Therapeutic Potential of Antioxidants in Denervation- or Fasting-Induced Skeletal Muscle Atrophy. *Front Physiol* 2018; **9**: 215 [PMID: 29593571 DOI: 10.3389/fphys.2018.00215]
 - 36 **Xu T**, Yang X, Wu C, Qiu J, Fang Q, Wang L, Yu S, Sun H. Pyrroloquinoline quinone attenuates cachexia-induced muscle atrophy via suppression of reactive oxygen species. *J Thorac Dis* 2018; **10**: 2752-2759 [PMID: 29997937 DOI: 10.21037/jtd.2018.04.112]

- 37 **Abrigo J**, Rivera JC, Simon F, Cabrera D, Cabello-Verrugio C. Transforming growth factor type beta (TGF- β) requires reactive oxygen species to induce skeletal muscle atrophy. *Cell Signal* 2016; **28**: 366-376 [PMID: 26825874 DOI: 10.1016/j.celsig.2016.01.010]
- 38 **Scalabrin M**, Engman V, Maccannell A, Critchlow A, Roberts LD, Yuldasheva N, Bowen TS. Temporal analysis of skeletal muscle remodeling post hindlimb ischemia reveals intricate autophagy regulation. *Am J Physiol Cell Physiol* 2022; **323**: C1601-C1610 [PMID: 36252128 DOI: 10.1152/ajpcell.00174.2022]
- 39 **Wang HH**, Sun YN, Qu TQ, Sang XQ, Zhou LM, Li YX, Ren FZ. Nobiletin Prevents D-Galactose-Induced C2C12 Cell Aging by Improving Mitochondrial Function. *Int J Mol Sci* 2022; **23**: 11963 [PMID: 36233264 DOI: 10.3390/ijms231911963]
- 40 **Huang Z**, Fang Q, Ma W, Zhang Q, Qiu J, Gu X, Yang H, Sun H. Skeletal Muscle Atrophy Was Alleviated by Salidroside Through Suppressing Oxidative Stress and Inflammation During Denervation. *Front Pharmacol* 2019; **10**: 997 [PMID: 31616291 DOI: 10.3389/fphar.2019.00997]
- 41 **Wu C**, Tang L, Ni X, Xu T, Fang Q, Xu L, Ma W, Yang X, Sun H. Salidroside Attenuates Denervation-Induced Skeletal Muscle Atrophy Through Negative Regulation of Pro-inflammatory Cytokine. *Front Physiol* 2019; **10**: 665 [PMID: 31293430 DOI: 10.3389/fphys.2019.00665]
- 42 **Yang X**, Ji Y, Wang W, Zhang L, Chen Z, Yu M, Shen Y, Ding F, Gu X, Sun H. Amyotrophic Lateral Sclerosis: Molecular Mechanisms, Biomarkers, and Therapeutic Strategies. *Antioxidants (Basel)* 2021; **10**: 1012 [PMID: 34202494 DOI: 10.3390/antiox10071012]
- 43 **Salles J**, Chanet A, Guillet C, Vaes AM, Brouwer-Brolsma EM, Rocher C, Giraudet C, Patrac V, Meugnier E, Montaurier C, Denis P, Le Bacquer O, Blot A, Jourdan M, Luiking Y, Furber M, Van Dijk M, Tardif N, Yves Boirie Y, Walrand S. Vitamin D status modulates mitochondrial oxidative capacities in skeletal muscle: role in sarcopenia. *Commun Biol* 2022; **5**: 1288 [PMID: 36434267 DOI: 10.1038/s42003-022-04246-3]
- 44 **Saeki S**, Tokutake K, Takasu M, Kurimoto S, Asami Y, Onaka K, Saeki M, Hirata H. Functional Reconstruction of Denervated Muscle by Xenotransplantation of Neural Cells from Porcine to Rat. *Int J Mol Sci* 2022; **23**: 8773 [PMID: 35955906 DOI: 10.3390/ijms23158773]
- 45 **Azzag K**, Bosnakovski D, Tungtur S, Salama P, Kyba M, Perlingeiro RCR. Transplantation of PSC-derived myogenic progenitors counteracts disease phenotypes in FSHD mice. *NPJ Regen Med* 2022; **7**: 43 [PMID: 36056021 DOI: 10.1038/s41536-022-00249-0]
- 46 **Rocca CJ**, Goodman SM, Dulin JN, Haquang JH, Gertsman I, Blondelle J, Smith JLM, Heyser CJ, Cherqui S. Transplantation of wild-type mouse hematopoietic stem and progenitor cells ameliorates deficits in a mouse model of Friedreich's ataxia. *Sci Transl Med* 2017; **9**: eaaj2347 [PMID: 29070698 DOI: 10.1126/scitranslmed.aaj2347]
- 47 **Yoon JK**, Schindler JW, Loperfido M, Baricordi C, DeAndrade MP, Jacobs ME, Treleaven C, Plasschaert RN, Yan A, Barese CN, Dogan Y, Chen VP, Fiorini C, Hull F, Barbarossa L, Unnisa Z, Ivanov D, Kutner RH, Guda S, Oborski C, Maiwald T, Michaud V, Rothe M, Schambach A, Pfeifer R, Mason C, Biasco L, van Til NP. Preclinical lentiviral hematopoietic stem cell gene therapy corrects Pompe disease-related muscle and neurological manifestations. *Mol Ther* 2024; **32**: 3847-3864 [PMID: 39295144 DOI: 10.1016/j.ymthe.2024.09.024]
- 48 **Kono Y**, Kajita H, Okada T, Nakagawa R, Fujita T, Konishi S. Mesenchymal Stem Cells Promote IL-6 Secretion and Suppress the Gene Expression of Proinflammatory Cytokines in Contractile C2C12 Myotubes. *Biol Pharm Bull* 2022; **45**: 962-967 [PMID: 35786604 DOI: 10.1248/bpb.b22-00118]
- 49 **Reis ALG**, Maximino JR, Lage LAPC, Gomes HR, Pereira J, Brofman PRS, Senegaglia AC, Rebelatto CLK, Daga DR, Paiva WS, Chadi G. Proteomic analysis of cerebrospinal fluid of amyotrophic lateral sclerosis patients in the presence of autologous bone marrow derived mesenchymal stem cells. *Stem Cell Res Ther* 2024; **15**: 301 [PMID: 39278909 DOI: 10.1186/s13287-024-03820-2]
- 50 **Berry JD**, Cudkovic ME, Windebank AJ, Staff NP, Owegi M, Nicholson K, McKenna-Yasek D, Levy YS, Abramov N, Kaspi H, Mehra M, Aricha R, Gothelf Y, Brown RH. NurOwn, phase 2, randomized, clinical trial in patients with ALS: Safety, clinical, and biomarker results. *Neurology* 2019; **93**: e2294-e2305 [PMID: 31740545 DOI: 10.1212/WNL.0000000000008620]
- 51 **Forotti G**, Nizzardo M, Bucchia M, Ramirez A, Trombetta E, Gatti S, Bresolin N, Comi GP, Corti S. CSF transplantation of a specific iPSC-derived neural stem cell subpopulation ameliorates the disease phenotype in a mouse model of spinal muscular atrophy with respiratory distress type 1. *Exp Neurol* 2019; **321**: 113041 [PMID: 31445043 DOI: 10.1016/j.expneurol.2019.113041]
- 52 **Mohseni R**, Hamidieh AA, Shoaee-Hassani A, Ghahvechi-Akbari M, Majma A, Mohammadi M, Nikougoftar M, Shervin-Badv R, Ai J, Montazerlotfelahi H, Ashrafi MR. An open-label phase 1 clinical trial of the allogeneic side population adipose-derived mesenchymal stem cells in SMA type 1 patients. *Neurol Sci* 2022; **43**: 399-410 [PMID: 34032944 DOI: 10.1007/s10072-021-05291-2]
- 53 **Yeo CJJ**, Tizzano EF, Darras BT. Challenges and opportunities in spinal muscular atrophy therapeutics. *Lancet Neurol* 2024; **23**: 205-218 [PMID: 38267192 DOI: 10.1016/S1474-4422(23)00419-2]
- 54 **Meesuk L**, Suwanprateeb J, Thammarakcharoen F, Tantrawatpan C, Kheolamai P, Palang I, Tantikanlayaporn D, Manochantr S. Osteogenic differentiation and proliferation potentials of human bone marrow and umbilical cord-derived mesenchymal stem cells on the 3D-printed hydroxyapatite scaffolds. *Sci Rep* 2022; **12**: 19509 [PMID: 36376498 DOI: 10.1038/s41598-022-24160-2]
- 55 **Oguma Y**, Kuroda Y, Wakao S, Kushida Y, Dezawa M. Single-cell RNA sequencing reveals different signatures of mesenchymal stromal cell pluripotent-like and multipotent populations. *iScience* 2022; **25**: 105395 [PMID: 36339265 DOI: 10.1016/j.isci.2022.105395]
- 56 **Tsai YC**, Cheng TS, Liao HJ, Chuang MH, Chen HT, Chen CH, Zhang KL, Chang CH, Lin PC, Huang CF. Mesenchymal Stem Cell Secreted-Extracellular Vesicles are Involved in Chondrocyte Production and Reduce Adipogenesis during Stem Cell Differentiation. *Tissue Eng Regen Med* 2022; **19**: 1295-1310 [PMID: 36346531 DOI: 10.1007/s13770-022-00490-0]
- 57 **Li J**, Wu Z, Zhao L, Liu Y, Su Y, Gong X, Liu F, Zhang L. The heterogeneity of mesenchymal stem cells: an important issue to be addressed in cell therapy. *Stem Cell Res Ther* 2023; **14**: 381 [PMID: 38124129 DOI: 10.1186/s13287-023-03587-y]
- 58 **Abrigo J**, Rivera JC, Aravena J, Cabrera D, Simon F, Ezquer F, Ezquer M, Cabello-Verrugio C. High Fat Diet-Induced Skeletal Muscle Wasting Is Decreased by Mesenchymal Stem Cells Administration: Implications on Oxidative Stress, Ubiquitin Proteasome Pathway Activation, and Myonuclear Apoptosis. *Oxid Med Cell Longev* 2016; **2016**: 9047821 [PMID: 27579157 DOI: 10.1155/2016/9047821]
- 59 **Li TS**, Shi H, Wang L, Yan C. Effect of Bone Marrow Mesenchymal Stem Cells on Satellite Cell Proliferation and Apoptosis in Immobilization-Induced Muscle Atrophy in Rats. *Med Sci Monit* 2016; **22**: 4651-4660 [PMID: 27898654 DOI: 10.12659/msm.898137]
- 60 **Schilling BK**, Schusterman MA 2nd, Kim DY, Repko AJ, Klett KC, Christ GJ, Marra KG. Adipose-derived stem cells delay muscle atrophy after peripheral nerve injury in the rodent model. *Muscle Nerve* 2019; **59**: 603-610 [PMID: 30681163 DOI: 10.1002/mus.26432]
- 61 **Artioli GG**, De Oliveira Silvestre JG, Guilherme JP, Baptista IL, Ramos GV, Da Silva WJ, Miyabara EH, Moriscot AS. Embryonic stem cells improve skeletal muscle recovery after extreme atrophy in mice. *Muscle Nerve* 2015; **51**: 346-352 [PMID: 24934406 DOI: 10.1002/mus.24320]
- 62 **Li WY**, Zhu GY, Yue WJ, Sun GD, Zhu XF, Wang Y. KLF7 overexpression in bone marrow stromal stem cells graft transplantation promotes sciatic nerve regeneration. *J Neural Eng* 2019; **16**: 056011 [PMID: 31296795 DOI: 10.1088/1741-2552/ab3188]
- 63 **Mierzejewski B**, Michalska Z, Jackowski D, Stremimska W, Janczyk-Ilach K, Koblowska M, Iwanicka-Nowicka R, Gromadka A, Ciemerych

- MA, Brzoska E. The miR151 and miR5100 Transfected Bone Marrow Stromal Cells Increase Myoblast Fusion in IGF1 Dependent Manner. *Stem Cell Rev Rep* 2022; **18**: 2164-2178 [PMID: 35190967 DOI: 10.1007/s12015-022-10350-y]
- 64 **de Munter JPM**, Mey J, Strekalova T, Kramer BW, Wolters EC. Why do anti-inflammatory signals of bone marrow-derived stromal cells improve neurodegenerative conditions where anti-inflammatory drugs fail? *J Neural Transm (Vienna)* 2020; **127**: 715-727 [PMID: 32253536 DOI: 10.1007/s00702-020-02173-3]
- 65 **Sykova E**, Cizkova D, Kubinova S. Mesenchymal Stem Cells in Treatment of Spinal Cord Injury and Amyotrophic Lateral Sclerosis. *Front Cell Dev Biol* 2021; **9**: 695900 [PMID: 34295897 DOI: 10.3389/fcell.2021.695900]
- 66 **Świerczek-Lasek B**, Tolak L, Bijoch L, Stefaniuk M, Szpak P, Kalaszczynska I, Streminska W, Ciemerych MA, Archacka K. Comparison of Muscle Regeneration after BMSC-Conditioned Medium, Syngeneic, or Allogeneic BMSC Injection. *Cells* 2022; **11**: 2843 [PMID: 36139418 DOI: 10.3390/cells11182843]
- 67 **Helal MAM**, Shaheen NEM, Abu Zahra FA. Immunomodulatory capacity of the local mesenchymal stem cells transplantation after severe skeletal muscle injury in female rats. *Immunopharmacol Immunotoxicol* 2016; **38**: 414-422 [PMID: 27560658 DOI: 10.1080/08923973.2016.1222617]
- 68 **Andrade BM**, Baldanza MR, Ribeiro KC, Porto A, Peçanha R, Fortes FS, Zapata-Sudo G, Campos-de-Carvalho AC, Goldenberg RC, Werneck-de-Castro JP. Bone marrow mesenchymal cells improve muscle function in a skeletal muscle re-injury model. *PLoS One* 2015; **10**: e0127561 [PMID: 26039243 DOI: 10.1371/journal.pone.0127561]
- 69 **Maeda Y**, Yonemochi Y, Nakajyo Y, Hidaka H, Ikeda T, Ando Y. CXCL12 and osteopontin from bone marrow-derived mesenchymal stromal cells improve muscle regeneration. *Sci Rep* 2017; **7**: 3305 [PMID: 28607396 DOI: 10.1038/s41598-017-02928-1]
- 70 **Klimczak A**, Kozłowska U, Kurpisz M. Muscle Stem/Progenitor Cells and Mesenchymal Stem Cells of Bone Marrow Origin for Skeletal Muscle Regeneration in Muscular Dystrophies. *Arch Immunol Ther Exp (Warsz)* 2018; **66**: 341-354 [PMID: 29536116 DOI: 10.1007/s00005-018-0509-7]
- 71 **Archacka K**, Grabowska I, Mierzejewski B, Graffstein J, Górzynska A, Krawczyk M, Różycka AM, Kalaszczynska I, Muras G, Streminska W, Jańczyk-Ilach K, Walczak P, Janowski M, Ciemerych MA, Brzoska E. Hypoxia preconditioned bone marrow-derived mesenchymal stromal/stem cells enhance myoblast fusion and skeletal muscle regeneration. *Stem Cell Res Ther* 2021; **12**: 448 [PMID: 34372911 DOI: 10.1186/s13287-021-02530-3]
- 72 **Moussa MH**, Hamam GG, Abd Elaziz AE, Rahoma MA, Abd El Samad AA, El-Waseef DAA, Hegazy MA. Comparative Study on Bone Marrow-Versus Adipose-Derived Stem Cells on Regeneration and Re-Innervation of Skeletal Muscle Injury in Wistar Rats. *Tissue Eng Regen Med* 2020; **17**: 887-900 [PMID: 33030680 DOI: 10.1007/s13770-020-00288-y]
- 73 **Grabowska I**, Zimowska M, Maciejewska K, Jabłonska Z, Bazga A, Ozieblo M, Streminska W, Bem J, Brzoska E, Ciemerych MA. Adipose Tissue-Derived Stromal Cells in Matrigel Impacts the Regeneration of Severely Damaged Skeletal Muscles. *Int J Mol Sci* 2019; **20**: 3313 [PMID: 31284492 DOI: 10.3390/ijms20133313]
- 74 **Archacka K**, Bem J, Brzoska E, Czerwinska AM, Grabowska I, Kasprzycka P, Hoinkis D, Siennicka K, Pojda Z, Bernas P, Binkowski R, Jastrzebska K, Kupiec A, Malesza M, Michalczywska E, Soszynska M, Ilach K, Streminska W, Ciemerych MA. Beneficial Effect of IL-4 and SDF-1 on Myogenic Potential of Mouse and Human Adipose Tissue-Derived Stromal Cells. *Cells* 2020; **9**: 1479 [PMID: 32560483 DOI: 10.3390/cells9061479]
- 75 **Yang W**, Zhang S, Ou T, Jiang H, Jia D, Qi Z, Zou Y, Qian J, Sun A, Ge J. Interleukin-11 regulates the fate of adipose-derived mesenchymal stem cells via STAT3 signalling pathways. *Cell Prolif* 2020; **53**: e12771 [PMID: 32270546 DOI: 10.1111/cpr.12771]
- 76 **Leong J**, Hong YT, Wu YF, Ko E, Dvoretzkiy S, Teo JY, Kim BS, Kim K, Jeon H, Boppart M, Yang YY, Kong H. Surface Tethering of Inflammation-Modulatory Nanostimulators to Stem Cells for Ischemic Muscle Repair. *ACS Nano* 2020; **14**: 5298-5313 [PMID: 32243129 DOI: 10.1021/acsnano.9b04926]
- 77 **Ciervo Y**, Gatto N, Allen C, Grierson A, Ferraiuolo L, Mead RJ, Shaw PJ. Adipose-derived stem cells protect motor neurons and reduce glial activation in both in vitro and in vivo models of ALS. *Mol Ther Methods Clin Dev* 2021; **21**: 413-433 [PMID: 33869658 DOI: 10.1016/j.omtm.2021.03.017]
- 78 **Wang X**, Zhang Y, Jin T, Botchway BOA, Fan R, Wang L, Liu X. Adipose-Derived Mesenchymal Stem Cells Combined With Extracellular Vesicles May Improve Amyotrophic Lateral Sclerosis. *Front Aging Neurosci* 2022; **14**: 830346 [PMID: 35663577 DOI: 10.3389/fnagi.2022.830346]
- 79 **Xie Q**, Liu R, Jiang J, Peng J, Yang C, Zhang W, Wang S, Song J. What is the impact of human umbilical cord mesenchymal stem cell transplantation on clinical treatment? *Stem Cell Res Ther* 2020; **11**: 519 [PMID: 33261658 DOI: 10.1186/s13287-020-02011-z]
- 80 **Kim MJ**, Kim ZH, Kim SM, Choi YS. Conditioned medium derived from umbilical cord mesenchymal stem cells regenerates atrophied muscles. *Tissue Cell* 2016; **48**: 533-543 [PMID: 27457384 DOI: 10.1016/j.tice.2016.06.010]
- 81 **Piao L**, Huang Z, Inoue A, Kuzuya M, Cheng XW. Human umbilical cord-derived mesenchymal stromal cells ameliorate aging-associated skeletal muscle atrophy and dysfunction by modulating apoptosis and mitochondrial damage in SAMP10 mice. *Stem Cell Res Ther* 2022; **13**: 226 [PMID: 35659361 DOI: 10.1186/s13287-022-02895-z]
- 82 **Cohen S**, Nathan JA, Goldberg AL. Muscle wasting in disease: molecular mechanisms and promising therapies. *Nat Rev Drug Discov* 2015; **14**: 58-74 [PMID: 25549588 DOI: 10.1038/nrd4467]
- 83 **Colleluori G**, Aguirre L, Phadnis U, Fowler K, Armamento-Villareal R, Sun Z, Brunetti L, Hyoung Park J, Kaiparettu BA, Putluri N, Auetumrongasawat V, Yarasheski K, Qualls C, Villareal DT. Aerobic Plus Resistance Exercise in Obese Older Adults Improves Muscle Protein Synthesis and Preserves Myocellular Quality Despite Weight Loss. *Cell Metab* 2019; **30**: 261-273.e6 [PMID: 31279675 DOI: 10.1016/j.cmet.2019.06.008]
- 84 **Song J**, Liu J, Cui C, Hu H, Zang N, Yang M, Yang J, Zou Y, Li J, Wang L, He Q, Guo X, Zhao R, Yan F, Liu F, Hou X, Sun Z, Chen L. Mesenchymal stromal cells ameliorate diabetes-induced muscle atrophy through exosomes by enhancing AMPK/ULK1-mediated autophagy. *J Cachexia Sarcopenia Muscle* 2023; **14**: 915-929 [PMID: 36708027 DOI: 10.1002/jcsm.13177]
- 85 **Nishimura K**, Fukuda A, Hisatake K. Mechanisms of the Metabolic Shift during Somatic Cell Reprogramming. *Int J Mol Sci* 2019; **20**: 2254 [PMID: 31067778 DOI: 10.3390/ijms20092254]
- 86 **Genet M**, Torres-Padilla ME. The molecular and cellular features of 2-cell-like cells: a reference guide. *Development* 2020; **147**: dev189688 [PMID: 32847823 DOI: 10.1242/dev.189688]
- 87 **Liu B**, Li M, Zhang L, Chen Z, Lu P. Motor neuron replacement therapy for amyotrophic lateral sclerosis. *Neural Regen Res* 2022; **17**: 1633-1639 [PMID: 35017408 DOI: 10.4103/1673-5374.332123]
- 88 **Mora C**, Serzanti M, Consiglio A, Memo M, Dell'Era P. Clinical potentials of human pluripotent stem cells. *Cell Biol Toxicol* 2017; **33**: 351-

- 360 [PMID: 28176010 DOI: 10.1007/s10565-017-9384-y]
- 89 **Novelli G**, Spitalieri P, Murdocca M, Centanini E, Sanguolo F. Organoid factory: The recent role of the human induced pluripotent stem cells (hiPSCs) in precision medicine. *Front Cell Dev Biol* 2022; **10**: 1059579 [PMID: 36699015 DOI: 10.3389/fcell.2022.1059579]
- 90 **Zhang L**, Ma XJ, Fei YY, Han HT, Xu J, Cheng L, Li X. Stem cell therapy in liver regeneration: Focus on mesenchymal stem cells and induced pluripotent stem cells. *Pharmacol Ther* 2022; **232**: 108004 [PMID: 34597754 DOI: 10.1016/j.pharmthera.2021.108004]
- 91 **Iberite F**, Gruppioni E, Ricotti L. Skeletal muscle differentiation of human iPSCs meets bioengineering strategies: perspectives and challenges. *NPJ Regen Med* 2022; **7**: 23 [PMID: 35393412 DOI: 10.1038/s41536-022-00216-9]
- 92 **Ohuchi K**, Funato M, Kato Z, Seki J, Kawase C, Tamai Y, Ono Y, Nagahara Y, Noda Y, Kameyama T, Ando S, Tsuruma K, Shimazawa M, Hara H, Kaneko H. Established Stem Cell Model of Spinal Muscular Atrophy Is Applicable in the Evaluation of the Efficacy of Thyrotropin-Releasing Hormone Analog. *Stem Cells Transl Med* 2016; **5**: 152-163 [PMID: 26683872 DOI: 10.5966/sctm.2015-0059]
- 93 **Deuse T**, Hu X, Gravina A, Wang D, Tediashvili G, De C, Thayer WO, Wahl A, Garcia JV, Reichenspurner H, Davis MM, Lanier LL, Schrepfer S. Hypoimmunogenic derivatives of induced pluripotent stem cells evade immune rejection in fully immunocompetent allogeneic recipients. *Nat Biotechnol* 2019; **37**: 252-258 [PMID: 30778232 DOI: 10.1038/s41587-019-0016-3]
- 94 **Pinton L**, Khedr M, Lionello VM, Sarcar S, Maffioletti SM, Dastidar S, Negroni E, Choi S, Khokhar N, Bigot A, Counsell JR, Bernardo AS, Zammit PS, Tedesco FS. 3D human induced pluripotent stem cell-derived bioengineered skeletal muscles for tissue, disease and therapy modeling. *Nat Protoc* 2023; **18**: 1337-1376 [PMID: 36792780 DOI: 10.1038/s41596-022-00790-8]
- 95 **Kim H**, Perlingeiro RCR. Generation of human myogenic progenitors from pluripotent stem cells for in vivo regeneration. *Cell Mol Life Sci* 2022; **79**: 406 [PMID: 35802202 DOI: 10.1007/s00018-022-04434-8]
- 96 **Sun C**, Kannan S, Choi IY, Lim H, Zhang H, Chen GS, Zhang N, Park SH, Serra C, Iyer SR, Lloyd TE, Kwon C, Lovering RM, Lim SB, Andersen P, Wagner KR, Lee G. Human pluripotent stem cell-derived myogenic progenitors undergo maturation to quiescent satellite cells upon engraftment. *Cell Stem Cell* 2022; **29**: 610-619.e5 [PMID: 35395188 DOI: 10.1016/j.stem.2022.03.004]
- 97 **Kim H**, Selvaraj S, Kiley J, Azzag K, Garay BI, Perlingeiro RCR. Genomic Safe Harbor Expression of PAX7 for the Generation of Engraftable Myogenic Progenitors. *Stem Cell Reports* 2021; **16**: 10-19 [PMID: 33275879 DOI: 10.1016/j.stemcr.2020.11.001]
- 98 **Rao L**, Qian Y, Khodabukus A, Ribar T, Bursac N. Engineering human pluripotent stem cells into a functional skeletal muscle tissue. *Nat Commun* 2018; **9**: 126 [PMID: 29317646 DOI: 10.1038/s41467-017-02636-4]
- 99 **Shahriyari M**, Islam MR, Sakib SM, Rinn M, Rika A, Krüger D, Kaurani L, Gisa V, Winterhoff M, Anandakumar H, Shomroni O, Schmidt M, Salinas G, Unger A, Linke WA, Zschützsch J, Schmidt J, Bassel-Duby R, Olson EN, Fischer A, Zimmermann WH, Tiburcy M. Engineered skeletal muscle recapitulates human muscle development, regeneration and dystrophy. *J Cachexia Sarcopenia Muscle* 2022; **13**: 3106-3121 [PMID: 36254806 DOI: 10.1002/jcsm.13094]
- 100 **Pepper JP**, Wang TV, Hennes V, Sun SY, Ichida JK. Human Induced Pluripotent Stem Cell-Derived Motor Neuron Transplant for Neuromuscular Atrophy in a Mouse Model of Sciatic Nerve Injury. *JAMA Facial Plast Surg* 2017; **19**: 197-205 [PMID: 27978547 DOI: 10.1001/jamafacial.2016.1544]
- 101 **Adami R**, Bottai D. NSC Physiological Features in Spinal Muscular Atrophy: SMN Deficiency Effects on Neurogenesis. *Int J Mol Sci* 2022; **23**: 15209 [PMID: 36499528 DOI: 10.3390/ijms232315209]
- 102 **Abati E**, Bresolin N, Comi GP, Corti S. Preconditioning and Cellular Engineering to Increase the Survival of Transplanted Neural Stem Cells for Motor Neuron Disease Therapy. *Mol Neurobiol* 2019; **56**: 3356-3367 [PMID: 30120734 DOI: 10.1007/s12035-018-1305-4]
- 103 **Forostyak S**, Forostyak O, Kwok JCF, Romanyuk N, Rehorova M, Kriska J, Dayanithi G, Raha-Chowdhury R, Jendelova P, Anderova M, Fawcett JW, Sykova E. Transplantation of Neural Precursors Derived from Induced Pluripotent Cells Preserve Perineuronal Nets and Stimulate Neural Plasticity in ALS Rats. *Int J Mol Sci* 2020; **21**: 9593 [PMID: 33339362 DOI: 10.3390/ijms21249593]
- 104 **Wang W**, Li M, Chen Z, Xu L, Chang M, Wang K, Deng C, Gu Y, Zhou S, Shen Y, Tao F, Sun H. Biogenesis and function of extracellular vesicles in pathophysiological processes of skeletal muscle atrophy. *Biochem Pharmacol* 2022; **198**: 114954 [PMID: 35167807 DOI: 10.1016/j.bcp.2022.114954]
- 105 **Assoni A**, Coatti G, Valadares MC, Beccari M, Gomes J, Pelatti M, Mitne-Neto M, Carvalho VM, Zatz M. Different Donors Mesenchymal Stromal Cells Secretomes Reveal Heterogeneous Profile of Relevance for Therapeutic Use. *Stem Cells Dev* 2017; **26**: 206-214 [PMID: 27762666 DOI: 10.1089/scd.2016.0218]
- 106 **Youssef El Baradie KB**, Hamrick MW. Therapeutic application of extracellular vesicles for musculoskeletal repair & regeneration. *Connect Tissue Res* 2021; **62**: 99-114 [PMID: 32602385 DOI: 10.1080/03008207.2020.1781102]
- 107 **Yang Q**, Su S, Liu S, Yang S, Xu J, Zhong Y, Yang Y, Tian L, Tan Z, Wang J, Yu Z, Shi Z, Liang F. Exosomes-loaded electroconductive nerve dressing for nerve regeneration and pain relief against diabetic peripheral nerve injury. *Bioact Mater* 2023; **26**: 194-215 [PMID: 36923267 DOI: 10.1016/j.bioactmat.2023.02.024]
- 108 **Yan B**, Zhang Y, Liang C, Liu B, Ding F, Wang Y, Zhu B, Zhao R, Yu XY, Li Y. Stem cell-derived exosomes prevent pyroptosis and repair ischemic muscle injury through a novel exosome/circHIPK3/FOXO3a pathway. *Theranostics* 2020; **10**: 6728-6742 [PMID: 32550900 DOI: 10.7150/thno.42259]
- 109 **Ma Y**, Dong L, Zhou D, Li L, Zhang W, Zhen Y, Wang T, Su J, Chen D, Mao C, Wang X. Extracellular vesicles from human umbilical cord mesenchymal stem cells improve nerve regeneration after sciatic nerve transection in rats. *J Cell Mol Med* 2019; **23**: 2822-2835 [PMID: 30772948 DOI: 10.1111/jcmm.14190]
- 110 **Ma H**, Jing Y, Zeng J, Ge J, Sun S, Cui R, Qian C, Qu S, Sheng H. Human umbilical cord mesenchymal stem cell-derived exosomes ameliorate muscle atrophy via the miR-132-3p/FoxO3 axis. *J Orthop Translat* 2024; **49**: 23-36 [PMID: 39420945 DOI: 10.1016/j.jot.2024.08.005]
- 111 **Chen J**, Ren S, Duscher D, Kang Y, Liu Y, Wang C, Yuan M, Guo G, Xiong H, Zhan P, Wang Y, Machens HG, Chen Z. Exosomes from human adipose-derived stem cells promote sciatic nerve regeneration via optimizing Schwann cell function. *J Cell Physiol* 2019; **234**: 23097-23110 [PMID: 31124125 DOI: 10.1002/jcp.28873]
- 112 **Mellows B**, Mitchell R, Antonioli M, Kretz O, Chambers D, Zeuner MT, Denecke B, Musante L, Ramachandra DL, Debacq-Chainiaux F, Holthofer H, Joch B, Ray S, Widera D, David AL, Huber TB, Dengiel J, De Coppi P, Patel K. Protein and Molecular Characterization of a Clinically Compliant Amniotic Fluid Stem Cell-Derived Extracellular Vesicle Fraction Capable of Accelerating Muscle Regeneration Through Enhancement of Angiogenesis. *Stem Cells Dev* 2017; **26**: 1316-1333 [PMID: 28679310 DOI: 10.1089/scd.2017.0089]
- 113 **Zhu A**, Liu N, Shang Y, Zhen Y, An Y. Signaling pathways of adipose stem cell-derived exosomes promoting muscle regeneration. *Chin Med J (Engl)* 2022; **135**: 2525-2534 [PMID: 36583914 DOI: 10.1097/CM9.0000000000002404]
- 114 **Lo Sicco C**, Reverberi D, Balbi C, Ulivi V, Principi E, Pascucci L, Becherini P, Bosco MC, Varesio L, Franzin C, Pozzobon M, Cancedda R,

- Tasso R. Mesenchymal Stem Cell-Derived Extracellular Vesicles as Mediators of Anti-Inflammatory Effects: Endorsement of Macrophage Polarization. *Stem Cells Transl Med* 2017; **6**: 1018-1028 [PMID: 28186708 DOI: 10.1002/sctm.16-0363]
- 115 **Figeac N**, Zammit PS. Coordinated action of Axin1 and Axin2 suppresses β -catenin to regulate muscle stem cell function. *Cell Signal* 2015; **27**: 1652-1665 [PMID: 25866367 DOI: 10.1016/j.cellsig.2015.03.025]
- 116 **Gan L**, Zheng L, Yao L, Lei L, Huang Y, Zeng Z, Fang N. Exosomes from adipose-derived mesenchymal stem cells improve liver fibrosis by regulating the miR-20a-5p/TGFBR2 axis to affect the p38 MAPK/NF- κ B pathway. *Cytokine* 2023; **172**: 156386 [PMID: 37852157 DOI: 10.1016/j.cyto.2023.156386]
- 117 **Fan W**, Gao XK, Rao XS, Shi YP, Liu XC, Wang FY, Liu YF, Cong XX, He MY, Xu SB, Shen WL, Shen Y, Yan SG, Luo Y, Low BC, Ouyang H, Bao Z, Zheng LL, Zhou YT. Hsp70 Interacts with Mitogen-Activated Protein Kinase (MAPK)-Activated Protein Kinase 2 To Regulate p38MAPK Stability and Myoblast Differentiation during Skeletal Muscle Regeneration. *Mol Cell Biol* 2018; **38**: e00211-18 [PMID: 30275345 DOI: 10.1128/MCB.00211-18]
- 118 **Nowosad A**, Jeannot P, Callot C, Creff J, Perchey RT, Joffre C, Codogno P, Manenti S, Besson A. p27 controls Ragulator and mTOR activity in amino acid-deprived cells to regulate the autophagy-lysosomal pathway and coordinate cell cycle and cell growth. *Nat Cell Biol* 2020; **22**: 1076-1090 [PMID: 32807902 DOI: 10.1038/s41556-020-0554-4]
- 119 **Wang C**, Song W, Chen B, Liu X, He Y. Exosomes Isolated From Adipose-Derived Stem Cells: A New Cell-Free Approach to Prevent the Muscle Degeneration Associated With Torn Rotator Cuffs. *Am J Sports Med* 2019; **47**: 3247-3255 [PMID: 31560856 DOI: 10.1177/0363546519876323]
- 120 **Bonafede R**, Turano E, Scambi I, Busato A, Bontempi P, Virla F, Schiaffino L, Marzola P, Bonetti B, Mariotti R. ASC-Exosomes Ameliorate the Disease Progression in SOD1(G93A) Murine Model Underlining Their Potential Therapeutic Use in Human ALS. *Int J Mol Sci* 2020; **21**: 3651 [PMID: 32455791 DOI: 10.3390/ijms21103651]
- 121 **Bonafede R**, Brandi J, Manfredi M, Scambi I, Schiaffino L, Merigo F, Turano E, Bonetti B, Marengo E, Cecconi D, Mariotti R. The Anti-Apoptotic Effect of ASC-Exosomes in an In Vitro ALS Model and Their Proteomic Analysis. *Cells* 2019; **8**: 1087 [PMID: 31540100 DOI: 10.3390/cells8091087]
- 122 **Luo Z**, Lin J, Sun Y, Wang C, Chen J. Bone Marrow Stromal Cell-Derived Exosomes Promote Muscle Healing Following Contusion Through Macrophage Polarization. *Stem Cells Dev* 2021; **30**: 135-148 [PMID: 33323007 DOI: 10.1089/scd.2020.0167]
- 123 **Li Z**, Liu C, Li S, Li T, Li Y, Wang N, Bao X, Xue P, Liu S. BMSC-Derived Exosomes Inhibit Dexamethasone-Induced Muscle Atrophy via the miR-486-5p/FoxO1 Axis. *Front Endocrinol (Lausanne)* 2021; **12**: 681267 [PMID: 34659106 DOI: 10.3389/fendo.2021.681267]
- 124 **Pan J**, Zhao M, Yi X, Tao J, Li S, Jiang Z, Cheng B, Yuan H, Zhang F. Acellular nerve grafts supplemented with induced pluripotent stem cell-derived exosomes promote peripheral nerve reconstruction and motor function recovery. *Bioact Mater* 2022; **15**: 272-287 [PMID: 35356813 DOI: 10.1016/j.bioactmat.2021.12.004]
- 125 **Yu M**, Gu G, Cong M, Du M, Wang W, Shen M, Zhang Q, Shi H, Gu X, Ding F. Repair of peripheral nerve defects by nerve grafts incorporated with extracellular vesicles from skin-derived precursor Schwann cells. *Acta Biomater* 2021; **134**: 190-203 [PMID: 34289422 DOI: 10.1016/j.actbio.2021.07.026]
- 126 **Wang W**, Shen D, Zhang L, Ji Y, Xu L, Chen Z, Shen Y, Gong L, Zhang Q, Shen M, Gu X, Sun H. SKP-SC-EVs Mitigate Denervated Muscle Atrophy by Inhibiting Oxidative Stress and Inflammation and Improving Microcirculation. *Antioxidants (Basel)* 2021; **11**: 66 [PMID: 35052570 DOI: 10.3390/antiox11010066]
- 127 **Iyer SR**, Scheiber AL, Yarowsky P, Henn RF 3rd, Otsuru S, Lovering RM. Exosomes Isolated From Platelet-Rich Plasma and Mesenchymal Stem Cells Promote Recovery of Function After Muscle Injury. *Am J Sports Med* 2020; **48**: 2277-2286 [PMID: 32543878 DOI: 10.1177/0363546520926462]
- 128 **Pan Y**, Li L, Cao N, Liao J, Chen H, Zhang M. Advanced nano delivery system for stem cell therapy for Alzheimer's disease. *Biomaterials* 2025; **314**: 122852 [PMID: 39357149 DOI: 10.1016/j.biomaterials.2024.122852]
- 129 **Bronckaers A**, Hilkens P, Martens W, Gervois P, Ratajczak J, Struys T, Lambrechts I. Mesenchymal stem/stromal cells as a pharmacological and therapeutic approach to accelerate angiogenesis. *Pharmacol Ther* 2014; **143**: 181-196 [PMID: 24594234 DOI: 10.1016/j.pharmthera.2014.02.013]
- 130 **Ullah M**, Liu DD, Thakor AS. Mesenchymal Stromal Cell Homing: Mechanisms and Strategies for Improvement. *iScience* 2019; **15**: 421-438 [PMID: 31121468 DOI: 10.1016/j.isci.2019.05.004]
- 131 **Goodman SB**, Lin T. Modifying MSC Phenotype to Facilitate Bone Healing: Biological Approaches. *Front Bioeng Biotechnol* 2020; **8**: 641 [PMID: 32671040 DOI: 10.3389/fbioe.2020.00641]
- 132 **Dabrowska S**, Andrzejewska A, Lukomska B, Janowski M. Neuroinflammation as a target for treatment of stroke using mesenchymal stem cells and extracellular vesicles. *J Neuroinflammation* 2019; **16**: 178 [PMID: 31514749 DOI: 10.1186/s12974-019-1571-8]
- 133 **Zhang F**, Leong W, Su K, Fang Y, Wang DA. A transduced living hyaline cartilage graft releasing transgenic stromal cell-derived factor-1 inducing endogenous stem cell homing in vivo. *Tissue Eng Part A* 2013; **19**: 1091-1099 [PMID: 23167782 DOI: 10.1089/ten.TEA.2012.0441]
- 134 **Chen L**, Li Y, Chen W, Han N, Li K, Guo R, Liu Z, Xiao Y. Enhanced recruitment and hematopoietic reconstitution of bone marrow-derived mesenchymal stem cells in bone marrow failure by the SDF-1/CXCR4. *J Tissue Eng Regen Med* 2020; **14**: 1250-1260 [PMID: 32633015 DOI: 10.1002/term.3096]
- 135 **Al-Kharboosh R**, ReFaey K, Lara-Velazquez M, Grewal SS, Imitola J, Quiñones-Hinojosa A. Inflammatory Mediators in Glioma Microenvironment Play a Dual Role in Gliomagenesis and Mesenchymal Stem Cell Homing: Implication for Cellular Therapy. *Mayo Clin Proc Innov Qual Outcomes* 2020; **4**: 443-459 [PMID: 32793872 DOI: 10.1016/j.mayocpiqo.2020.04.006]
- 136 **Ogay V**, Sekenova A, Li Y, Issabekova A, Saporov A. The Therapeutic Potential of Mesenchymal Stem Cells in the Treatment of Atherosclerosis. *Curr Stem Cell Res Ther* 2021; **16**: 897-913 [PMID: 33511957 DOI: 10.2174/1574888X16999210128193549]
- 137 **Sarsenova M**, Issabekova A, Abisheva S, Rutsakaya-Moroshan K, Ogay V, Saporov A. Mesenchymal Stem Cell-Based Therapy for Rheumatoid Arthritis. *Int J Mol Sci* 2021; **22**: 11592 [PMID: 34769021 DOI: 10.3390/ijms222111592]
- 138 **Szydlak R**. Biological, chemical and mechanical factors regulating migration and homing of mesenchymal stem cells. *World J Stem Cells* 2021; **13**: 619-631 [PMID: 34249231 DOI: 10.4252/wjsc.v13.i6.619]
- 139 **Jovic D**, Yu Y, Wang D, Wang K, Li H, Xu F, Liu C, Liu J, Luo Y. A Brief Overview of Global Trends in MSC-Based Cell Therapy. *Stem Cell Rev Rep* 2022; **18**: 1525-1545 [PMID: 35344199 DOI: 10.1007/s12015-022-10369-1]
- 140 **Shi Y**, Wang Y, Li Q, Liu K, Hou J, Shao C, Wang Y. Immunoregulatory mechanisms of mesenchymal stem and stromal cells in inflammatory diseases. *Nat Rev Nephrol* 2018; **14**: 493-507 [PMID: 29895977 DOI: 10.1038/s41581-018-0023-5]
- 141 **Pezzi A**, Amarin B, Laureano Á, Valim V, Dahmer A, Zambonato B, Sehn F, Wilke I, Bruschi L, Silva MALD, Filippi-Chiela E, Silla L.

- Effects Of Hypoxia in Long-Term In Vitro Expansion of Human Bone Marrow Derived Mesenchymal Stem Cells. *J Cell Biochem* 2017; **118**: 3072-3079 [PMID: 28240374 DOI: 10.1002/jcb.25953]
- 142 Siegel G, Kluba T, Hermanutz-Klein U, Bieback K, Northoff H, Schäfer R. Phenotype, donor age and gender affect function of human bone marrow-derived mesenchymal stromal cells. *BMC Med* 2013; **11**: 146 [PMID: 23758701 DOI: 10.1186/1741-7015-11-146]
- 143 Stubbendorff M, Deuse T, Hua X, Phan TT, Bieback K, Atkinson K, Eiermann TH, Velden J, Schröder C, Reichenspurner H, Robbins RC, Volk HD, Schrepfer S. Immunological properties of extraembryonic human mesenchymal stromal cells derived from gestational tissue. *Stem Cells Dev* 2013; **22**: 2619-2629 [PMID: 23711207 DOI: 10.1089/scd.2013.0043]
- 144 Zhuang X, Hu X, Zhang S, Li X, Yuan X, Wu Y. Mesenchymal Stem Cell-Based Therapy as a New Approach for the Treatment of Systemic Sclerosis. *Clin Rev Allergy Immunol* 2023; **64**: 284-320 [PMID: 35031958 DOI: 10.1007/s12016-021-08892-z]
- 145 Dreia K, Stanaszek L, Nowakowski A, Kuczynska Z, Lukomska B. Experimental Strategies of Mesenchymal Stem Cell Propagation: Adverse Events and Potential Risk of Functional Changes. *Stem Cells Int* 2019; **2019**: 7012692 [PMID: 30956673 DOI: 10.1155/2019/7012692]
- 146 Yang YK, Ogando CR, Wang See C, Chang TY, Barabino GA. Changes in phenotype and differentiation potential of human mesenchymal stem cells aging in vitro. *Stem Cell Res Ther* 2018; **9**: 131 [PMID: 29751774 DOI: 10.1186/s13287-018-0876-3]
- 147 Al Tanoury Z, Rao J, Tassy O, Gobert B, Gapon S, Garnier JM, Wagner E, Hick A, Hall A, Gussoni E, Pourquie O. Differentiation of the human PAX7-positive myogenic precursors/satellite cell lineage in vitro. *Development* 2020; **147**: dev187344 [PMID: 32541004 DOI: 10.1242/dev.187344]
- 148 Shamsian A, Sahebnasagh R, Norouzy A, Hussein SH, Ghahremani MH, Azizi Z. Cancer cells as a new source of induced pluripotent stem cells. *Stem Cell Res Ther* 2022; **13**: 459 [PMID: 36064437 DOI: 10.1186/s13287-022-03145-y]
- 149 Xiong J, Hu H, Guo R, Wang H, Jiang H. Mesenchymal Stem Cell Exosomes as a New Strategy for the Treatment of Diabetes Complications. *Front Endocrinol (Lausanne)* 2021; **12**: 646233 [PMID: 33995278 DOI: 10.3389/fendo.2021.646233]
- 150 Rezaie J, Nejati V, Mahmoodi M, Ahmadi M. Mesenchymal stem cells derived extracellular vesicles: A promising nanomedicine for drug delivery system. *Biochem Pharmacol* 2022; **203**: 115167 [PMID: 35820499 DOI: 10.1016/j.bcp.2022.115167]
- 151 Li Y, Kamei Y, Kambe M, Ebisawa K, Oishi M, Takanari K. Peripheral Nerve Regeneration Using Different Germ Layer-Derived Adult Stem Cells in the Past Decade. *Behav Neurol* 2021; **2021**: 5586523 [PMID: 34539934 DOI: 10.1155/2021/5586523]
- 152 Shan Y, Zhang M, Tao E, Wang J, Wei N, Lu Y, Liu Q, Hao K, Zhou F, Wang G. Pharmacokinetic characteristics of mesenchymal stem cells in translational challenges. *Signal Transduct Target Ther* 2024; **9**: 242 [PMID: 39271680 DOI: 10.1038/s41392-024-01936-8]
- 153 Winkler T, Perka C, von Roth P, Agres AN, Plage H, Preininger B, Pumberger M, Geissler S, Hagai EL, Ofir R, Pinzur L, Eyal E, Stoltenburg-Didinger G, Meisel C, Consentius C, Streitz M, Reinke P, Duda GN, Volk HD. Immunomodulatory placental-expanded, mesenchymal stromal cells improve muscle function following hip arthroplasty. *J Cachexia Sarcopenia Muscle* 2018; **9**: 880-897 [PMID: 30230266 DOI: 10.1002/jcsm.12316]
- 154 Jiang L, Mee T, Zhou X, Jia X. Augmenting Peripheral Nerve Regeneration with Adipose-Derived Stem Cells. *Stem Cell Rev Rep* 2022; **18**: 544-558 [PMID: 34417730 DOI: 10.1007/s12015-021-10236-5]
- 155 Kolagar TA, Farzaneh M, Nikkar N, Khoshnam SE. Human Pluripotent Stem Cells in Neurodegenerative Diseases: Potentials, Advances and Limitations. *Curr Stem Cell Res Ther* 2020; **15**: 102-110 [PMID: 31441732 DOI: 10.2174/1574888X14666190823142911]
- 156 Tan F, Li X, Wang Z, Li J, Shahzad K, Zheng J. Clinical applications of stem cell-derived exosomes. *Signal Transduct Target Ther* 2024; **9**: 17 [PMID: 38212307 DOI: 10.1038/s41392-023-01704-0]
- 157 Luo W, Zhang H, Wan R, Cai Y, Liu Y, Wu Y, Yang Y, Chen J, Zhang D, Luo Z, Shang X. Biomaterials-Based Technologies in Skeletal Muscle Tissue Engineering. *Adv Healthc Mater* 2024; **13**: e2304196 [PMID: 38712598 DOI: 10.1002/adhm.202304196]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

