

Supplementary Material

Supplementary Table 1 Summary of all pediatric patients who underwent endoscopic ultrasound-guided pseudocyst drainage

No.	Age (years)	Sex	Location pseudocyst	of Size (cm × cm)	EUS-guided intervention
1	18	Male	Pancreatic body	8.8 × 7.0	Stent placement
2	12	Male	Head and neck	6.1 × 6.0	Stent placement
3	11	Male	Pancreatic body	14.0 × 10.0	Stent placement
4	13	Female	Pancreatic body	6.0 × 7.0	Stent placement
5	14	Female	Pancreatic body	10.0 × 10.0	Stent placement
6	12	Female	Pancreatic body	11.0 × 12.0	Stent placement
7	16	Male	Pancreatic body	8.7 × 7.0	Stent placement
8	14	Female	Pancreatic tail	8.0 × 11.0	Stent placement
9	16	Male	Pancreatic tail	7.0 × 6.0	Stent placement
10	16	Male	Pancreatic tail	5.0 × 6.0	Stent placement
11	8	Male	Pancreatic tail	5.8 × 3.6	Complete aspiration
12	18	Male	Pancreatic body	2.0 × 1.2	Complete aspiration
13	17	Male	Head and neck	3.0 × 3.0	Complete aspiration
14	15	Female	Head and neck	6.4 × 4.0	Complete aspiration
15	15	Female	Head and neck	3.0 × 2.6	Complete aspiration
16	6	Female	Head and neck	3.0 × 1.5	Complete aspiration
17	8	Male	Pancreatic body	6.0 × 5.0	Complete aspiration
18	15	Male	Pancreatic tail	6.0 × 4.0	Complete aspiration
19	13	Female	Pancreatic tail	5.5 × 5.0	Complete aspiration

No.	Age (years)	Sex	Location pseudocyst	of Size (cm × cm)	EUS-guided intervention
20	18	Female	Pancreatic tail	2.4 × 2.4	Complete aspiration
21	15	Female	Pancreatic tail	6.7 × 5.6	Complete aspiration (recurrence case)

¹Patients with a history of abdominal trauma leading to pseudocyst formation; ²Patients with no history of pancreatitis; ³Pseudocyst drainage *via* lumen-apposing metal stent. EUS: Endoscopic ultrasound.