Supplementary Table 1 Timeline of historical and current information

Time	Event	Intervention	Outcome
point			
May 2022	Patient visited our clinic because	None	
	of abnormal vaginal bleeding		
	and an abdominal mass		
Jun 2022	Transvaginal us scan showed a	Optimal debulking was	
	$13 \text{ cm} \times 8 \text{ cm} \times 12 \text{ cm}$ solid mass	achieved with no	Rapid
	posterior to the uterus. Scrum	macroscopic residual	postoperative
	beta-hCG was 214405.5 mIU/mL	tumor (R0) was	recovery
		performe	
Jul 2022	Postoperatively, the scrum beta-	The adjuvant	No adverse
	hCG decreased to 796.7 mIU/ml.	chemotherapy of BEP	reaction
	Immunohistochemical	protocol was	
	comfirmed the diagnosis of	administrated	
	primary pure non-gestational		
	ovarian choriocarcinoma		
Sep 2022	The scrum beta-hCG decreased	Two cycles of adjuvant	No adverse
	to normal Level	chemotherapy had been	reaction
		performed	
Dec 2022		Four cycles of adjuvant	No sign of
		chemotherapy had been	recurrence and
		performed	metastasis

beta-hCG: Beta-human chorionic gonadotropin; BEP: Bleomycin, etoposide, and cisplatin.