

93258_Auto_Edited.docx

WORD COUNT

1628

TIME SUBMITTED

02-JUL-2024 11:33AM

PAPER ID

109985284

3

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 93258

Manuscript Type: EDITORIAL

Solid pancreatic masses: What's hidden beneath? Insights into rare pancreatic lesions

Balaban DV. Insights into rare pancreatic lesions

Daniel V Balaban

2

Daniel V Balaban, Department of Internal Medicine and Gastroenterology, Carol Davila University of Medicine and Pharmacy, Bucharest 020021, Romania

Author contributions: Balaban DV revised the case reports and wrote the manuscript.

1

Corresponding author: Daniel Vasile Balaban, MD, PhD, Adjunct Associate Professor, Department of Internal Medicine and Gastroenterology, Carol Davila University of Medicine and Pharmacy, 37 Dionisie Lupu, Bucharest 020021, Romania. vasile.balaban@umfcd.ro

Received: February 22, 2024

Revised: June 27, 2024

Abstract

The diagnostic approach to solid pancreatic masses has significantly evolved from the era when a focal pancreatic mass was almost synonymous to pancreatic ductal adenocarcinoma, to a wide spectrum of pancreatic lesions, some of them having good prognosis. With the advent of advanced diagnostic tools, particularly refined imaging and tissue acquisition techniques, a broader spectrum of differential diagnoses has been recognized, encompassing conditions ranging from neuroendocrine tumors or inflammatory masses, to rare entities like metastatic clear cell sarcoma or solitary fibrous tumors. We herein discuss case presentations reporting on some rare pancreatic lesions, which were diagnosed by combining clinical and imaging features, endoscopic ultrasound-guided tissue sampling and confirmed on surgical specimens. Further reports on these rare pancreatic tumors will contribute to better understanding of their pathogenesis and effective management.

Key Words: Pancreatic mass; Pancreas; Metastasis; Diagnosis; Endoscopic ultrasound

Balaban DV. Solid pancreatic masses: What's hidden beneath? Insights into rare pancreatic lesions. *World J Clin Cases* 2024; In press

Core Tip: The diagnostic landscape for solid pancreatic masses has expanded beyond the traditional assumption of pancreatic ductal adenocarcinoma and few other more common lesions. Advancements in imaging, tissue sampling and histopathological techniques, have unveiled a wide spectrum of pancreatic lesions, including rare tumors. In this editorial, we comment on case reports of some rare pancreatic lesions, such as metastatic clear cell sarcoma of the pancreas and pancreatic solitary fibrous tumor, emphasizing the importance of a comprehensive diagnostic approach, integrating medical history, cross-sectional imaging, and endoscopic ultrasound-guided tissue sampling.

INTRODUCTION

In the landscape of pancreatic pathology, the diagnosis of focal pancreatic masses has undergone tremendous progress in the last years. While a few decades ago, the presence of a focal mass in the pancreas was almost synonymous to a diagnosis of pancreatic ductal adenocarcinoma (PDAC), nowadays we have a wide spectrum of differential diagnosis for solid pancreatic tumors. This historical perspective was largely due to the limited diagnostic tools at that time, which led to PDAC often being considered the primary, if not sole, diagnosis in front of a pancreatic mass. Indeed, PDAC accounts for most of these lesions, but we can encounter other diagnosis such as neuroendocrine tumors, autoimmune pancreatitis, metastasis, lymphoma, inflammatory masses, tuberculosis, acinar cell carcinoma, pancreatoblastoma and other rare tumors[1]. While PDAC is well known for its aggressive tumor biology and poor prognosis, there are particular histological variants and also non-PDAC lesions associated with better outcomes[2].

Advancements in imaging techniques, consisting in refinement of cross-sectional imaging and the development and large-scale use of endoscopic ultrasound (EUS), have unveiled a wide spectrum of solid pancreatic masses. In the era of EUS, with enhanced imaging techniques and tissue acquisition by means of fine needle aspiration or fine needle biopsy (FNB), we can secure the diagnosis for a focal pancreatic lesion with high accuracy. In a recent meta-analysis, EUS-FNB has shown 93.1% sensitivity and 100% specificity in the diagnosis of pancreatic cancer[3].

REPORTS ON RARE PANCREATIC LESIONS

In light of these introductory remarks, we highlight on some case reports published in *World J Clin Cases*. First, there is the case presentation on metastatic clear cell sarcoma (CCS) of the pancreas, by Liu *et al*[4], which describes the rare case of CCS with pancreatic metastasis. While CCS is known to metastasize to lungs, bones and brain, pancreatic involvement is infrequent, as is primary pancreatic CCS. The authors highlight the relevance of the past medical history of the patient in defining the diagnosis and present the therapeutic strategy, consisting in distal pancreatectomy with splenectomy. While CCS bears a poor prognosis, obtaining

negative margins on the surgical specimen for the pancreatic tumor in the reported case is of paramount importance from an oncologic perspective. Another point made by the authors is regarding the requirement of long-term follow-up of patients with CCS, as metastasis can occur late.

In cases of suspected pancreatic metastasis, a thorough medical history, along with careful physical examination, are very important prerequisites for imaging and histology confirmation of a clinically suspected secondary pancreatic tumor. Most common primary tumor that metastasize to the pancreas is renal cell carcinoma, with a particular good prognosis, followed by colorectal cancer, melanoma, breast and lung cancer[5,6]. While pancreatic metastasis can be asymptomatic and detected incidentally on imaging during oncologic follow-up for the primary tumor, they can become clinically manifest as in the case report by Liu *et al*[4]. Differential diagnosis is based on imaging features, enhancement patterns and confirmed by histopathological sampling. EUS-guided tissue acquisition has proven high diagnostic accuracy in diagnosing pancreatic metastasis, particularly with the procurement of core tissue using FNB needles[6].

Another rare lesion encountered in the pancreas is solitary fibrous tumor, as presented in the case by Wang *et al*[7]. The solitary fibrous tumor is of mesenchymal origin, primary located in the pleura, but can develop in extrapleural sites also. In contrast with the previous case, the patient presented by Wang *et al*[7] did not have any complaints, and was detected with a pancreatic mass on physical examination, which was later confirmed on computed-tomography (CT) scan. While the CT showed a mixed density mass, with differential diagnosis of a solid pseudopapillary tumor and neuroendocrine tumor, the definite diagnosis was obtained by EUS-FNB, with histopathology and immunohistochemistry confirming a solitary fibrous tumor. As some solid pancreatic masses have similar structure and enhancing patterns, imaging features alone cannot confirm the precise type of lesion, but can narrow the differential diagnosis, and in most cases tissue sampling is required for positive diagnosis and therapeutic decision. As in the previous case report, distal pancreatectomy with splenectomy, this time by laparoscopic approach, was carried out, without recurrence on follow-up imaging. The authors point out the specific

immunohistochemistry profile of this rare pancreatic tumor, which prompts for differential diagnosis with other rare lesions such as sarcoma, neural or vascular tumors. Notably, another case of pancreatic solitary fibrous tumor, this time metastasized after a primary tumor in the central nervous system, was also reported recently by Yi *et al*[8].

Due to rarity of some pancreatic lesions as the ones presented, optimal management and follow-up are not standardized. Besides oncological surveillance, long term morbidity after pancreatic resection, comprising pancreatic exocrine insufficiency and metabolic complications, should also be monitored.

CONCLUSION

Solid pancreatic masses are more than just PDAC and other more commonly encountered tumors. Variability in clinical presentation and imaging features, also rarity of some lesions, make it difficult to conclude a diagnosis. Good history taking, refined imaging and EUS-guided tissue acquisition can accurately provide a preoperative definite diagnosis for focal pancreatic masses. Reports of cases of these rare pancreatic lesions will contribute to better understanding of their biology and to develop effective therapeutic strategies.

REFERENCES

- 1 **Iglesias-Garcia J**, de la Iglesia-Garcia D, Olmos-Martinez JM, Lariño-Noia J, Dominguez-Muñoz JE. Differential diagnosis of solid pancreatic masses. *Minerva Gastroenterol Dietol* 2020; **66**: 70-81 [PMID: 31994370 DOI: 10.23736/S1121-421X.20.02646-X]
- 2 **Schawkat K**, Manning MA, Glickman JN, Morteale KJ. Pancreatic Ductal Adenocarcinoma and Its Variants: Pearls and Perils. *Radiographics* 2020; **40**: 1219-1239 [PMID: 32678699 DOI: 10.1148/rg.2020190184]
- 3 **Facciorusso A**, Bajwa HS, Menon K, Buccino VR, Muscatiello N. Comparison between 22G aspiration and 22G biopsy needles for EUS-guided sampling of pancreatic lesions: A meta-analysis. *Endosc Ultrasound* 2020; **9**: 167-174 [PMID: 31031330 DOI: 10.4103/eus.eus_4_19]

4 **Liu YJ**, Zou C, Wu YY. Metastatic clear cell sarcoma of the pancreas: A rare case report. *World J Clin Cases* 2024; **12**: 1448-1453 [PMID: 38576799 DOI: 10.12998/wjcc.v12.i8.1448]

5 **Zerbi A**, Pecorelli N. Pancreatic metastases: An increasing clinical entity. *World J Gastrointest Surg* 2010; **2**: 255-259 [PMID: 21160884 DOI: 10.4240/wjgs.v2.i8.255]

6 **Spadaccini M**, Conti Bellocchi MC, Mangiavillano B, Fantin A, Rahal D, Manfrin E, Gavazzi F, Bozzarelli S, Crinò SF, Terrin M, Di Leo M, Bonifacio C, Facciorusso A, Realdon S, Cristofori C, Auriemma F, Fugazza A, Frulloni L, Hassan C, Repici A, Carrara S. Secondary Tumors of the Pancreas: A Multicenter Analysis of Clinicopathological and Endosonographic Features. *J Clin Med* 2023; **12** [PMID: 37109171 DOI: 10.3390/jcm12082829]

7 **Wang WW**, Zhou SP, Wu X, Wang LL, Ruan Y, Lu J, Li HL, Ni XL, Qiu LL, Zhou XH. Imaging, pathology, and diagnosis of solitary fibrous tumor of the pancreas: A case report and review of literature. *World J Clin Cases* 2024; **12**: 995-1003 [PMID: 38414609 DOI: 10.12998/wjcc.v12.i5.995]

8 **Yi K**, Lee J, Kim DU. Metastatic pancreatic solitary fibrous tumor: A case report. *World J Clin Cases* 2023; **11**: 8416-8424 [PMID: 38130626 DOI: 10.12998/wjcc.v11.i35.8416]

Footnotes

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Specialty type: Medicine, research and experimental

Country of origin: Romania

Peer-review report's classification

Scientific Quality: Grade B

Novelty: Grade B

Creativity or Innovation: Grade B

Scientific Significance: Grade B

P-Reviewer: Luo H **S-Editor:** Fan M **L-Editor:** P-Editor:

5%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|---|---|---------------|
| 1 | Daniel Vasile Balaban, Iulia Enache, Marina Ciochina, Alina Popp, Mariana Jinga. "Pancreatic involvement in celiac disease", World Journal of Gastroenterology, 2022
<small>Crossref</small> | 22 words — 2% |
| 2 | referencecitationanalysis.com
<small>Internet</small> | 17 words — 1% |
| 3 | f6publishing.blob.core.windows.net
<small>Internet</small> | 13 words — 1% |
| 4 | propertibazar.com
<small>Internet</small> | 12 words — 1% |
-

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES OFF

EXCLUDE MATCHES < 12 WORDS