World Journal of *Gastrointestinal Oncology*

World J Gastrointest Oncol 2024 December 15; 16(12): 4532-4781





Published by Baishideng Publishing Group Inc

World Journal of Gastrointestinal Oncology

Contents

Monthly Volume 16 Number 12 December 15, 2024

EDITORIAL

4532	Mixed neuroendocrine non-neuroendocrine tumors: The quest for evidence
	Cives M, Porta C, Palmirotta R
4537	Is nutritional status a new indicator to use in clinical practice for colorectal cancer patients?
	Berardi R, Chiariotti R, Mentrasti G
4543	Gene targets with therapeutic potential in hepatocellular carcinoma
	Shodry S, Hasan YTN, Ahdi IR, Ulhaq ZS
4548	Estimating prognosis of gastric neuroendocrine neoplasms using machine learning: A step towards precision medicine
	Wang HN, An JH, Zong L
4553	Exploring Xiaojianzhong decoction's potential in gastric cancer treatment: Integrative insights and experi- mental validation
	Cheng CH, Hao WR, Cheng TH
4559	Critical considerations for the management of gastrointestinal mixed neuroendocrine non-neuroendocrine neoplasms and pure neuroendocrine carcinomas
	Pavlidis ET, Galanis IN, Pavlidis TE
	REVIEW
4565	REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer
4565	REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer <i>Cui JY, Ma J, Gao XX, Sheng ZM, Pan ZX, Shi LH, Zhang BG</i>
4565	REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer <i>Cui JY, Ma J, Gao XX, Sheng ZM, Pan ZX, Shi LH, Zhang BG</i>
4565	REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer <i>Cui JY, Ma J, Gao XX, Sheng ZM, Pan ZX, Shi LH, Zhang BG</i> ORIGINAL ARTICLE
4565	REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer <i>Cui JY, Ma J, Gao XX, Sheng ZM, Pan ZX, Shi LH, Zhang BG</i> ORIGINAL ARTICLE Case Control Study
4565 4579	REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer Cui JY, Ma J, Gao XX, Sheng ZM, Pan ZX, Shi LH, Zhang BG ORIGINAL ARTICLE Case Control Study Prognostic utility of gamma-glutamyl transpeptidase to platelet ratio in patients with solitary hepatitis B virus-related hepatocellular carcinoma after hepatectomy
4565 4579	REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer <i>Cui JY, Ma J, Gao XX, Sheng ZM, Pan ZX, Shi LH, Zhang BG</i> ORIGINAL ARTICLE Case Control Study Prognostic utility of gamma-glutamyl transpeptidase to platelet ratio in patients with solitary hepatitis B virus-related hepatocellular carcinoma after hepatectomy <i>Yang CK, Wei ZL, Shen XQ, Jia YX, Wu QY, Wei YG, Su H, Qin W, Liao XW, Zhu GZ, Peng T</i>
4565 4579	REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer <i>Cui JY, Ma J, Gao XX, Sheng ZM, Pan ZX, Shi LH, Zhang BG</i> ORIGINAL ARTICLE Case Control Study Prognostic utility of gamma-glutamyl transpeptidase to platelet ratio in patients with solitary hepatitis B virus-related hepatocellular carcinoma after hepatectomy <i>Yang CK, Wei ZL, Shen XQ, Jia YX, Wu QY, Wei YG, Su H, Qin W, Liao XW, Zhu GZ, Peng T</i> Retrospective Cohort Study
4565 4579 4597	 REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer <i>Cui JY, Ma J, Gao XX, Sheng ZM, Pan ZX, Shi LH, Zhang BG</i> ORIGINAL ARTICLE Case Control Study Prognostic utility of gamma-glutamyl transpeptidase to platelet ratio in patients with solitary hepatitis B virus-related hepatocellular carcinoma after hepatectomy <i>Yang CK, Wei ZL, Shen XQ, Jia YX, Wu QY, Wei YG, Su H, Qin W, Liao XW, Zhu GZ, Peng T</i> Retrospective Cohort Study Prognostic prediction models for postoperative patients with stage I to III colorectal cancer based on
4565 4579 4597	REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer Cui JY, Ma J, Gao XX, Sheng ZM, Pan ZX, Shi LH, Zhang BG ORIGINAL ARTICLE Case Control Study Prognostic utility of gamma-glutamyl transpeptidase to platelet ratio in patients with solitary hepatitis B virus-related hepatocellular carcinoma after hepatectomy Yang CK, Wei ZL, Shen XQ, Jia YX, Wu QY, Wei YG, Su H, Qin W, Liao XW, Zhu GZ, Peng T Retrospective Cohort Study Prognostic prediction models for postoperative patients with stage I to III colorectal cancer based on machine learning
4565 4579 4597	REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer <i>Cui JY, Ma J, Gao XX, Sheng ZM, Pan ZX, Shi LH, Zhang BG</i> ORIGINAL ARTICLE Case Control Study Prognostic utility of gamma-glutamyl transpeptidase to platelet ratio in patients with solitary hepatitis B virus-related hepatocellular carcinoma after hepatectomy <i>Yang CK, Wei ZL, Shen XQ, Jia YX, Wu QY, Wei YG, Su H, Qin W, Liao XW, Zhu GZ, Peng T</i> Retrospective Cohort Study Prognostic prediction models for postoperative patients with stage I to III colorectal cancer based on machine learning <i>Ji XL, Xu S, Li XY, Xu JH, Han RS, Guo YJ, Duan LP, Tian ZB</i>
4565 4579 4597 4614	 REVIEW Unraveling the role of cancer-associated fibroblasts in colorectal cancer <i>Cui JY, Ma J, Gao XX, Sheng ZM, Pan ZX, Shi LH, Zhang BG</i> ORIGINAL ARTICLE Case Control Study Prognostic utility of gamma-glutamyl transpeptidase to platelet ratio in patients with solitary hepatitis B virus-related hepatocellular carcinoma after hepatectomy <i>Yang CK, Wei ZL, Shen XQ, Jia YX, Wu QY, Wei YG, Su H, Qin W, Liao XW, Zhu GZ, Peng T</i> Retrospective Cohort Study Prognostic prediction models for postoperative patients with stage I to III colorectal cancer based on machine learning <i>Ji XL, Xu S, Li XY, Xu JH, Han RS, Guo YJ, Duan LP, Tian ZB</i> Local excision for middle-low rectal cancer after neoadjuvant chemoradiation: A retrospective study from a single tertiary center

Chen N, Li CL, Wang L, Yao YF, Peng YF, Zhan TC, Zhao J, Wu AW



World Journal of Gastrointestinal Oncology Contents Monthly Volume 16 Number 12 December 15, 2024 **Retrospective Study** 4625 Risk factors for hepatocellular carcinoma in cirrhosis: A comprehensive analysis from a decade-long study Zhou DQ, Liu JY, Zhao F, Zhang J, Liu LL, Jia JR, Cao ZH 4636 Prognosis of radiotherapy for esophageal cancer in elderly patients exceeding seventy-five years old Hu LL, Rong F, Liu L, Zhang L, Zhang LL, Yang Q, Xia ZL, Wang H 4650 Nomogram model based on y-glutamyl transferase to albumin ratio predicts survival in hepatocellular carcinoma patients with transarterial chemoembolization treatment Wu ZY, Li H, Chen JL, Su K, Weng ML, Han YW Deep learning model combined with computed tomography features to preoperatively predicting the risk 4663 stratification of gastrointestinal stromal tumors Li Y, Liu YB, Li XB, Cui XN, Meng DH, Yuan CC, Ye ZX 4675 Temozolomide and capecitabine regimen as first-line treatment in advanced gastroenteropancreatic neuroendocrine tumors at a Latin American reference center Cruz-Diaz WE, Paitan V, Medina J, Flores R, Haro-Varas J, Mantilla R, Castro-Oliden V **Basic Study** 4685 Vitamin D 1,25-Dihydroxyvitamin D₃ reduces lipid accumulation in hepatocytes by inhibiting M1 macrophage polarization Luo WJ, Dong XW, Ye H, Zhao QS, Zhang QB, Guo WY, Liu HW, Xu F 4700 Matrine promotes colorectal cancer apoptosis by downregulating shank-associated RH domain interactor expression Zhou YC, Wang QQ, Zhou GYJ, Yin TF, Zhao DY, Sun XZ, Tan C, Zhou L, Yao SK 4716 Enhancing the radiosensitivity of colorectal cancer cells by reducing spermine synthase through promoting autophagy and DNA damage Guo YB, Wu YM, Lin ZZ **META-ANALYSIS** 4728 Efficacy and safety of transhepatic arterial chemoembolization with drug-loaded microspheres in unresectable primary liver cancer Deng J, Mi YH, Xie L, Sun XX, Liu DH, Long HJ, He LY, Wu DH, Shang HC **CASE REPORT** 4738 Mixed pancreatic ductal adenocarcinoma and well-differentiated neuroendocrine tumor: A case report Zhao X, Bocker Edmonston T, Miick R, Joneja U 4746 Signet-ring cell carcinoma of the transverse colon in a 10-year-old girl: A case report Lv L, Song YH, Gao Y, Pu SQ, A ZX, Wu HF, Zhou J, Xie YC



Contents

Monthly Volume 16 Number 12 December 15, 2024

LETTER TO THE EDITOR

4753 Combinations of lenvatinib and immune checkpoint inhibitors plus transarterial chemoembolization, is it the prime time for unresectable hepatocellular carcinoma?

Centrone N, Serrano Uson Junior PL

- 4757 Advancing hepatocellular carcinoma treatment with hepatic arterial infusion chemotherapy Caliskan Yildirim E, Ergun Y
- 4762 Timely identification and treatment of uterine artery pseudoaneurysm after hysteroscopic procedures Byeon H
- 4766 Current efficacy of hepatic arterial infusion chemotherapy in hepatocellular carcinoma Dias E Silva D, Borad M, Uson Junior PLS
- 4770 Use of traditional Chinese medicine bezoars and bezoar-containing preparations in hepatocarcinoma Li DH, Wen QE, Feng RQ, Qiao C, Tian XT
- Crosslink among cyclin-dependent kinase 9, ATP binding cassette transporter G2 and Beclin 1 in colorectal 4778 cancer

Shao ZB, He K, Su YB, Shi Z



World Journal of Gastrointestinal Oncology

Contents

Monthly Volume 16 Number 12 December 15, 2024

ABOUT COVER

Editorial Board Member of World Journal of Gastrointestinal Oncology, Zilvinas Dambrauskas, MD, PhD, Professor, Department of Surgery and Institute for Digestive System Research, Lithuanian University of Health Sciences, Kaunas 50161, Lithuania. zilvinas.dambrauskas@lsmuni.lt

AIMS AND SCOPE

The primary aim of World Journal of Gastrointestinal Oncology (WJGO, World J Gastrointest Oncol) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, etc.

INDEXING/ABSTRACTING

The WJGO is now abstracted and indexed in PubMed, PubMed Central, Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 edition of Journal Citation Reports[®] cites the 2023 journal impact factor (JIF) for WJGO as 2.5; JIF without journal self cites: 2.5; 5-year JIF: 2.8; JIF Rank: 72/143 in gastroenterology and hepatology; JIF Quartile: Q3; and 5-year JIF Quartile: Q2. The WJGO's CiteScore for 2023 is 4.2 and Scopus CiteScore rank 2023: Gastroenterology is 80/167; Oncology is 196/404.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xiang Li; Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL World Journal of Gastrointestinal Oncology	INSTRUCTIONS TO AUTHORS https://www.wjgnet.com/bpg/gerinfo/204		
ISSN	GUIDELINES FOR ETHICS DOCUMENTS		
ISSN 1948-5204 (online)	https://www.wjgnet.com/bpg/GerInfo/287		
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH		
February 15, 2009	https://www.wjgnet.com/bpg/gerinfo/240		
FREQUENCY	PUBLICATION ETHICS		
Monthly	https://www.wjgnet.com/bpg/GerInfo/288		
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT		
Monjur Ahmed, Florin Burada	https://www.wjgnet.com/bpg/gerinfo/208		
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE		
https://www.wjgnet.com/1948-5204/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242		
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS		
December 15, 2024	https://www.wjgnet.com/bpg/GerInfo/239		
COPYRIGHT	ONLINE SUBMISSION		
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com		

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



 \mathcal{O} WŨ

World Journal of **Gastrointestinal** Oncology

Submit a Manuscript: https://www.f6publishing.com

World J Gastrointest Oncol 2024 December 15; 16(12): 4700-4715

DOI: 10.4251/wjgo.v16.i12.4700

ISSN 1948-5204 (online)

ORIGINAL ARTICLE

Basic Study Matrine promotes colorectal cancer apoptosis by downregulating shank-associated RH domain interactor expression

Yuan-Chen Zhou, Qian-Qian Wang, Ge-Yu-Jia Zhou, Teng-Fei Yin, Dong-Yan Zhao, Xi-Zhen Sun, Chang Tan, Lei Zhou, Shu-Kun Yao

Specialty type: Oncology	Yuan-Chen Zhou, Qian-Qian Wang, Chang Tan, Shu-Kun Yao, Graduate School, Peking University China-Japan Friendship School of Clinical Medicine, Beijing 100029, China			
Provenance and peer review: Unsolicited article; Externally peer reviewed.	Ge-Yu-Jia Zhou, Department of Gastroenterology, China-Japan Friendship Hospital (Institute of Clinical Medical Sciences), Beijing 100029, China			
Peer-review model: Single blind	Teng-Fei Yin, Department of Gastroenterology, Qilu Hospital, Shandong University, Jinan 250012, Shandong Province, China			
Peer-review report's classification Scientific Quality: Grade C Novelty: Grade C	Dong-Yan Zhao , Department of Gastroenterology, The First Affiliated Hospital of Soochow University, Suzhou 215006, Jiangsu Province, China			
Creativity or Innovation: Grade B Scientific Significance: Grade B	Xi-Zhen Sun, Department of Gastroenterology, Beijing Jishuitan Hospital, Beijing 100035, China			
P-Reviewer: Kong MW	Lei Zhou, Department of General Surgery, China-Japan Friendship Hospital, Beijing 100029, China			
Received: June 16, 2024				
Revised: September 17, 2024	Shu-Kun Yao, Department of Gastroenterology, China-Japan Friendship Hospital, Beijing			
Accepted: October 16, 2024	100029, China			
Published online: December 15, 2024	Co-corresponding authors: Lei Zhou and Shu-Kun Yao.			
Processing time: 148 Days and 19.7	Corresponding author: Shu-Kun Yao, MD, PhD, Chief Physician, Doctor, Professor, Graduate			
Hours	School, Peking University China-Japan Friendship School of Clinical Medicine, No. 2 Yinghua East Road, Chaoyang District, Beijing 100029, China. shukunyao@126.com			
	Abstract			

Abstract

BACKGROUND

The 5-year survival rate of patients with colorectal cancer (CRC) in China is only 56.9%, highlighting the need for new therapeutic drugs. Previous studies have shown that matrine exhibits antitumor effects by inducing apoptosis. However, the mechanism by which matrine regulates antiapoptotic proteins in CRC remains unclear.

AIM

To identify apoptotic proteins from proteomics and investigate the role of matrine in impeding CRC apoptosis by regulating these proteins.



WJGO | https://www.wjgnet.com

METHODS

Tumor and adjacent normal tissues were collected from 52 patients with CRC who underwent surgery between January and December 2021. Data-independent acquisition quantitative proteomic analysis was performed to identify differentially expressed apoptotic proteins. The selected apoptotic proteins were identified through their association with tumor-node-metastasis (TNM) stage and prognosis, then confirmed by immunohistochemical (IHC) staining in validation cohort. *In vitro*, the role of matrine or apoptotic proteins on cancer cells were analyzed.

RESULTS

Compared to normal tissues, 88 anti-apoptotic proteins from proteomic results were selected. Among them, Shankassociated RH domain interactor (SHARPIN) was identified because of its relationship with TNM stage and overall survival in TCGA database. In the IHC-confirmed cohort, SHARPIN was highly expressed in CRC tissues and localized in the cytoplasm. Higher SHARPIN expression was associated with TNM stage, carbohydrate antigen 153 levels, and gross type compared to low expression. SHARPIN knockdown promoted apoptosis, significantly upregulated the expression of Bcl-2 associated agonist of cell death, Bcl-2 associated X protein, caspase 3, and caspase 8, and downregulated B-cell lymphoma-2 (P < 0.05). Importantly, matrine treatment promoted apoptosis and reversed the proliferation, invasion, and migration of CRC cells by repressing SHARPIN.

CONCLUSION

SHARPIN was identified as an upregulated anti-apoptotic protein in CRC, and matrine exhibited anticancer effects by downregulating its expression. Thus, matrine appears to be a promising drug for CRC.

Key Words: Colorectal cancer; Proteomics; Shank-associated RH domain interactor; Matrine; Apoptosis

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Despite advances in therapy for colorectal cancer (CRC), the 5-year survival rate for CRC patients in China remains only 56.9%. This study explored the effects of matrine on CRC by targeting a newly identified anti-apoptotic protein, Shank-associated RH domain interactor (SHARPIN). SHARPIN was discovered through proteomic analysis and its expression was validated in both the TCGA database and our patient cohort using immunohistochemistry. Inhibiting SHARPIN expression led to increased apoptosis and reduced proliferation, invasion, and migration of CRC cells *in vitro*. Matrine's ability to inhibit SHARPIN and induce apoptosis highlights its potential as a promising therapeutic agent for CRC.

Citation: Zhou YC, Wang QQ, Zhou GYJ, Yin TF, Zhao DY, Sun XZ, Tan C, Zhou L, Yao SK. Matrine promotes colorectal cancer apoptosis by downregulating shank-associated RH domain interactor expression. *World J Gastrointest Oncol* 2024; 16(12): 4700-4715

URL: https://www.wjgnet.com/1948-5204/full/v16/i12/4700.htm **DOI:** https://dx.doi.org/10.4251/wjgo.v16.i12.4700

INTRODUCTION

Colorectal cancer (CRC) is the third most common cancer type worldwide and ranks second in China[1]. It imposes a significant burden on both individuals and the healthcare system because of its high incidence and mortality rate. Despite notable advancements in systemic therapy, CRC prognosis remains poor. In China, although 5-year survival rate increased from 47.2% to 56.9% between 2003 and 2015, it still lagged that of a developed country[2]. The poor prognosis of CRC can be attributed to factors, such as advanced TNM stage, genetic dysregulation or mutation, and drug resistance. Hence, it is imperative to explore novel targets and corresponding drugs to enhance the treatment efficacy.

Cell death resistance is a crucial hallmark of cancer[3]. The dysregulation of apoptotic proteins promotes CRC progression, especially at early-stage, and identified as promising targets for improving therapeutic outcomes. Proteomics offers a means of characterizing protein profiles, including those involved in anti-apoptosis, in CRC tissues. However, there is a paucity of drugs designed to induce apoptosis in the treatment of CRC. Matrine, a natural alkaloid derived from plants such as Sophora flavescens (kushen), has shown anticancer effects without obvious adverse effects. Notably, few studies have explored the effect of matrine on the induction of apoptosis in CRC cells. This study aimed to select antiapoptotic proteins from CRC proteomic profiles and investigate the ability of matrine to induce apoptosis in CRC cells by inhibiting these anti-apoptotic proteins.

Znishideng® WJGO | https://www.wjgnet.com

MATERIALS AND METHODS

Patient recruitment and sample collection

From January 2021 to December 2021, we recruited patients diagnosed with CRC (n = 158) and health control subjects (n = 35) at our hospital. First, 52 paired tumor and adjacent tissues (5 cm away from the tumor) were collected from patients with CRC for proteomic analysis. Four CRC samples were missing, and 9 samples did not contain tumor tissue. Finally, 93 tumor tissues from patients with CRC (Figure 1) and 35 normal tissues from healthy subjects were used as a validation cohort to examine protein expression using immunohistochemistry (IHC). Clean Tissues were transferred by liquid nitrogen and stored in a -80 °C freezer. Other tissues were fixed and processed into sections.

Surgical resection samples diagnosed with CRC were included. Inflammatory bowel disease-associated, hereditary, and neuroendocrine CRC were excluded. Additionally, patients who had received chemoradiotherapy, targeted therapy, immunotherapy, or participated in clinical trials within the past 6 months were also excluded. All enrolled patients provided written informed consent. This study was conducted in accordance with the Declaration of Helsinki and approved by the Clinical Research Ethics Committee of the China-Japan Friendship Hospital (2018-116-K85).

Data independent acquisition analysis

The 52 pairs of CRC tissue samples were analyzed using high-performance liquid chromatography-mass spectrometry/ mass spectrometry to identify differentially expressed proteins (DEPs). Samples were prepared through various processes, including protein denaturation, reduction, alkylation, tryptic digestion, and peptide clean-up, using an iST Sample Preparation Kit (PreOmics, Germany). The eluted peptides were lyophilized using SpeedVac. The peptides were then redissolved in solvent A (0.1% formic acid in water) and analyzed using an Orbitrap Fusion Lumos Tribrid coupled to an EASY-nano LC 1200 system (Thermo Fisher Scientific, MA, United States). The mass spectrometer was run in data independent acquisition (DIA) mode with a hybrid data strategy[4].

Raw DIA data were processed and analyzed using Spectronaut 15.0 (Biognosys AG, Switzerland) with default settings. The spectra were configured to search the UniProt-Homo sapiens database (version201907, 20428 entries), assuming trypsin as the digestion enzyme. The Q value false discovery rate (FDR) cutoff was set to 1% at both the precursor and protein levels. Decoy generation was set to mutated, which is similar to scrambled generation but only applies a random number of amino acid position swamps (min = 2, max = length/2). A normalization strategy was set for local normalization. The average of the top three filtered peptides that passed the 1% Q value cutoff was used to calculate the major group quantities. DEPs were selected if their P_{adj} value was < 0.05, and absolute fold change (FC) > 2 after the *t*-test.

Bioinformatic analysis

Quality control showed a low coefficient of variation and good reliability of the proteomic results: (1) DEPs were analyzed using R packages such as BiocManager, org.Hs.eg.db, colorspace, stringi, and ggplot2, and pairwise comparisons between two groups. The differential protein screening criteria were a *t*-test with Benjamini & Hochberg-corrected *P* value < 0.05 and an absolute FC > 2. Principal component analysis (PCA) was performed to determine the differences between tumor and normal samples. The upregulated and downregulated DEPs are displayed as volcano plots, and hierarchical clustering analysis was performed for expressed proteins and samples simultaneously, with the results displayed as heatmaps; and (2) Functional annotation and enrichment analyses were performed on all identified DEPs, and their functions were marked using Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG).

RNA sequencing quantification and relevant clinical and survival data[5] of CRC tissues (n = 647) and normal colorectal tissues (n = 51) were obtained from TCGA database. (https://portal.gdc.cancer.gov). The ggplot2 and survinier packages were used to visualize the analyzed data.

IHC

Methanal-Fixed and Paraffin-Embedded tumors were stained for the Shank-associated RH domain interactor (SHARPIN) protein. Antigen retrieval was performed using an automatic antigen repair instrument (LBP-5196-II; Guangzhou LBP Medicine Science & Technology Co., Ltd) for 15 minutes in Tris-EDTA buffer (pH 9.0), followed by treatment with an enhanced endogenous peroxidase-blocking buffer (P0100B, Beyotime, Shanghai, China) for 10 minutes. Specimens were incubated overnight at 4 °C with polyclonal antibodies against human SHARPIN (1:100; 14626-1-AP, Proteintech, Wuhan, China). After rinsing three times with phosphate buffered saline (PBS), the slides were incubated with goat anti-rabbit IgG (1:2000, ab6721; Abcam) for 1 hour at room temperature. A coloring reaction was performed using a 3,3N-Diaminobenzidine Tertrahydrochloride chromogenic kit (ZLI-9017; Zhongshan Golden Bridge, Beijing, China). Nuclei were stained with Harris hematoxylin and mounted with neutral gum. Finally, the slides were photographed using a BX53 upright bright-field microscope (Olympus).

Evaluation of IHC staining

The ImageJ software was used for image quantification. SHARPIN expression was quantified using the H-score method in ImageJ; weak immunostaining was assigned a score of 1, moderate immunoreactivity was assigned a score of 2, and strong immunostaining was assigned a score of 3. The percentage of tumor cells stained at each intensity level was multiplied by the respective numeric score of the staining intensity, and the results were summed to quantify the total H-score.

Zaishideng® WJGO | https://www.wjgnet.com



Figure 1 Flow chart. CRC: Colorectal cancer; IHC: Immunohistochemistry.

Data collection

Demographic data, clinicopathological features, and tumor markers, including age, sex, gross type, differentiation, tumor size, tumor location, histopathological type and manifestation, microsatellite instability, TNM stage, carcinoembryonic antigen, carbohydrate antigen 199 (CA199), CA125, CA153, and CA724 were collected from the medical record system.

Cell culture

Two CRC cells, HCT116 and SW480, obtained from the Procell Life Science & Technology Co., Ltd (Wuhan, China), were cultured in McCoy's 5A or DMEM supplemented 10% fetal bovine serum (FBS) (Gibco), 1% penicillin/streptomycin solution (C125C5, NCM Biotech) at 37.7 °C in a humidified atmosphere with 5% CO₂.

Cell transfection and matrine intervention

CRC cells were transfected with SHARPIN small interfering RNA (GenePharma) or negative control siRNA (GenePharma). The sequence pairs of the SHARPIN were as followed: Sense, 5'-CCUGAGGCAGAUCUUCCUATT-3', antisense, 5'-UAGGAAGAUCUGCCUCAGGTT-3'. Lipofectamine 3000 Transfection Reagent (Invitrogen) was used to transfect CRC cells with the above siRNAs, according to the manufacturer's instructions. After 24 hours, the CRC cells were collected for subsequent experiments.

Matrine (MCE) stock solution was diluted with dimethyl sulfoxide and stored in -20 °C in the dark. CRC cells were treated with matrine at concentration of 2.5 mg/mL for 48 hours, which refer to previously described[6].

Quantitative reverse transcription-PCR

Total RNA was extracted from the cells using TRIzol reagent (Vazyme, Nanjing, China). The following primers were used for SHARPIN: Forward 5'- TGTTCTCAGAGCTCGGTTT -3' and reverse 5'- AAGTTCCCCGTCCATCTT -3'. Fluorescence was detected using a CFX Connect instrument (Bio-Rad). Each sample was run in triplicate and compared using GAPDH as the reference gene. Results were analyzed using the $2^{-\Delta CT}$ method for the relative quantification of mRNA expression.

Western blotting

Cell lysis buffer for western blotting, IP (P0013, Beyotime), and a protease and phosphatase inhibitor cocktail (P1045, Beyotime) were used for protein extraction. The BCA Protein Assay Kit (P0012S, Beyotime) was used to measure the amount of protein.

Total proteins were subjected to sodium dodecyl sulfate/polyacrylamide gel electrophoresis and electrotransferred to PVDF membranes (Merck Millipore, China). Nonspecific binding sites were blocked using 5% nonfat milk for 2 hours and the membranes were incubated at 4 °C overnight with SHARPIN Monoclonal antibody (1:1000, Immunoway, United States), Caspase 3/p17/p19 Polyclonal antibody (1:1000, Proteintech), Caspase 8/p43/p18 Polyclonal antibody (1:1000, Proteintech), B-cell lymphoma-2 (Bcl-2) Polyclonal antibody (1:1000, Proteintech), Bcl-2 associated agonist of cell death (BAD) Polyclonal antibody (1:1000, Proteintech), Bcl-2 associated X protein (BAX) Polyclonal antibody (1:1000, Proteintech), GAPDH antibody (1:4000; Abcam), β-actin (1:1000, Proteintech). The PVDF membranes were then treated for 1 hour with a goat anti-rabbit secondary antibody (1:10000; D21109-35, IRDye® 800CW, LI-COR, United States). Fluorescence imaging was performed using an LI-COR Odyssey CLx imaging system. ImageJ software was used to quantify the chemiluminescent signals of the protein bands, using GAPDH, β -actin or tubulin as an internal control.

Flow cytometry analysis

An Annexin V-fluorescein isothiocyanate/propidium iodide apoptosis kit (556547, BD, United States) was used to detect the proportion of apoptotic cells after transfection. All procedures were performed according to manufacturer's instructions. Flow cytometry (FACSCanto II, BD, United States) and FlowJo (V_10.8.1) software were used to determine cell apoptosis rates.



Cell counting kit-8 proliferation assay

The collected CRC cells were seeded in each well of a 96-well plate at a density of 1×10^4 cells/well. Then, at 0 hour, 24 hours, 48 hours and 72 hours, 10 µL of cell counting kit-8 (CCK-8) reagent (C0037, Beyotome) and 90 µL of medium were added after the previous medium was removed. Then, the cells were incubated for 1 hours at 37 °C with 5% CO₂, and the optical density was detected at 450 nm using a microplate reader (Tecan infinite M200, Switzerland).

Transwell invasion assay

CRC cells were suspended in 200 µL of serum-free medium and seeded in the upper chambers coated with matrigel (Corning, United States). The bottom 24-well plates were filled with 600 µL of medium containing 10% FBS as a chemoattractant. After incubation for 72 hours, the cells remaining in the upper chamber were gently removed using cotton swabs, and the cells on the lower surface were photographed and counted after fixing with 4% paraformaldehyde and staining with 0.1% crystal violet.

Wound healing assay

CRC cells were cultured in 6-well plates for 24 hours, and cell transfection was performed as described above. After incubation for 48 hours, a wound was scratched with a 200 µL sterile pipette tip. The cells were then washed with PBS 3 times and incubated in a serum-free medium. Representative images were captured 0 and 24 hours after scratching. The total wound area was measured using the ImageJ software, and the relative migration rate was calculated.

Statistical analysis

Statistical programs for R software (version 4.0.3) and GraphPad Prism (version 9) were used to perform statistical analyses. Quantitative data with a normal distribution are presented as the mean ± SD and were analyzed using Student's t-test or one-way analysis of variance, while variables with a non-normal distribution are presented as the median and interquartile range and were compared using a Mann-Whitney U test or Kruskal-Wallis's test. Categorical variables were analyzed using the χ^2 test or a nonparametric test. Receiver operating characteristic curves were constructed to examine the diagnostic efficiency of RNA molecules by assessing the area under the curve. Statistical significance was set at P <0.05.

RESULTS

Proteomics profiling of CRC patients

52 pairs of cancerous and matched adjacent normal tissues were collected from patients with CRC after surgical treatment for quantitative proteomic detection. 46.2% CRC patients were male, with a mean age of 64.6 ± 15.1 yeas old (Supplementary Table 1). A total of 7666 proteins were identified, and PCA demonstrated a clear separation between the CRC and normal groups (Figure 2A and B). Among these, 1960 DPEs with an FDR of < 1% were identified between the two groups (Figure 2C).

GO and KEGG enrichment analyses were performed for 1960 DEPs. GO enrichment analysis (Figure 2D) revealed the significant enrichment of various biological processes, molecular functions, and cellular components. In the biological processes, enrichment was observed in response to stimuli (oxygen-containing compounds, oxidative stress, and corticosteroids), RNA metabolic processes (ncRNA metabolic processes, rRNA metabolic processes, etc.), and other processes. In terms of molecular functions, enrichment was noted for binding (proteins, ions, etc.) and catalytic activity (acting on RNA). Among the cellular components, enrichment was found in organelles (chromosomal part, nucleolus, etc.), extracellular regions (extracellular space, extracellular matrix, etc.), and cell junctions (anchoring junction, adheres junction, etc.). KEGG pathway enrichment analysis (Figure 2E) indicated the activation of several pathways, including the PI3K-Akt signaling pathway, focal adhesion, cell cycle, and extracellular matrix-receptor interaction.

The enrichment of responses to stimuli in biological processes is often associated with cell death. Resistance to cell death is a crucial hallmark of cancer and the upregulation of anti-apoptotic proteins is known to promote tumor progression.

Selecting anti-apoptotic proteins in tissue proteomics profiling and TCGA database of CRC patients

To identify these anti-apoptotic proteins, 88 proteins were selected by negative regulation of apoptotic proteins based on GO enrichment analysis and were upregulated in tumor tissue (Supplementary Table 2). SHARPIN, metalloproteinase inhibitor 1 (TIMP1), nucleolar complex protein 2 homolog (NOC2 L), and DNA ligase 4 (LIG4) were significantly different among the TNM stages (Supplementary Table 3). As shown in Figure 3A, the expression of SHARPIN was significantly higher in stage III than in stage I. TIMP1 and NOC2 L expression levels were significantly higher in stage III than in stage II. With the advancement of TNM staging, the expression of SHARPIN was gradually increased, but not that of the others.

SHARPIN, TIMP1, NOC2 L, and LIG4 were further confirmed using TCGA database. As shown in Figure 3B, expect for LIG4, the expression of SHARPIN, TIMP1, NOC2 L was higher in tumor tissues than in normal tissues. As shown in Figure 3C, only SHARPIN was associated with the TNM stage. In the survival analysis, higher expression of SHARPIN and TIMP1 significantly decreased the overall survival (Figure 3D). Thus, upregulation SHARPIN was associated with the pathological stage and prognosis of CRC.



WJGO https://www.wjgnet.com





Zhou YC et al. Matrine impedes CRC by repressing SHARPIN





Figure 2 Proteomic profiles and bioinformatic analysis of colorectal cancer patients. A: Hierarchical cluster analysis shows the global differentially expressed proteins. Upregulated (red) and downregulated (green) differentially expressed proteins in the CRC group compared to the normal group; B: Principal component analysis shows the protein expression in the cancer tissues (green) and normal tissues (orange) of colorectal cancer (CRC) patients; C: Volcano plot shows the fold changes of identified proteins. Significantly upregulated or downregulated proteins are represented in red and blue, respectively, while proteins without significant changes are shown in gray; D: Gene Ontology analysis shows the enrichment results for differentially expressed proteins; E: Kyoto encyclopedia of genes and genomes analysis shows that differentially expressed proteins are significantly enriched in multiple pathways, °*P* < 0.001. CRC: Colorectal cancer; PCA: Principal component analysis; HCA: Hierarchical cluster analysis; GO: Gene ontology; KEGG: Kyoto encyclopedia of genes and genomes; PI3K-Akt: Phosphatidylinositol 3-kinase serine/threonine kinase; ECM: Extracellular matrix.

Figure 3 The expression of shank-associated RH domain interactor, metalloproteinase inhibitor 1, nucleolar complex protein 2 homolog, and DNA ligase 4 proteins and its association with tumor-node-metastasis and overall survival in colorectal cancer. A: The expression of 4 proteins among tumor-node-metastasis (TNM) stages in 52 colorectal cancer (CRC) patients; B: The expression of 4 proteins in TCGA; C: The expression of 4 proteins among TNM stages in TCGA; D: The association between expression of 4 proteins and overall survival in TCGA; E: The immunohistochemistry of shank-associated RH domain interactor in 93 CRC tissues and 35 normal tissues. $^{a}P < 0.05$; $^{b}P < 0.01$; CRC: Colorectal cancer; HC: Health control; SHARPIN: Shank-associated RH domain interactor; TIMP1: Metalloproteinase inhibitor 1; NOC2 L: Nucleolar complex protein 2 homolog; LIG4: DNA ligase 4.

Zaishideng® WJGO | https://www.wjgnet.com

December 15, 2024 Volume 16 Issue 12

High expression of SHARPIN and its association with clinicopathological characteristics in CRC

IHC was conducted to investigate the expression of SHARPIN between CRC tumor tissues (n = 93) and normal tissues (n = 35). Compared with normal tissues (Figure 3E), CRC tissues showed significantly higher SHARPIN expression (P < 0.01). SHARPIN was hardly stained in normal tissues but was highly expressed in the cytoplasm of CRC cells.

93 CRC patients were divided into two groups according to their mean H-score. As indicated in Table 1, the average age was 67.5 ± 10.9 years, and 49 participants (52.7%) were male. SHARPIN expression correlated with serum CA153 Levels, gross type, T stage, N stage, and TNM stage (all *P* < 0.05; Table 1).

Downregulated SHARPIN induced CRC cells apoptosis

To further investigate the effects of SHARPIN on the apoptosis of two CRC cell lines (HCT116 and SW480), siRNA was used to knockdown the expression of SHARPIN. The efficiency of SHARPIN knockdown was confirmed by Quantitative reverse transcription-PCR and western blot analysis (Supplementary Figure 1). Apoptotic cells were assessed by flow cytometry, and the expression of apoptotic proteins (Caspase 3, caspase 8, BAD, BAX, and Bcl-2) was determined by western blotting. The results showed that SHARPIN knockdown promoted apoptosis in CRC cells, as indicated by the increased proportion of apoptotic cells in the siSHARPIN group compared to that in the control group (Figure 4A; P < 0.05). Additionally, the expression levels of Caspase 3, Caspase 8, BAD, and BAX were significantly upregulated in the siSHARPIN group, whereas the expression level of Bcl-2 was downregulated in siSHARPIN + Matrine group (Figure 4B).

Matrine induced apoptosis of CRC cells by downregulating SHARPIN

Subsequently, we explored the role of SHARPIN in suppressing matrine-induced apoptosis in CRC cells. We observed the apoptosis-promoting effects of matrine in CRC cells. In CRC cell lines treated with matrine, there was a significant downregulation of SHARPIN expression (Figure 4C) and significant increase in the number of apoptotic cells (Figure 4A). The siSHARPIN + matrine group enhanced this effect. As shown in Figure 4, caspase 3 was significantly upregulated and Bcl-2 was significantly downregulated in the matrine group. Furthermore, in the siSHARPIN + matrine group, the expression of caspase 3, caspase 8, BAD, and BAX was significantly upregulated, whereas that of Bcl-2 was significantly downregulated.

Matrine inhibited progression of CRC cells by downregulating SHARPIN

Next, we examined the effects of matrine and SHARPIN knockdown on the proliferation, invasion, and migration of CRC cells. The viability of CRC cells was assessed using the CCK-8 assay, whereas cell invasion and migration were evaluated using transwell invasion and wound-healing assays, respectively.

The viability of CRC cells was significantly inhibited in the siSHARPIN, matrine, and siSHARPIN + matrine groups compared to that in the control group (P < 0.01; Figure 5A). In the transwell invasion assay, the number of invasive cells markedly decreased in the siSHARPIN, matrine, and siSHARPIN + matrine groups (Figure 5B). Furthermore, the wound healing rates of CRC cells were significantly reduced in the siSHARPIN, matrine, and s

DISCUSSION

The incidence and mortality rates of CRC are on the rise, and despite advances in treatment strategies, improvements in prognosis have been modest. Resistance to cell death is a critical hallmark of cancer, and there are relatively few drugs specifically designed to induce apoptosis in cancer cells. Apoptotic drugs hold promise for tumor treatment, and further exploration of molecules involved in regulating the apoptotic pathway is essential for advancing CRC therapies. In the present study, SHARPIN was both found to be upregulated in CRC tissues based on our proteomic and TCGA database and validated by IHC staining in our CRC cohort. Additionally, the expression of SHARPIN was associated with different TNM stages and a poor prognosis. *In vitro* experiments, this study demonstrated that the downregulation of SHARPIN induces apoptosis in CRC cells. Matrine treatment also induced apoptosis and inhibited the proliferation, invasion, and migration of CRC cells by downregulating SHARPIN expression. Therefore, the role of SHARPIN in CRC was initially validated in our study, providing further evidence supporting the potential of matrine in preventing CRC and colorectal adenoma. Our study identifies SHARPIN as a potential anti-apoptotic molecule, shedding light on the mechanisms of matrine's impact on CRC. Moreover, matrine has a favorable safety profile with minimal side effects. In the future, it is interesting to explore the effect of matrine in preventing tumor progression in high-risk patients.

SHARPIN is enriched at synaptic sites in mature neurons[7]. Located in the cytoplasm, SHARPIN is an approximately 40 kDa conserved protein that is ubiquitously expressed in various human tissues[8]. Furthermore, SHARPIN is amplified in some cancers, such as prostate, breast, and ovarian cancers. Zhang *et al*[9] demonstrated that upregulated SHARPIN activated the NF- κ B pathway and suppressed cell apoptosis, potentially contributing to prostate cancer progression[9]. Overexpression of SHARPIN has been shown to increase the breast cancer risk[10] and promote the breast cancer progression by regulating *p*53 and *ERa*[11]. Knockdown of SHARPIN expression leads to apoptosis and inhibits the proliferation, invasion, and migration of ovarian cancer cells[12]. However, the role of SHAPRIN in CRC has not yet been reported. In the present study, SHARPIN expression was significantly higher in tumor tissues than in normal tissues in our proteomic data (*n* = 52), TCGA database (*n* = 698), and IHC-confirmed cohort (*n* = 93). Additionally, higher SHARPIN expression was associated with poor overall survival in CRC patients compared to that in the low expression group.

WJGO https://www.wjgnet.com

Figure 4 Matrine induces apoptosis and regulates apoptotic proteins by downregulating shank-associated RH domain interactor. A: The apoptosis rate of colorectal cancer (CRC) cells in 4 groups [control group (con group), SHARPIN knockdown group (siSHARPIN group), matrine, siSHARPIN + matrine]; B: The expression of caspase 3, caspase 8, B-cell lymphoma-2 (Bcl-2) associated X protein, Bcl-2 associated agonist of cell death, Bcl-2 proteins in CRC cells in 4 groups; C: The suppressing effect of matrine on shank-associated RH domain interactor (SHARPIN) in CRC cells. ^aP < 0.05; ^bP < 0.01; ^cP < 0.05 vs con, ^dP < 0.01 vs con; *P < 0.05 vs siSHARPIN, *P < 0.01 vs siSHARPIN; *P < 0.05 vs matrine, *P < 0.01 vs matrine; Con: Control group; siSHARPIN: SHARPIN knockdown group; matrine: Matrine group; siSHARPIN + matrine: Matrine and SHARPIN knockdown group; CRC: Colorectal cancer; SHARPIN: Shank-associated RH domain interactor; Bcl-2: B-cell lymphoma-2; BAD: Bcl-2 associated agonist of cell death; BAX: Bcl-2 associated X protein.

Baisbideng® WJGO https://www.wjgnet.com

Figure 5 Matrine impedes proliferation, invasion and migration in colorectal cancer by downregulating shank-associated RH domain interactor interactor. A: The viability of colorectal cancer (CRC) cells using the cell counting kit-8 assay in 4 groups [control (con), si-Shank-associated RH domain interactor (SHARPIN), matrine, siSHARPIN + matrine]; B: The invasion cells of CRC in 4 groups; C: The wound healing of CRC cells in 4 groups. $^{o}P < 0.05 vs \operatorname{con}$, $^{d}P < 0.01 vs \operatorname{con}$; $^{e}P < 0.05 vs \operatorname{siSHARPIN}$, $^{f}P < 0.01 vs \operatorname{siSHARPIN}$; $^{g}P < 0.05 vs \operatorname{matrine}$; b: The invasion cells of CRC in 4 group; C: The wound healing of CRC cells in 4 groups. $^{o}P < 0.05 vs \operatorname{con}$, $^{d}P < 0.01 vs \operatorname{con}$; $^{e}P < 0.05 vs \operatorname{siSHARPIN}$; $^{f}P < 0.01 vs \operatorname{matrine}$; b: The invasion cells of CRC in 4 group; C: The wound healing of CRC cells in 4 group; SHARPIN: Shank-associated RH domain interactor; siSHARPIN; SHARPIN, $^{f}P < 0.01 vs \operatorname{matrine}$; $^{h}P < 0.01 vs \operatorname{matrine}$; Con: Control group; SHARPIN: Shank-associated RH domain interactor; siSHARPIN: SHARPIN knockdown group; matrine: Matrine group; siSHARPIN + matrine: Matrine and SHARPIN knockdown group; CRC: Colorectal cancer.

In a 14-year-old boy, a homozygous deficiency of SHARPIN led to a shift in signaling from pro-survival to cell death in immune cells, contributing to an observed discrepancy in immune function[13]. Because of the effect of SHARPIN deficiency on cell death, knockdown expression of SHARPIN may induce apoptosis in cancer. As the component of the linear ubiquitin chain activation complex, SHARPIN plays an important role in inflammatory response and apoptosis by regulating canonical NF-κB signaling[8,14,15]. For anti-tumor effect, NF-kB pathway was activated by SHARPIN overex-pression with upregulated anti-apoptosis proteins in prostate cancer[9,16]. As a *p*53 mediator, SHARPIN may facilitate its degradation or ubiquitination in a mouse double minute 2 homolog-dependent manner[17], which is an inhibitor of *p*53 transcriptional activation[18]. SHARPIN is a negative regulator of PTEN in human tumor cell lines and primary cervical cancer cells both *in vitro* and *in vivo*. SHARPIN activates the PI3K/AKT pathway and induces tumorigenesis by interacting with PTEN.

Matrine is an alkaloid isolated from the traditional Chinese medicine, Sophora flavescens Aiton. In China, matrine injections are used to treat hepatitis. Owing to its anti-inflammatory effects, matrine also exhibits antitumor properties, especially apoptosis. Previous studies have shown that matrine induces autophagy and apoptosis by downregulating β -catenin and mediating the JNK-Bcl-2/Bcl-xL-Bax/Bak pathway in hepatocellular carcinoma cells[19,20]. In CRC cells, matrine induces apoptosis by regulating associated proteins. Zhang *et al*[21] demonstrated matrine reduce the ratio of Bcl-2/Bax and increase caspase-9 *in vitro*. Gu *et al*[22] explored effects of matrine on Bcl-2, Bax and caspase-3 in several cell lines. And Chang *et al*[23] showed that matrine activated apoptosis by regulating BAX,Bcl-2 and Cyto C. However, the precise mechanism underlying the matrine-induced apoptosis in CRC cells remains unclear. In the current study, we found that matrine induced cell apoptosis by inhibiting SHARPIN expression, while also repressing the proliferation, invasion, and migration of CRC cells. Targeted SHARPIN drugs might lead to various adverse effects, while matrine

WJGO https://www.wjgnet.com

Table 1 Clinicalpathological charate	eristics of colorectal cance	er according to Shank-ass	ociated RH domain interac	tor expression, <i>n</i> (%)
Variables	Total (<i>n</i> = 93)	Low (<i>n</i> = 50)	High (<i>n</i> = 43)	P value
Sex, male	49 (52.7)	28 (56)	21 (48.8)	0.49
Age, years	67.5 ± 10.9	67.4 ± 10.6	67.7 ± 11.4	0.89
Location				0.38
Left	75 (80.6)	42 (84)	33 (76.7)	
Right	18 (19.4)	8 (16)	10 (23.3)	
Gross type				0.03
Fungating	32 (34.4)	21 (42)	11 (25.6)	
Ulceroinfiltrative	54 (58.1)	23 (46)	31 (72.1)	
Ulcerofungating	7 (7.5)	6 (12)	1 (2.3)	
Differentiation				0.443
Poor	10 (11.1)	3 (10.3)	4 (12.5)	
Moderate	70 (77.8)	21 (72.4)	26 (81.2)	
Well	10 (11.1)	5 (17.2)	2 (6.2)	
Tumor size, ≥ 5 cm	48 (51.6)	24 (48)	24 (55.8)	0.45
Histopathologic type				0.76
AC	79 (84.9)	43 (86)	36 (83.7)	
MAC	14 (15.1)	7 (14)	7 (16.3)	
Venous invasion	21 (22.6)	10 (20)	11 (25.6)	0.52
Perineural invasion	48 (51.6)	24 (48)	24 (55.8)	0.45
MSI	7 (7.5)	3 (6)	4 (9.3)	0.70
T stage				0.02
1	4 (4.3)	4 (8)	0 (0)	
2	17 (18.3)	13 (26)	4 (9.3)	
3	60 (64.5)	29 (58)	31 (72.1)	
4	12 (12.9)	4 (8)	8 (18.6)	
N stage				0.02
0	63 (67.7)	40 (80)	23 (53.5)	
1	18 (19.4)	7 (14)	11 (25.6)	
2	12 (12.9)	3 (6)	9 (20.9)	
M stage				0.09
0	87 (93.5)	49 (98)	38 (88.4)	
1	6 (6.5)	1 (2)	5 (11.6)	
TNM stage				0.01
Ι	20 (21.5)	16 (32)	4 (9.3)	
П	41 (44.1)	22 (44)	19 (44.2)	
III	26 (28.0)	11 (22)	15 (34.9)	
IV	6 (6.5)	1 (2)	5 (11.6)	
CA125, ng/mL	9.0 (7.0, 14.5)	9.0 (7.0, 14.5)	9.0 (7.1, 13.7)	0.96
CA199, U/mL	14.1 (9.2, 24.0)	14.5 (8.0, 23.0)	13.6 (10.0, 26.0)	0.97
CA724, U/mL	1.7 (1.5, 4.7)	1.9 (1.5, 4.9)	1.6 (1.5, 2.8)	0.61
CEA, ng/mL	4.8 (2.5, 11.1)	5.5 (2.6, 11.5)	3.5 (2.5, 8.8)	0.26

CA153, U/mL	8.1 (6.0, 12.4)	7.2 (5.3, 10.6)	9.2 (6.9, 13.3)	0.04

Two groups were divided the mean of the H-score. AC: Adenocarcinoma; MAC: Mucinous adenocarcinoma; MSI: Microsatellite instability; CEA Carcinoembryonic antigen; CA199: Carbohydrate antigen 19-9; CA724: Carbohydrate antigen 724; TNM: Tumor-node-metastasis.

shows potential in multiple therapeutic roles, making it a promising candidate for treatment.

Our study has some limitations. First, the follow-up time of the enrolled 52 patients was not available, and the association between SHARPIN expression and CRC survival needs to be investigated in Chinese population. Second, the therapeutic effects of matrine on CRC need to be validated *in vivo* and in clinical trials. Matrine injection has been used to improve hepatitis; however, the dosage of matrine that inhibits tumor progression remains to be explored. Third, the mechanism of matrine repressing SHARPIN expression remains to be conducted in the further study.

CONCLUSION

In conclusion, this study identified overexpressed SHARPIN in CRC tissues. High expression SHARPIN is associated with TNM stage and poor prognosis. Matrine promotes cell apoptosis and inhibits proliferation, invasion, and migration by repressing SHARPIN expression in CRC. Thus, matrine could be considered as a novel therapeutic agent for CRC.

ACKNOWLEDGEMENTS

The authors thank Wei-Liang Sun (China-Japan Friendship Hospital, Beijing, PhD) and Tian-Hui Zhou (Graduate School of Beijing University of Chinese Medicine, Beijing University of Chinese Medicine, MD) for their technical assistance.

FOOTNOTES

Author contributions: Zhou YC conceived and designed the study, performed sample collection, experiments, acquired and analyzed data, and prepared the original draft; Wang QQ, Zhou GY, Zhao DY, Yin TF, Tan C, and Sun XZ participated in sample acquisition, data analysis, and manuscript revision; All authors read and approved the final manuscript. Both Yao SK and Zhou L have played important and indispensable roles in the experimental design, data interpretation and manuscript preparation as the co-corresponding authors. Yao SK applied for and obtained the funds for this research project.

Supported by National Key Development Plan for Precision Medicine Research, No. 2017YFC0910002.

Institutional review board statement: The study was reviewed and approved by the Clinical Research Ethics Committee of the China-Japan Friendship Hospital (No. 2018-116-K85).

Conflict-of-interest statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as potential conflicts of interest.

Data sharing statement: Technical appendix, statistical code, and dataset available from the corresponding author at shukunyao@126. com.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country of origin: China

ORCID number: Yuan-Chen Zhou 0000-0001-6024-6246; Qian-Qian Wang 0000-0002-7709-2121; Ge-Yu-Jia Zhou 0000-0002-4196-1436; Teng-Fei Yin 0000-0003-1140-8637; Dong-Yan Zhao 0000-0002-7026-068X; Xi-Zhen Sun 0000-0001-9967-5726; Chang Tan 0000-0002-0969-1851; Shu-Kun Yao 0000-0002-8512-2589.

S-Editor: Li L L-Editor: A P-Editor: Zhao S

Raishideng® WJGO | https://www.wjgnet.com

REFERENCES

- Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* 2021; 71: 209-249 [PMID: 33538338 DOI: 10.3322/caac.21660]
- 2 Zeng H, Chen W, Zheng R, Zhang S, Ji JS, Zou X, Xia C, Sun K, Yang Z, Li H, Wang N, Han R, Liu S, Li H, Mu H, He Y, Xu Y, Fu Z, Zhou Y, Jiang J, Yang Y, Chen J, Wei K, Fan D, Wang J, Fu F, Zhao D, Song G, Chen J, Jiang C, Zhou X, Gu X, Jin F, Li Q, Li Y, Wu T, Yan C, Dong J, Hua Z, Baade P, Bray F, Jemal A, Yu XQ, He J. Changing cancer survival in China during 2003-15: a pooled analysis of 17 population-based cancer registries. *Lancet Glob Health* 2018; 6: e555-e567 [PMID: 29653628 DOI: 10.1016/S2214-109X(18)30127-X]
- 3 Hanahan D. Hallmarks of Cancer: New Dimensions. *Cancer Discov* 2022; 12: 31-46 [PMID: 35022204 DOI: 10.1158/2159-8290.CD-21-1059]
- 4 Prakash A, Peterman S, Ahmad S, Sarracino D, Frewen B, Vogelsang M, Byram G, Krastins B, Vadali G, Lopez M. Hybrid data acquisition and processing strategies with increased throughput and selectivity: pSMART analysis for global qualitative and quantitative analysis. J Proteome Res 2014; 13: 5415-5430 [PMID: 25244318 DOI: 10.1021/pr5003017]
- 5 Liu J, Lichtenberg T, Hoadley KA, Poisson LM, Lazar AJ, Cherniack AD, Kovatich AJ, Benz CC, Levine DA, Lee AV, Omberg L, Wolf DM, Shriver CD, Thorsson V; Cancer Genome Atlas Research Network, Hu H. An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. *Cell* 2018; **173**: 400-416.e11 [PMID: 29625055 DOI: 10.1016/j.cell.2018.02.052]
- **Zhang X**, Xu H, Bi X, Hou G, Liu A, Zhao Y, Wang G, Cao X. Src acts as the target of matrine to inhibit the proliferation of cancer cells by regulating phosphorylation signaling pathways. *Cell Death Dis* 2021; **12**: 931 [PMID: 34642304 DOI: 10.1038/s41419-021-04221-6]
- 7 Seymour RE, Hasham MG, Cox GA, Shultz LD, Hogenesch H, Roopenian DC, Sundberg JP. Spontaneous mutations in the mouse Sharpin gene result in multiorgan inflammation, immune system dysregulation and dermatitis. *Genes Immun* 2007; 8: 416-421 [PMID: 17538631 DOI: 10.1038/sj.gene.6364403]
- Wang Z, Potter CS, Sundberg JP, Hogenesch H. SHARPIN is a key regulator of immune and inflammatory responses. *J Cell Mol Med* 2012; 16: 2271-2279 [PMID: 22452937 DOI: 10.1111/j.1582-4934.2012.01574.x]
- 9 Zhang Y, Huang H, Zhou H, Du T, Zeng L, Cao Y, Chen J, Lai Y, Li J, Wang G, Guo Z. Activation of nuclear factor κB pathway and downstream targets survivin and livin by SHARPIN contributes to the progression and metastasis of prostate cancer. *Cancer* 2014; **120**: 3208-3218 [PMID: 24925528 DOI: 10.1002/cncr.28796]
- 10 De Melo J, Tang D. Elevation of SIPL1 (SHARPIN) Increases Breast Cancer Risk. *PLoS One* 2015; 10: e0127546 [PMID: 25992689 DOI: 10.1371/journal.pone.0127546]
- 11 **Tian Z**, Tang J, Yang Q, Li X, Zhu J, Wu G. Atypical ubiquitin-binding protein SHARPIN promotes breast cancer progression. *Biomed Pharmacother* 2019; **119**: 109414 [PMID: 31518875 DOI: 10.1016/j.biopha.2019.109414]
- 12 Wang G, Zhuang Z, Cheng J, Yang F, Zhu D, Jiang Z, Du W, Shen S, Huang J, Hua L, Chen Y. Overexpression of SHARPIN promotes tumor progression in ovarian cancer. *Exp Mol Pathol* 2022; 104806 [PMID: 35798064 DOI: 10.1016/j.yexmp.2022.104806]
- 13 Oda H, Manthiram K, Chavan PP, Rieser E, Veli Ö, Kaya Ö, Rauch C, Nakabo S, Kuehn HS, Swart M, Wang Y, Çelik NI, Molitor A, Ziaee V, Movahedi N, Shahrooei M, Parvaneh N, Alipour-Olyei N, Carapito R, Xu Q, Preite S, Beck DB, Chae JJ, Nehrebecky M, Ombrello AK, Hoffmann P, Romeo T, Deuitch NT, Matthíasardóttir B, Mullikin J, Komarow H, Stoddard J, Niemela J, Dobbs K, Sweeney CL, Anderton H, Lawlor KE, Yoshitomi H, Yang D, Boehm M, Davis J, Mudd P, Randazzo D, Tsai WL, Gadina M, Kaplan MJ, Toguchida J, Mayer CT, Rosenzweig SD, Notarangelo LD, Iwai K, Silke J, Schwartzberg PL, Boisson B, Casanova JL, Bahram S, Rao AP, Peltzer N, Walczak H, Lalaoui N, Aksentijevich I, Kastner DL. Biallelic human SHARPIN loss of function induces autoinflammation and immunodeficiency. *Nat Immunol* 2024; 25: 764-777 [PMID: 38609546 DOI: 10.1038/s41590-024-01817-w]
- 14 Tokunaga F, Nakagawa T, Nakahara M, Saeki Y, Taniguchi M, Sakata S, Tanaka K, Nakano H, Iwai K. SHARPIN is a component of the NFκB-activating linear ubiquitin chain assembly complex. *Nature* 2011; **471**: 633-636 [PMID: 21455180 DOI: 10.1038/nature09815]
- 15 Ikeda F, Deribe YL, Skånland SS, Stieglitz B, Grabbe C, Franz-Wachtel M, van Wijk SJ, Goswami P, Nagy V, Terzic J, Tokunaga F, Androulidaki A, Nakagawa T, Pasparakis M, Iwai K, Sundberg JP, Schaefer L, Rittinger K, Macek B, Dikic I. SHARPIN forms a linear ubiquitin ligase complex regulating NF-κB activity and apoptosis. *Nature* 2011; **471**: 637-641 [PMID: 21455181 DOI: 10.1038/nature09814]
- 16 Li J, Lai Y, Cao Y, Du T, Zeng L, Wang G, Chen X, Chen J, Yu Y, Zhang S, Zhang Y, Huang H, Guo Z. SHARPIN overexpression induces tumorigenesis in human prostate cancer LNCaP, DU145 and PC-3 cells *via* NF-κB/ERK/Akt signaling pathway. *Med Oncol* 2015; **32**: 444 [PMID: 25550157 DOI: 10.1007/s12032-014-0444-3]
- 17 Yang H, Yu S, Wang W, Li X, Hou Y, Liu Z, Shi Y, Mu K, Niu G, Xu J, Wang H, Zhu J, Zhuang T. SHARPIN Facilitates p53 Degradation in Breast Cancer Cells. *Neoplasia* 2017; 19: 84-92 [PMID: 28063307 DOI: 10.1016/j.neo.2016.12.002]
- 18 Lessel D, Wu D, Trujillo C, Ramezani T, Lessel I, Alwasiyah MK, Saha B, Hisama FM, Rading K, Goebel I, Schütz P, Speit G, Högel J, Thiele H, Nürnberg G, Nürnberg P, Hammerschmidt M, Zhu Y, Tong DR, Katz C, Martin GM, Oshima J, Prives C, Kubisch C. Dysfunction of the MDM2/p53 axis is linked to premature aging. *J Clin Invest* 2017; **127**: 3598-3608 [PMID: 28846075 DOI: 10.1172/JCI92171]
- Xie BS, He XX, Ai ZL, Yao SK. Involvement of β-catenin in matrine-induced autophagy and apoptosis in WB-F344 cells. *Mol Med Rep* 2014;
 9: 2547-2553 [PMID: 24718323 DOI: 10.3892/mmr.2014.2107]
- 20 Yang J, Yao S. JNK-Bcl-2/Bcl-xL-Bax/Bak Pathway Mediates the Crosstalk between Matrine-Induced Autophagy and Apoptosis *via* Interplay with Beclin 1. *Int J Mol Sci* 2015; 16: 25744-25758 [PMID: 26516844 DOI: 10.3390/ijms161025744]
- 21 Zhang S, Cheng B, Li H, Xu W, Zhai B, Pan S, Wang L, Liu M, Sun X. Matrine inhibits proliferation and induces apoptosis of human colon cancer LoVo cells by inactivating Akt pathway. *Mol Biol Rep* 2014; **41**: 2101-2108 [PMID: 24452711 DOI: 10.1007/s11033-014-3059-z]
- Gu YY, Chen MH, May BH, Liao XZ, Liu JH, Tao LT, Man-Yuen Sze D, Zhang AL, Mo SL. Matrine induces apoptosis in multiple colorectal cancer cell lines in vitro and inhibits tumour growth with minimum side effects *in vivo via* Bcl-2 and caspase-3. *Phytomedicine* 2018; **51**: 214-225 [PMID: 30466620 DOI: 10.1016/j.phymed.2018.10.004]
- 23 Chang C, Liu SP, Fang CH, He RS, Wang Z, Zhu YQ, Jiang SW. Effects of matrine on the proliferation of HT29 human colon cancer cells and its antitumor mechanism. *Oncol Lett* 2013; **6**: 699-704 [PMID: 24137393 DOI: 10.3892/ol.2013.1449]

Raishideng® WJGO | https://www.wjgnet.com

Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

