



High cost of silence: Needlestick injuries and the persistence of hepatitis B virus risk among healthcare trainees

Janaina Luz Narciso-Schiavon

Peer review: Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific quality: Grade A, Grade C, Grade C, Grade C

Novelty: Grade A, Grade C, Grade C, Grade C

Creativity or innovation: Grade A, Grade C, Grade C, Grade C

Scientific significance: Grade A, Grade C, Grade C, Grade C

P-Reviewer: Said ZNANA, PhD, Professor, Professor Emeritus, Egypt; Zhu J, Academic Fellow, MD, China

Received: April 29, 2026

Revised: July 18, 2026

Accepted: August 25, 2026

Published online: September 25, 2026

Processing time: 144 Days and 7.9 Hours

Copyright: ©Author(s) 2026. This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution-NonCommercial \(CC BY-NC 4.0\)](#) license. No commercial re-use. See [permissions](#). Published by Baishideng Publishing Group Inc.



Janaina Luz Narciso-Schiavon, Department of Internal Medicine, Gastroenterology Division, Federal University of Santa Catarina, Florianópolis 88040-900, Santa Catarina, Brazil

ORCID number: Janaina Luz Narciso-Schiavon [0000-0002-6228-4120](https://orcid.org/0000-0002-6228-4120).

Corresponding author: Janaina Luz Narciso-Schiavon, MD, PhD, Associate Professor, Department of Internal Medicine, Gastroenterology Division, Federal University of Santa Catarina, Florianópolis 88040-900, Santa Catarina, Brazil. janaina.narciso@uol.com.br

Abstract

The study by Jaho *et al* published in the *World Journal of Virology* highlights a compelling paradox in occupational health: While a robust 86.6% of 134 Albanian health sciences students demonstrate seroprotection (antibody to hepatitis B surface antigen ≥ 10 IU/L), a staggering 64.5% report at least one needlestick injury during training. This high prevalence of exposure is compounded by a culture of silence, with one-third of incidents remaining unreported. This editorial analyzes why high seroprotection rates should not lead to institutional complacency. It argues that the specific vulnerability observed among midwifery students and the lack of systematic post-vaccination serological testing represent critical system gaps. Instead of relying solely on historical vaccination records, institutions should adopt guideline-based strategies for assessing hepatitis B immune status before clinical exposure, while fostering non-punitive reporting pathways. Bridging the gap between immunological protection and clinical behaviour is essential to safeguard those at the frontline of care.

Key Words: Hepatitis B; Students; Health occupations; Health personnel; Needlestick injuries; Occupational exposure; Serologic tests; Vaccination; Hepatitis B vaccines

Core Tip: High hepatitis B virus seroprotection rates among healthcare trainees may create a false sense of safety. The study by Jaho *et al* shows that protective antibody levels (antibody to hepatitis B surface antigen) often coexist with frequent needlestick injuries and underreporting. Protection in clinical training is not only biological but also procedural. It requires verification of immune status before exposure and clear, accessible post-exposure pathways. Without these systems, a meaningful minority of trainees remains at risk despite apparent vaccination success.

Citation: Narciso-Schiavon JL. High cost of silence: Needlestick injuries and the persistence of hepatitis B virus risk among healthcare trainees. *World J Virol* 2026; 15(3): 122794

URL: <https://www.wjgnet.com/2220-3249/full/v15/i3/122794.htm>

DOI: <https://dx.doi.org/10.5501/wjv.122794>

This editorial refers to "Serological evidence of hepatitis B virus exposure and vaccination coverage trends in health sciences students in Albania" by Jaho et al, 2026; <https://doi.org/10.5501/wjv.v15.i2.118992>.

INTRODUCTION

The study by Jaho et al[1] published in the *World Journal of Virology* evaluated 134 Albanian health sciences students (mean age 23.8 ± 5.7 years). As Albania implemented universal hepatitis B vaccination by 2000[2], most participants were likely vaccinated during childhood. Nevertheless, the study breaks the false sense of security that permeates many educational institutions[1]. As expected, 86.6% of students had protective antibody to hepatitis B surface antigen (anti-HBs) levels (≥ 10 IU/L), and most (76.1%) had titers > 100 IU/L, indicating a robust vaccine-induced immune response. In addition, the absence of circulating hepatitis B surface antigen (HBsAg) suggests a controlled epidemiological situation. However, the same cohort reported frequent needlestick and sharps injuries, showing an alarming disconnect: Serological protection exists alongside uncontrolled percutaneous exposure and underreporting of approximately one-third of needlestick injuries, a finding that may indicate barriers to reporting within the training environment. Although these findings are clinically relevant, they should be interpreted considering the original study's limitations, including its single-center design, the relatively small serologically evaluated sample ($n = 134$), and the self-reported assessment of needlestick injuries, which may introduce recall bias and limit the applicability of the findings to other settings. Taking care of students therefore requires an inseparable trio: Vaccination policy, occupational safety, and continuous education (Figure 1).

VACCINATION POLICY

Regarding the vaccination policy, the hepatitis B vaccine has been used worldwide for more than 40 years, leading to an approximately 10-fold reduction in hepatitis B infections among vaccinated individuals[3]. The vaccine has demonstrated excellent efficacy, inducing protective levels of antibody to HBsAg (anti-HBs ≥ 10 IU/L) in more than 95% of infants, children, and young adults[4-6]. Non-responders (individuals who do not develop anti-HBs ≥ 10 IU/L after the primary vaccination series) remain at risk of acquiring hepatitis B and should be revaccinated, with seroconversion rates of 50%-99% depending on the regimen used[7,8]. The rate of anti-HBs positivity in healthcare workers vaccinated in infancy is significantly lower than the initial vaccine response rate, ranging from 16%-60% depending on the time since vaccination, with most studies showing that only 28%-51% maintain detectable protective levels when tested in adulthood[9,10]. This long-term decline in detectable anti-HBs levels is well documented in healthcare populations. A similar pattern has also been observed in a large United States cohort of nearly 1000 healthcare workers entering the workforce. Among 986 individuals born in 1991 or later with documented completion of the hepatitis B vaccine series, 51% had anti-HBs levels < 10 IU/L at the time of evaluation. Following administration of a fourth vaccine dose, 88% mounted an anamnestic response, whereas only a small proportion (approximately 5%) failed to respond[11]. In addition, despite progressive decline in detectable anti-HBs levels, immune memory is preserved, and protection against symptomatic acute and chronic hepatitis B infection persists for 30 years or more in most immunocompetent individuals who originally responded to the vaccine[7,10]. However, these exceptions are exactly the reason why vaccination assessment should be performed before healthcare workers are exposed. A randomized study showed that, in the few individuals who remain susceptible after revaccination, alternative vaccination strategies can still lead to protective immunity[12]. This finding is crucial, as it invalidates the passive acceptance of biological susceptibility in high-risk environments.

OCCUPATIONAL SAFETY

The main message of this study is that hepatitis B virus (HBV) prevention in healthcare trainees cannot be reduced to one question of whether vaccination was once received. In blood-exposed settings, protection is both biological and procedural. It depends on immune status, as well as on whether that status is known before exposure, whether non-responders are identified, and whether post-exposure protocols are simple enough for injured trainees to use. Unlike the general population, healthcare trainees at risk of occupational blood exposure should have their hepatitis B immune status documented before clinical placement, in accordance with current national and international recommendations. Individuals without documented post-vaccination serological testing may benefit from anti-HBs assessment to identify those who require revaccination or specific post-exposure management, whereas documented vaccine responders generally remain protected through immune memory despite declining antibody levels[9]. The World Health Orga-




Figure 1 Three pillars for preventing hepatitis B virus occupational risk among healthcare trainees. Anti-HBs: Antibody to hepatitis B surface antigen.

nization position paper also supports targeted serologic testing in groups in whom knowledge of immune status has practical value, including individuals with occupational exposure to blood[13].

The 86.6% seroprotection rate reported by Jaho *et al*[1] can give a wrong impression if seen as a complete success. In a clinical training environment, it leaves a meaningful minority without documented protective titers at the very moment when exposure risk is becoming real. The study becomes even more compelling when this finding is read together with the high burden of needlestick injury and underreporting. A trainee with uncertain immune status and a delayed or unreported exposure cannot be protected by reassuring population averages. Among the major blood-borne viruses, HBV deserves particular attention because it is the most infectious following occupational percutaneous exposure[14]. The estimated risk of transmission after a needlestick injury ranges from 19%-37% when the source patient is HBeAg-positive, compared with approximately 0.2%-1.8% for hepatitis C virus (HCV) and 0.3% for human immunodeficiency virus (HIV). Importantly, these estimates apply primarily to susceptible individuals exposed to highly infectious HBeAg-positive source patients. In healthcare trainees with documented protective immunity after hepatitis B vaccination, the absolute risk is substantially reduced, and post-exposure management is guided by the individual's immune status, source-patient testing, the characteristics of the exposure, and timely post-exposure prophylaxis when indicated[8,9]. The high transmission potential in susceptible individuals reflects the combination of high circulating viral loads, remarkable environmental stability, and efficient transmission through percutaneous and mucosal exposure. HBV is also unique because highly effective preventive strategies are available, including pre-exposure vaccination, post-vaccination serological testing, and post-exposure prophylaxis with hepatitis B immune globulin and vaccination for susceptible individuals. Moreover, it is essential to emphasize that HCV and HIV are also transmitted by needlestick injuries, reinforcing the importance of prompt reporting and appropriate post-exposure evaluation[15,16].

CONTINUOUS EDUCATION

Continuous education is equally essential. Another important lesson from Jaho *et al*[1] is behavioral rather than immunological. Even when vaccination programs perform well, unsafe exposures still occur, and trainees often do not report them. This pattern is not exclusive to Albania. Studies in healthcare students and residents in different settings have documented the same combination of sharps injuries and underreporting, often related to fear of blame, concern about academic consequences, uncertainty about reporting routes, or the idea that minor injuries are not important[17,18]. These observations are consistent with a culture of underreporting that has been documented in healthcare students and residents in different settings. The Albanian findings are also consistent with global epidemiological evidence showing that percutaneous injuries remain a common occupational hazard among healthcare workers worldwide. A large meta-analysis estimated a global one-year prevalence of 36.4%, with substantial regional variation, including 43.2% in Asia, 34.5% in Africa, and 31.8% in Europe[19]. Underreporting has likewise been recognized as a pervasive international problem, suggesting that the true burden of occupational exposures is likely underestimated across many healthcare systems[20]. Thus, the findings from Albania are consistent with a broader global pattern rather than representing an isolated local phenomenon. Underreporting is not only an individual problem. It usually reflects a process that is too slow, unclear, or difficult to use. In such conditions, students may feel that reporting is not worth the effort. As a result, important steps are missed, including source evaluation, timely prophylaxis, and understanding where injuries are actually happening[21]. In addition, the study suggests that occupational risk may not be equally distributed across training areas[1]. Some groups may be exposed earlier or more frequently to sharps injuries, while having less confirmed protection. Surveillance may be particularly valuable in settings where exposure risk and gaps in immunity are more likely to occur[22].

The main message of this study is practical. Training institutions need a clear pathway for serological verification and management of blood exposure. Evidence from current guidelines supports a structured approach: Confirm vaccination history when records are available; distinguish individuals with documented vaccine response from those without previous post-vaccination serological testing; assess anti-HBs when immune status is uncertain or documentation is unavailable; identify true non-responders who require specific post-exposure management; and provide individualized management for immunocompromised trainees according to current recommendations[9,21]. Individuals who remain non-responsive after the recommended revaccination strategy should undergo evaluation for HBV infection, including testing for HBsAg. If current HBV infection is excluded, they should be considered susceptible to HBV, counselled regarding their ongoing occupational risk, and instructed to seek immediate occupational health evaluation after any blood or body fluid exposure for timely post-exposure prophylaxis[23]. Documenting persistent non-response may facilitate timely and appropriate post-exposure management by reducing uncertainty about immune status at the time of occupational exposure and enabling prompt implementation of appropriate post-exposure prophylaxis[23,24]. Simultaneously, verification of immune status alone is insufficient. Reporting systems must be simple, fast, and clearly non-punitive. Trainees should know before clinical placement whom to contact and what to do after an exposure[21,23]. A practical framework summarizing these recommendations is presented in Figure 2. Electronic reporting systems and digital surveillance tools may facilitate timely reporting of occupational exposures, improve adherence to post-exposure protocols, and strengthen institutional surveillance of exposure events[25].




Figure 2 Proposed approach to hepatitis B virus prevention and post-exposure management among healthcare trainees. Anti-HBs: Antibody to hepatitis B surface antigen; HBV: Hepatitis B virus; HCV: Hepatitis C virus; HIV: Human immunodeficiency virus; HBIG: Hepatitis B immune globulin.

CONCLUSION

Ultimately, although the success of hepatitis B immunization is undeniable, high seroprotection rates should not be used to justify gaps in post-exposure safety. Even trainees who are immune to HBV remain vulnerable to unsafe sharps practices, delayed reporting, and exposure to other blood-borne pathogens. Vaccination prevents disease, but effective protection also depends on standard precautions, verified immunity when appropriate, and clear, timely post-exposure pathways.

REFERENCES

- 1 **Jaho J**, Balilaj L, Bulla A, Capraro Y, Lamfu L, Baiocchi L. Serological evidence of hepatitis B virus exposure and vaccination coverage trends in health sciences students in Albania. *World J Virol* 2026; **15**: 118992 [RCA] [PMID: 42395040 DOI: 10.5501/wjv.v15.i2.118992] [Full Text] [Full Text(PDF)]
- 2 **Van Damme P**. Hepatitis B: vaccination programmes in Europe--an update. *Vaccine* 2001; **19**: 2375-2379 [RCA] [PMID: 11257363 DOI: 10.1016/s0264-410x(00)00457-6] [Full Text]
- 3 **Gerlich WH**. [Hepatitis B vaccines-history, achievements, challenges, and perspectives]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2022; **65**: 170-182 [RCA] [PMID: 35015108 DOI: 10.1007/s00103-021-03484-w] [Full Text] [Full Text(PDF)]
- 4 **Seto WK**, Lo YR, Pawlotsky JM, Yuen MF. Chronic hepatitis B virus infection. *Lancet* 2018; **392**: 2313-2324 [RCA] [PMID: 30496122 DOI: 10.1016/S0140-6736(18)31865-8] [Full Text]
- 5 **Trépo C**, Chan HL, Lok A. Hepatitis B virus infection. *Lancet* 2014; **384**: 2053-2063 [RCA] [PMID: 24954675 DOI: 10.1016/S0140-6736(14)60220-8] [Full Text]
- 6 **Jeng WJ**, Papatheodoridis GV, Lok ASF. Hepatitis B. *Lancet* 2023; **401**: 1039-1052 [RCA] [PMID: 36774930 DOI: 10.1016/S0140-6736(22)01468-4] [Full Text]
- 7 **Raven SFH**, Hoebe CJP, Vossen ACTM, Visser LG, Hautvast JLA, Roukens AHE, van Steenberghe JE. Serological response to three alternative series of hepatitis B revaccination (Fendrix, Twinrix, and HBVaxPro-40) in healthy non-responders: a multicentre, open-label, randomised, controlled, superiority trial. *Lancet Infect Dis* 2020; **20**: 92-101 [RCA] [PMID: 31629649 DOI: 10.1016/S1473-3099(19)30417-7] [Full Text]
- 8 **Terrault NA**, Lok ASF, McMahon BJ, Chang KM, Hwang JP, Jonas MM, Brown RS Jr, Bzowej NH, Wong JB. Update on prevention, diagnosis, and treatment of chronic hepatitis B: AASLD 2018 hepatitis B guidance. *Hepatology* 2018; **67**: 1560-1599 [RCA] [PMID: 29405329 DOI: 10.1002/hep.29800] [Full Text] [Full Text(PDF)]
- 9 **Schillie S**, Vellozzi C, Reingold A, Harris A, Haber P, Ward JW, Nelson NP. Prevention of Hepatitis B Virus Infection in the United States: Recommendations of the Advisory Committee on Immunization Practices. *MMWR Recomm Rep* 2018; **67**: 1-31 [RCA] [PMID: 29939980 DOI: 10.15585/mmwr.rr6701a1] [Full Text] [Full Text(PDF)]
- 10 **Phattraprayoon N**, Kakheaw J, Soonklang K, Cheirsilpa K, Ungtrakul T, Auewarakul C, Mahanonda N. Duration of Hepatitis B Vaccine-Induced Protection among Medical Students and Healthcare Workers following Primary Vaccination in Infancy and Rate of Immunity Decline. *Vaccines (Basel)* 2022; **10**: 267 [RCA] [PMID: 35214725 DOI: 10.3390/vaccines10020267] [Full Text] [Full Text(PDF)]
- 11 **Ritscher AM**, LeClair-Netzel M, Friedlander NJ, Howard Stewart DN, Wagner M, Kalscheur N, Caldera F, Hayney MS. Cross-sectional study of hepatitis B antibody status in health care workers immunized as children at an academic medical center in Wisconsin. *Vaccine* 2020; **38**: 1597-1600 [RCA] [PMID: 31955961 DOI: 10.1016/j.vaccine.2019.12.036] [Full Text]
- 12 **Marks KM**, Kang M, Umbleja T, Cox A, Vigil KJ, Ta NT, Omoz-Oarhe A, Perazzo H, Kosgei J, Hatlen T, Price J, Katsidzira L, Supparatpinoy K, Knowles K, Alston-Smith BL, Rathod P, Sherman KE; ACTG 5379 (BEE-HIVE) Study Team. HepB-CpG vs HepB-Alum Vaccine in People With HIV and Prior Vaccine Nonresponse: The BEE-HIVE Randomized Clinical Trial. *JAMA* 2025; **333**: 295-306 [RCA] [PMID: 39616603 DOI: 10.1001/jama.2024.24490] [Full Text] [Full Text(PDF)]
- 13 **World Health Organization**. Hepatitis B vaccines: WHO position paper, July 2017 - Recommendations. *Vaccine* 2019; **37**: 223-225 [RCA] [PMID: 28743487 DOI: 10.1016/j.vaccine.2017.07.046] [Full Text]
- 14 **Wu HM**, Nilles E. International Travel to Deliver Healthcare in Resource-Limited Settings. 2025 Apr 23. In: CDC Yellow Book, 2026 edition: Health Information for International Travel [Internet]. Atlanta (GA): Centers for Disease Control and Prevention (US); 2025 [PMID: 41818539]
- 15 **Dolman GE**, Koffas A, Phipps E, Kennedy PTF. Clinical and occupational health management of healthcare workers living with chronic hepatitis B: UK policy and international comparisons. *J Viral Hepat* 2021; **28**: 976-981 [RCA] [PMID: 33665937 DOI: 10.1111/jvh.13494] [Full Text]
- 16 **Mengistu DA**, Tolera ST, Demmu YM. Worldwide Prevalence of Occupational Exposure to Needle Stick Injury among Healthcare Workers: A Systematic Review and Meta-Analysis. *Can J Infect Dis Med Microbiol* 2021; **2021**: 9019534 [RCA] [PMID: 33564345 DOI: 10.1155/2021/9019534] [Full Text] [Full Text(PDF)]
- 17 **Papadopoli R**, Bianco A, Pepe D, Pileggi C, Pavia M. Sharps and needle-stick injuries among medical residents and healthcare professional students: pattern and reporting in Italy-a cross-sectional analytical study. *Occup Environ Med* 2019; **76**: 739-745 [RCA] [PMID: 31439689 DOI: 10.1136/oemed-2019-105873] [Full Text]
- 18 **Black Thomas LM**. Underreporting of Bloodborne Pathogen Exposures in Nursing Students. *Nurse Educ* 2020; **45**: 78-82 [RCA] [PMID: 31145179 DOI: 10.1097/NNE.0000000000000696] [Full Text]
- 19 **Auta A**, Adewuyi EO, Tor-Anyiin A, Edor JP, Kureh GT, Khanal V, Oga E, Adeyoye D. Global prevalence of percutaneous injuries among healthcare workers: a systematic review and meta-analysis. *Int J Epidemiol* 2018; **47**: 1972-1980 [RCA] [PMID: 30272173 DOI: 10.1093/ije/dyy208] [Full Text]
- 20 **Abdelmalik MA**, Alhowaymel FM, Fadlalmola H, Mohammed MO, Abbakr I, Alenezi A, Mohammed AM, Abaoud AF. Global prevalence of needle stick injuries among nurses: A comprehensive systematic review and meta-analysis. *J Clin Nurs* 2023; **32**: 5619-5631 [RCA] [PMID: 36841963 DOI: 10.1111/jocn.16661] [Full Text]
- 21 **Puro V**, De Carli G, Cicalini S, Soldani F, Balslev U, Begovac J, Boaventura L, Campins Marti M, Hernández Navarrete MJ, Kammerlander

- R, Larsen C, Lot F, Lunding S, Marcus U, Payne L, Pereira AA, Thomas T, Ippolito G. European recommendations for the management of healthcare workers occupationally exposed to hepatitis B virus and hepatitis C virus. *Euro Surveill* 2005; **10**: 11-12 [RCA] [PMID: 29208102 DOI: 10.2807/esm.10.10.00573-en] [FullText]
- 22 **Henderson DK**, Dembry LM, Sifri CD, Palmore TN, Dellinger EP, Yokoe DS, Grady C, Heller T, Weber D, Del Rio C, Fishman NO, Deloney VM, Lundstrom T, Babcock HM. Management of healthcare personnel living with hepatitis B, hepatitis C, or human immunodeficiency virus in US healthcare institutions. *Infect Control Hosp Epidemiol* 2022; **43**: 147-155 [RCA] [PMID: 33050959 DOI: 10.1017/ice.2020.458] [FullText]
- 23 **Jeng WJ**, Yip TC, Lok AS. Hepatitis B: A Review. *JAMA* 2026; **335**: 1879-1892 [RCA] [PMID: 42081318 DOI: 10.1001/jama.2026.6070] [FullText]
- 24 **Moore Ii R**, Porter CL, Darling JM. Hepatitis B: Part I. Updates on Screening and Prevention. *Am Fam Physician* 2026; **113**: 229-234 [RCA] [PMID: 41839073]
- 25 **Bammann E**, Nikrad H, Aragon D, Young H. Implementation of a secure electronic form for bloodborne pathogen exposure reporting associated with increased reports among healthcare workers: A quasi-experimental study. *Am J Infect Control* 2025; **53**: 869-874 [RCA] [PMID: 40081496 DOI: 10.1016/j.ajic.2025.03.008] [FullText]

FOOTNOTES

Specialty type: Virology

Country of origin: Brazil

Author contributions: Narciso-Schiavon JL solely contributed to the conception, literature review, manuscript writing, revision of the manuscript, and approved the final version for publication.

AI contribution statement: Artificial intelligence tools (ChatGPT, OpenAI) were used exclusively for language editing and clarity improvement. The author critically reviewed and verified all AI-assisted suggestions, approved the final content, and takes full responsibility for the integrity, accuracy, and scientific validity of the manuscript.

Conflict-of-interest statement: The author declares no conflict of interest.

S-Editor: Liu H

L-Editor: A

P-Editor: Zhao YQ



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

