

World Journal of *Gastroenterology*

World J Gastroenterol 2024 November 14; 30(42): 4518-4596



EDITORIAL

- 4518 Sepsis-associated liver injury: Mechanisms and potential therapeutic targets
Chen JW, Liu CY, Li S, Wu SW, Cai C, Lu MQ

ORIGINAL ARTICLE**Retrospective Study**

- 4523 Comparing gastrointestinal dysfunction score and acute gastrointestinal injury grade for predicting short-term mortality in critically ill patients
Shen C, Wang X, Xiao YY, Zhang JY, Xia GL, Jiang RL

Prospective Study

- 4532 Strategic insights into the cultivation of pancreatic cancer organoids from endoscopic ultrasonography-guided biopsy tissue
Yang JL, Zhang JF, Gu JY, Gao M, Zheng MY, Guo SX, Zhang T

Basic Study

- 4544 Transfection of 12/15-lipoxygenase effectively alleviates inflammatory responses during experimental acute pancreatitis
Sun HW, Bai YY, Qin ZL, Li RZ, Madzikatire TB, Akuetteh PDP, Li Q, Kong HR, Jin YP

CASE REPORT

- 4557 Esophageal melanosis: Two case reports and review of literature
Kazacheuskaya L, Arora K

LETTER TO THE EDITOR

- 4566 Colon cancer screening programs prevent cancer
Flynn DJ, Feuerstein JD
- 4569 Does hemorrhoidal artery embolization really benefit patients with hemorrhoids?
Lei ML, Dong LL, Zhang HP, Yu YB
- 4576 Metabolic dysfunction-associated steatotic liver disease: The question of long-term high-normal alanine aminotransferase as a screening test
Moyana TN
- 4583 Endoscopic polidocanol foam sclerobanding for treatment of internal hemorrhoids: A novel outpatient procedure
Mou AN, Wang YT

- 4587** Evaluation of the GATIS score for predicting prognosis in rectal neuroendocrine neoplasms
Feng YN, Liu LH, Zhang HW
- 4591** *Helicobacter pylori*, esophageal precancerous lesions, and proton pump inhibitor overuse
Zhang F, Zhang H, Liu YM, Tang FS

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Ce-Fan Zhou, PhD, Professor, School of Life and Health Sciences, Vice Dean of Biomedical Research Institute, National "111" Center for Cellular Regulation and Molecular Pharmaceutics, Hubei University of Technology, Wuhan 430068, Hubei Province, China.
cefan@hbut.edu.cn

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (*WJG, World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. *WJG* mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The *WJG* is now abstracted and indexed in Science Citation Index Expanded (SCIE), MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for *WJG* as 4.3; Quartile: Q1. The *WJG*'s CiteScore for 2023 is 7.8.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Cover Editor: *Jia-Ru Fan*.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski

EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF

Jian-Gao Fan (Chronic Liver Disease)

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

November 14, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

PUBLISHING PARTNER

Shanghai Pancreatic Cancer Institute and Pancreatic Cancer Institute, Fudan University
Biliary Tract Disease Institute, Fudan University

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/bpg/GerInfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

PUBLISHING PARTNER'S OFFICIAL WEBSITE

<https://www.shca.org.cn>
<https://www.zs-hospital.sh.cn>



Metabolic dysfunction-associated steatotic liver disease: The question of long-term high-normal alanine aminotransferase as a screening test

Terence N Moyana

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade C

Novelty: Grade C

Creativity or Innovation: Grade C

Scientific Significance: Grade C

P-Reviewer: Zhao QW

Received: July 17, 2024

Revised: September 26, 2024

Accepted: October 9, 2024

Published online: November 14, 2024

Processing time: 105 Days and 10.8 Hours



Terence N Moyana, Department of Pathology and Laboratory Medicine, University of Ottawa and The Ottawa Hospital, Ottawa K1H 8L6, Ontario, Canada

Corresponding author: Terence N Moyana, FRCPC, MD, Full Professor, Department of Pathology and Laboratory Medicine, University of Ottawa and The Ottawa Hospital, 501 Smyth Road, General Campus, Ottawa K1H 8L6, Ontario, Canada. tmoyana@corla.ca

Abstract

The growing prevalence of metabolic dysfunction-associated steatotic liver disease (MASLD) is being driven by the obesity epidemic. The quest for solutions continues particularly with regard to early detection. This editorial comments on the utility of long-term high-normal alanine aminotransferase (ALT) in screening for MASLD. Chen *et al* found that new onset MASLD can be detected by repetitively high normal ALT. Implicit in this concept is the question of what should be the accepted upper limit of normal (ULN) for ALT. It was previously set at 40 IU/L based on studies that included people with subclinical liver disease but the new consensus is 30/19 U/L in healthy males/females. Thus, when Chen *et al* defines the ULN as 40 U/L, others may view it as excessively high. It is important to recognize the variables affecting ULN *e.g.* instrumentation, diurnal variations, exercise and ageing. These variables matter when the distinctions are subtle *e.g.* normal *vs* high-normal. In this regard, the utility of long-term high normal ALT as a disease marker could be enhanced by combining it with other biomarkers, imaging and MASLD genetics to create machine learning classifiers. All in all, Chen *et al*'s work on long-term high normal ALT as a marker of new-onset MASLD deserves merit.

Key Words: Metabolic dysfunction-associated steatotic liver disease; Alanine aminotransferase; Pathophysiology; Screening; Upper limit of normal; Biomarkers; Predictive models

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: High normal alanine aminotransferase (ALT) has potential utility in screening for new-onset metabolic dysfunction-associated steatotic liver disease (MASLD). In view of its biochemistry, the upper limit of normal for ALT has to be interpreted with caution since it can be influenced by various methodologic and physiologic factors. Its utility as a screening tool could be enhanced by combining it with other serum biomarkers, cardiometabolic risk factors, imaging techniques and MASLD genetics to create predictive nomograms or machine learning classifiers.

Citation: Moyana TN. Metabolic dysfunction-associated steatotic liver disease: The question of long-term high-normal alanine aminotransferase as a screening test. *World J Gastroenterol* 2024; 30(42): 4576-4582

URL: <https://www.wjgnet.com/1007-9327/full/v30/i42/4576.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v30.i42.4576>

TO THE EDITOR

Metabolic dysfunction-associated steatotic liver disease (MASLD) is one of the most common liver diseases world-wide and its prevalence is steadily rising. This is largely being driven by lifestyle changes (overnutrition, food composition and physical underactivity) leading to the obesity epidemic[1,2]. Consequently, it is a growing cause of morbidity and mortality. The quest for solutions continues. One approach is early detection and timely intervention and in this regard Chen *et al*[3] investigated the utility of long-term high-normal alanine aminotransferase (ALT) as a screening tool for this condition[3]. There is a fairly long history of ALT as a surrogate marker for MASLD[4-7]. This editorial comments on the role of this enzyme in screening for MASLD.

Nomenclature

The nomenclature for nonalcoholic fatty liver disease (NAFLD) was recently changed to MASLD[8]. The main reasons driving the change were: (1) The terms “nonalcoholic” and “fatty” were considered to be stigmatizing; (2) NAFLD was quite limiting as a diagnosis because some such patients consume alcohol to a greater or lesser extent *i.e.* an overlap of MASLD with alcoholic liver disease (ALD) resulting in MetALD; and (3) The term failed to recognize the root cause of the disease, namely insulin resistance (IR)[1,2,8]. In fact, one of the major changes in moving from the old to the new terminology is that the definition of the disease now includes at least one of 5 cardiometabolic risk factors (CMRFs) pertaining to body mass index (BMI), type 2 diabetes mellitus (T2DM), elevated blood pressure, high plasma triglycerides and low high-density lipoprotein-cholesterol[8]. In light of the epidemiology of the disease as highlighted by Chen *et al* [3], it is also important that these CMRFs be adjusted according to ethnicity *e.g.* Hispanics, Europeans, Africans and Asians[3,9,10]. At the same time, it should be noted that the nomenclature used by Chen *et al*[3], even in the title, does not quite reflect the new terminology.

The entities comprising MASLD

MASLD represents a dynamic spectrum of liver disease[1,2,8]. At one end is metabolic dysfunction-associated steatotic liver (MASL) or simple steatosis, which is defined as the presence $\geq 5\%$ steatosis with no evidence of hepatocellular injury. Further along the spectrum is metabolic dysfunction-associated steatohepatitis (MASH) which is indicative of the onset of hepatocyte membrane damage and the inflammatory cascade with or without some degree of fibrosis. At the other end is cirrhosis characterized by advanced fibrosis and its attendant complications. Along the spectrum, fibrosis is the dominant morphologic feature that predicts morbidity and mortality[1,2,8]. Screening protocols such as those advocated by Chen *et al*[3] and McGinty and Przemioslo[11] aim to detect the disease before it gets to cirrhosis.

The rationale for ALT in screening for MASLD

ALT occurs abundantly in the cytosol of hepatocytes where it catalyzes the transfer of amino groups to form oxaloacetate [7,12]. Its activity in the liver is 3000 times greater than in the serum and as such, it is a sensitive marker of hepatocyte injury/loss of membrane integrity. Its plasma half-life is 47 hours which is significantly longer than that of serum aspartate aminotransferase (AST) (17 hours). In acute liver cell injury, AST usually rises immediately but within a day or 2, ALT will become higher due to its longer half-life[7,12]. Thus as a general rule, with ongoing liver damage, ALT is more commonly elevated than AST though exceptions occur (*e.g.* ALD).

Pathogenesis of liver damage in MASLD leading to changes in ALT

A key feature of MASLD is the impaired response of tissues to insulin *i.e.* IR[1,2,8]. The metabolic consequences of IR are hyperglycemia, dyslipidemia and visceral adiposopathy with the latter typically manifesting as abdominal fat. A vicious cycle occurs resulting in decreased uptake of glucose and release of free fatty acids which further aggravates hepatic fatty infiltration. Accumulation of fat in the liver can cause oxidative stress, release of cytokines and lipid peroxidation culminating in hepatocyte injury and/or death *i.e.* lipotoxicity[1,2,13,14]. This triggers the inflammatory process that can lead to MASH and, if left unchecked to fibrosis. Traditionally, MASL has been associated with normal ALT levels and MASH abnormal levels. The postulate by Chen *et al*[3] is that long-term high normal ALT is associated with new onset MASLD. It does not specifically distinguish between MASL and MASH. However, it could be argued that their cohort with abnormally high ALT (as opposed to high-normal) most likely represented MASH.

Fibrosis remodeling and regression

Fibrosis starts appearing at the MASH stage of the disease due in part to the activation of stellate cells and microvascular injury[1,15]. It begins as fine filmy wisps of fibrosis, so-called chicken-wire fibrosis that can be pericellular, perisinusoidal or perivenular. This is a very important stage in the disease process because the fibrosis is still quite delicate and amenable to regression[1,15-17]. Due to the concurrent activation of the inflammatory process during MASH, this is the period when ALT is most likely to be elevated thus offering the prospect of timely intervention. With cirrhosis, ALT can be quite variable depending on the balance between inflammatory activity and fibrosis *e.g.* burnt out disease with shrunken liver due to hepatocyte loss. Notwithstanding the timely concepts of remodeling and regression, it is important to note that regression as currently envisaged is a fairly modest process and does not imply resolution of cirrhosis since large areas of parenchymal extinction cannot be repopulated by hepatocytes[15,17,18].

DISCUSSION

Chen *et al*[3] defined MASLD based on abdominal ultrasound (US) findings and at least one of several CMRFs which is similar to what is envisaged in the new terminology[8]. Their study found that long-term high normal ALT could identify cases of new-onset MASLD. This is very helpful because such individuals can then be targeted for lifestyle changes/intervention. However, based on pathophysiology and natural history of MASLD, a number of discussion points come up with respect to the validity of long-term high normal ALT as a screening tool for this condition.

ALT as a screening test for MASLD

Screening is defined as the presumptive identification of unrecognized disease by tests which can be carried out easily and conveniently[12,19], and as emphasized by McGinty and Przemioslo[11], should be cost-effective. A screening test is not intended to be diagnostic but to classify individuals with a high probability of disease from those with a low probability. Based on the pathophysiology of MASLD and the biochemistry of ALT, changes in ALT would not be expected to occur until there is loss of hepatocyte membrane integrity[1,2,13,14,20]. Most studies do indeed reaffirm this but others show that the levels can be normal in subjects across the entire spectrum of MASLD[21,22]. Since physicians tend to pay more attention to those patients with elevated ALTs, it may be useful to devise a way for them to look at ALT levels in a different light. Essentially, this is part of the message being conveyed by Chen *et al*[3]. They are postulating that new onset MASLD can be detected by repetitively high normal ALT, and that this should be the trigger point for intervention/action rather than waiting for a truly abnormal level. Implicit in this concept is the question of what should be the accepted upper limit of normal (ULN) for ALT.

ULN for ALT

Ideally, the ULN should be determined in individuals without definite liver disease but such subjects are difficult to identify in population-based studies. It was previously set at 40 IU/L regardless of sex or BMI based on studies that were carried out before the introduction of viral hepatitis C testing and prior to the concept of MASLD[23-26]. Inevitably, this included a proportion of people with subclinical liver disease (*e.g.* viral hepatitis carriers and MASLD) thereby limiting the sensitivity of ALT as a tool for assessing liver damage[6,7,12]. In light of the changing epidemiology of liver disease, this called for a re-evaluation of the ULN after excluding individuals with subclinical liver disease. There appears to be convergence towards a new ULN of 30/19 U/L in healthy males and females (M/F) though there is no unanimity[6,25-28]. Thus, when Chen *et al*[3] define their ULN as 40 U/L, others may view it as excessively high.

Variables affecting ALT values

Methodology: Chen *et al*[3] rightly considered some of the factors affecting the ULN for ALT *e.g.* variability in methods and instrumentation depending on geographic location[28,29]. For example: (1) There is progressive ALT activity loss the longer the time interval from collection of the blood sample to centrifugation[27,28]; (2) Freezing the serum specimens after collection followed by thawing prior to assay can cause a 46% loss of ALT activity compared to only a 6% loss with refrigeration at 4 °C[27]; (3) Conducting the test at 37 °C instead of 30 °C raises enzyme activity and results in higher ALT values[28] and (4) Laboratories using different instruments and reagents may report different ALT levels[29]. These are well-known issues in clinical chemistry and efforts have been made to standardize analyzer and assay measurements but with limited success[28,30-32]. In this regard, recommendations have been made to use a reference interval rather than a fixed cutoff until such a time as there are internationally harmonized ALT methods[32,33].

Physiology of ALT: (1) It is also found in the kidney and to a lesser extent in cardiac and skeletal muscle. This may not matter as much when ALT is significantly elevated *e.g.* $\times 5$, $\times 10$ or $\times 15$ normal as can occur in certain conditions[12,31]. However, in high-normal or borderline cases, it may matter; (2) ALT activity varies day to day, by as much as 10% to 30%; (3) Within a given day, there is significant diurnal variation, with ALT activities being up to 45% higher in the afternoon than early morning[12,34]; (4) Seasonal variations also occur[35]; (5) ALT activity can vary quite substantially depending on exercise: No exercise, customary exercise levels and strenuous activity[12,36]; (6) Fasting has a bearing on ALT levels; hence the recommendation that any mildly increased ALT should be re-evaluated in the fasting state[37]; and (7) ALT activity is substantially higher in adult males than females, and also increases with age up to a certain point[38]. Chen *et al*'s study rightly recognizes the M/F differences and establishes a lower end cut-off of 18 years[3]. However, it is not clear to what extent Chen *et al*[3] took into account all of the above physiologic variables.

Ageing and osteoporosis can alter ALT activity in the serum: In fact, after age 60, the reference limits change in women [39]. Furthermore, other investigators have suggested that elevations in ALT in middle-aged and elderly men and women with MASLD could be related to reduced bone mineral density as a result of curtailed osteoblast activity[40-42] rather than being an indication of MASLD.

Non-invasive tests for MASLD

While liver biopsy is the gold standard for diagnosing MASLD, it has limitations due to its invasiveness, inherent complications (e.g. pain, bleeding, perforation), spatial variability in sampling and impracticality for serial monitoring[43-45]. Recent approaches using endoscopic US-guided biopsies have some advantages over the traditional percutaneous samples but do not completely mitigate these drawbacks[46,47], hence the need for non-invasive tests. They can be classified as: (1) Serum biomarkers; and (2) Imaging techniques.

Serum biomarkers: Although ALT is the enzyme most commonly used for screening of MASLD, it's utility can be augmented by other biomarkers potentially analyzable from the same routine blood samples. They include but are not limited to AST, gamma-glutamyl transferase, fasting blood glucose, hemoglobin A1c, triglycerides and cholesterol. These biomarkers are increasing being used in conjunction with other parameters in various algorithms for assessing MASLD e.g. the fatty liver index[48-50], liver fat score[49], and hepatic steatosis index[49,51].

Imaging techniques: Various imaging techniques such as abdominal US, computed tomography (CT) and magnetic resonance imaging (MRI) can be used in the assessment of MASLD. Of these: (1) US is the fastest and least expensive and can detect hepatic fat by comparing liver hyper-echogenicity in relation to the kidney. However, it's utility is limited by low sensitivity for mild disease. Furthermore, accuracy decreases as BMI increases[52,53]. Vibration-controlled transient elastography (VCTE) (Fibroscan) is an US-based method that utilizes shear wave velocity to perform liver stiffness measurement. When equipped with a module called controlled attenuation parameter, it can be very useful for quantifying the levels of liver fat and fibrosis[44,54,55]. The test is instantaneous and reliable which makes it suitable for population-wide screening as well as point-of-care testing[44,55]; (2) On CT, fat reduces liver attenuation causing it to be hypodense and can provide detailed information, potentially averting needle biopsies[43]. However, it is more expensive than US especially when applied to the general population; and (3) Unlike US and CT which use proxies to assess liver fat, magnetic resonance can measure the proton density fat fraction which is highly accurate in quantifying liver fat and may be even more precise than liver biopsy[43,45]. A significant drawback is that MRI-based techniques are costly in terms of equipment/scanners and require technical and professional expertise that is usually only available in tertiary care centres. As such, they may not be suitable for large scale population screening.

Prediction models and machine learning classifiers in MASLD

Given all the variables that affect ALT measurements, it is prudent to consider other approaches for assessing risk in MASLD. This concept has been captured by using combined biomarker parameters. Scores such as Agile 3 + and Agile 4 combine laboratory data (e.g. ALT, AST, fasting glucose, triglycerides, high- and low density lipoproteins), demographic features (e.g. age, sex, BMI, waist-to-hip ratio, T2DM, hypertension) and liver stiffness measurements as determined by VCTE[43,56]. The data can also be used to create multiparameter predictive nomograms or machine learning classifiers. Such predictive models are promising to be effective not only as screening tools for assessing risk in MASLD but also for identifying patients for referral to specialists/hepatologists[1,57-59].

Future directions

While liver biopsy is the gold standard for diagnosing MASLD, it is not practical for population studies. A more useful and practical alternative would be serum biomarkers which in addition to ALT, could include AST/ALT ratio, Fibrosis-4, Hepascore, C-reactive protein, ferritin, serum hyaluronic acid and alpha-2-macroglobulin[1,60-62]. From a medical imaging perspective, US is gaining wider acceptance as a biomarker since it can be used in population-based studies. While CT and MRI are still quite expensive, continuing improvements in imaging technologies could improve accessibility, enhance accuracy and lower costs. A potentially exciting field is MASLD genetics: It is known that the risk of developing MASLD is quite variable among ethnic groups and individuals, and is determined by both environmental and genetic factors[10]. Candidate genes include *PNPLA3*, *TM6SF2*, *HSD17B13*, *MBOAT7* and *GCKR* which can be used for creating polygenic risk scores to gain further insights into MASLD prevalence and progression[10,63]. From a prognostic viewpoint, regression of disease from MASH to MASL does occur but whether remodeling of advanced fibrosis/cirrhosis can be achieved is still an open question.

CONCLUSION

The MASLD epidemic appears to be growing. There are ongoing efforts to reverse this trend particularly changes in lifestyle. Screening to identify people at risk can be helpful, but to be cost-effective, it has to be targeted. In this regard, individuals with CMRFs stand to benefit the most from ALT screening and this is part of the message conveyed by Chen *et al*[3] and McGinty and Przemioslo[11]. The question is what is the actionable level for ALT for professional intervention and whether it can be augmented by other approaches such as MASLD genetics, predictive nomograms and/or machine learning classifiers.

FOOTNOTES

Author contributions: As the sole author, Moyana TN is responsible for all aspects of the work, including conception, design, research, writing, and finalization of the manuscript.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: Canada

ORCID number: Terence N Moyana 0009-0006-8426-8638.

S-Editor: Fan M

L-Editor: A

P-Editor: Yuan YY

REFERENCES

- Farrell GC, Larter CZ. Nonalcoholic fatty liver disease: from steatosis to cirrhosis. *Hepatology* 2006; **43**: S99-S112 [PMID: 16447287 DOI: 10.1002/hep.20973]
- Reddy YK, Marella HK, Jiang Y, Ganguli S, Snell P, Podila PSB, Maliakkal B, Satapathy SK. Natural History of Non-Alcoholic Fatty Liver Disease: A Study With Paired Liver Biopsies. *J Clin Exp Hepatol* 2020; **10**: 245-254 [PMID: 32405181 DOI: 10.1016/j.jceh.2019.07.002]
- Chen JF, Wu ZQ, Liu HS, Yan S, Wang YX, Xing M, Song XQ, Ding SY. Cumulative effects of excess high-normal alanine aminotransferase levels in relation to new-onset metabolic dysfunction-associated fatty liver disease in China. *World J Gastroenterol* 2024; **30**: 1346-1357 [PMID: 38596503 DOI: 10.3748/wjg.v30.i10.1346]
- Siest G, Schiele F, Galteau MM, Panek E, Steinmetz J, Fagnani F, Gueguen R. Aspartate aminotransferase and alanine aminotransferase activities in plasma: statistical distributions, individual variations, and reference values. *Clin Chem* 1975; **21**: 1077-1087 [PMID: 1137913 DOI: 10.1093/clinchem/21.8.1077]
- Sherman KE. Alanine aminotransferase in clinical practice. A review. *Arch Intern Med* 1991; **151**: 260-265 [PMID: 1992953 DOI: 10.1001/archinte.151.2.260]
- Prati D, Taioli E, Zanella A, Della Torre E, Butelli S, Del Vecchio E, Vianello L, Zanuso F, Mozzi F, Milani S, Conte D, Colombo M, Sirchia G. Updated definitions of healthy ranges for serum alanine aminotransferase levels. *Ann Intern Med* 2002; **137**: 1-10 [PMID: 12093239 DOI: 10.7326/0003-4819-137-1-200207020-00006]
- Liu Z, Que S, Xu J, Peng T. Alanine aminotransferase-old biomarker and new concept: a review. *Int J Med Sci* 2014; **11**: 925-935 [PMID: 25013373 DOI: 10.7150/ijms.8951]
- Lazarus JV, Newsome PN, Francque SM, Kanwal F, Terrault NA, Rinella ME. Reply: A multi-society Delphi consensus statement on new fatty liver disease nomenclature. *Hepatology* 2024; **79**: E93-E94 [PMID: 37983810 DOI: 10.1097/HEP.0000000000000696]
- Bell CN, Kerr J, Young JL. Associations between Obesity, Obesogenic Environments, and Structural Racism Vary by County-Level Racial Composition. *Int J Environ Res Public Health* 2019; **16** [PMID: 30857286 DOI: 10.3390/ijerph16050861]
- Trépo E, Valenti L. Update on NAFLD genetics: From new variants to the clinic. *J Hepatol* 2020; **72**: 1196-1209 [PMID: 32145256 DOI: 10.1016/j.jhep.2020.02.020]
- McGinty G, Przemioslo R. Effects of excess high-normal alanine aminotransferase levels in relation to new-onset metabolic dysfunction-associated fatty liver disease: Clinical implications. *World J Gastroenterol* 2024; **30**: 3264-3267 [PMID: 39086753 DOI: 10.3748/wjg.v30.i27.3264]
- Kim WR, Flamm SL, Di Bisceglie AM, Bodenheimer HC; Public Policy Committee of the American Association for the Study of Liver Disease. Serum activity of alanine aminotransferase (ALT) as an indicator of health and disease. *Hepatology* 2008; **47**: 1363-1370 [PMID: 18366115 DOI: 10.1002/hep.22109]
- Takahashi Y, Fukusato T. Histopathology of nonalcoholic fatty liver disease/nonalcoholic steatohepatitis. *World J Gastroenterol* 2014; **20**: 15539-15548 [PMID: 25400438 DOI: 10.3748/wjg.v20.i42.15539]
- Parthasarathy G, Revelo X, Malhi H. Pathogenesis of Nonalcoholic Steatohepatitis: An Overview. *Hepatol Commun* 2020; **4**: 478-492 [PMID: 32258944 DOI: 10.1002/hep4.1479]
- Wang S, Friedman SL. Hepatic fibrosis: A convergent response to liver injury that is reversible. *J Hepatol* 2020; **73**: 210-211 [PMID: 32402525 DOI: 10.1016/j.jhep.2020.03.011]
- Wong VW, Wong GL, Choi PC, Chan AW, Li MK, Chan HY, Chim AM, Yu J, Sung JJ, Chan HL. Disease progression of non-alcoholic fatty liver disease: a prospective study with paired liver biopsies at 3 years. *Gut* 2010; **59**: 969-974 [PMID: 20581244 DOI: 10.1136/gut.2009.205088]
- Khan S, Saxena R. Regression of Hepatic Fibrosis and Evolution of Cirrhosis: A Concise Review. *Adv Anat Pathol* 2021; **28**: 408-414 [PMID: 34326286 DOI: 10.1097/PAP.0000000000000312]
- Manne V, Akhtar E, Saab S. Cirrhosis regression in patients with viral hepatitis B and C: a systematic review. *J Clin Gastroenterol* 2014; **48**: e76-e84 [PMID: 24921210 DOI: 10.1097/MCG.0000000000000162]
- Maxim LD, Niebo R, Utell MJ. Screening tests: a review with examples. *Inhal Toxicol* 2014; **26**: 811-828 [PMID: 25264934 DOI: 10.3109/08958378.2014.955932]
- Pratt DS, Kaplan MM. Evaluation of abnormal liver-enzyme results in asymptomatic patients. *N Engl J Med* 2000; **342**: 1266-1271 [PMID: 10811111 DOI: 10.1056/NEJM200005113421001]

- 10781624 DOI: 10.1056/NEJM200004273421707]
- 21 **Ma X**, Liu S, Zhang J, Dong M, Wang Y, Wang M, Xin Y. Proportion of NAFLD patients with normal ALT value in overall NAFLD patients: a systematic review and meta-analysis. *BMC Gastroenterol* 2020; **20**: 10 [PMID: 31937252 DOI: 10.1186/s12876-020-1165-z]
 - 22 **Kunde SS**, Lazenby AJ, Clements RH, Abrams GA. Spectrum of NAFLD and diagnostic implications of the proposed new normal range for serum ALT in obese women. *Hepatology* 2005; **42**: 650-656 [PMID: 16037946 DOI: 10.1002/hep.20818]
 - 23 **Ludwig J**, Viggiano TR, McGill DB, Oh BJ. Nonalcoholic steatohepatitis: Mayo Clinic experiences with a hitherto unnamed disease. *Mayo Clin Proc* 1980; **55**: 434-438 [PMID: 7382552]
 - 24 **Alberti A**, Morsica G, Chemello L, Cavalletto D, Noventa F, Pontisso P, Ruol A. Hepatitis C viraemia and liver disease in symptom-free individuals with anti-HCV. *Lancet* 1992; **340**: 697-698 [PMID: 1355801 DOI: 10.1016/0140-6736(92)92234-7]
 - 25 **Park JH**, Choi J, Jun DW, Han SW, Yeo YH, Nguyen MH. Low Alanine Aminotransferase Cut-Off for Predicting Liver Outcomes; A Nationwide Population-Based Longitudinal Cohort Study. *J Clin Med* 2019; **8** [PMID: 31514449 DOI: 10.3390/jcm8091445]
 - 26 **Valenti L**, Pelusi S, Bianco C, Ceriotti F, Berzuini A, Iogna Prat L, Trotti R, Malvestiti F, D'Ambrosio R, Lampertico P, Colli A, Colombo M, Tsochatzis EA, Fraquelli M, Prati D. Definition of Healthy Ranges for Alanine Aminotransferase Levels: A 2021 Update. *Hepatol Commun* 2021; **5**: 1824-1832 [PMID: 34520121 DOI: 10.1002/hep4.1794]
 - 27 **Williams KM**, Williams AE, Kline LM, Dodd RY. Stability of serum alanine aminotransferase activity. *Transfusion* 1987; **27**: 431-433 [PMID: 3629675 DOI: 10.1046/j.1537-2995.1987.27587320539.x]
 - 28 **Shimizu Y**, Ichihara K. Elucidation of stability profiles of common chemistry analytes in serum stored at six graded temperatures. *Clin Chem Lab Med* 2019; **57**: 1388-1396 [PMID: 30860975 DOI: 10.1515/cclm-2018-1109]
 - 29 **Green JL**, Campagna E, Bogdan GM, Dart RC, Heard K. Effects of clinical laboratory choice on study outcome: an interlaboratory evaluation of aminotransferase levels. *Pharmacotherapy* 2008; **28**: 453-457 [PMID: 18363529 DOI: 10.1592/phco.28.4.453]
 - 30 **Lee JK**, Shim JH, Lee HC, Lee SH, Kim KM, Lim YS, Chung YH, Lee YS, Suh DJ. Estimation of the healthy upper limits for serum alanine aminotransferase in Asian populations with normal liver histology. *Hepatology* 2010; **51**: 1577-1583 [PMID: 20162730 DOI: 10.1002/hep.23505]
 - 31 **Hunt CM**, Lee TH, Morgan TR, Campbell S. One ALT Is Not Like the Other. *Gastroenterology* 2023; **165**: 320-323 [PMID: 37094663 DOI: 10.1053/j.gastro.2023.04.009]
 - 32 **Ceriotti F**, Hinzmann R, Panteghini M. Reference intervals: the way forward. *Ann Clin Biochem* 2009; **46**: 8-17 [PMID: 19103955 DOI: 10.1258/acb.2008.008170]
 - 33 **Panteghini M**, Adeli K, Ceriotti F, Sandberg S, Horvath AR. American Liver Guidelines and Cutoffs for "Normal" ALT: A Potential for Overdiagnosis. *Clin Chem* 2017; **63**: 1196-1198 [PMID: 28515102 DOI: 10.1373/clinchem.2017.274977]
 - 34 **Córdoba J**, O'Riordan K, Dupuis J, Borensztajn J, Blei AT. Diurnal variation of serum alanine transaminase activity in chronic liver disease. *Hepatology* 1998; **28**: 1724-1725 [PMID: 9890798 DOI: 10.1002/hep.510280640]
 - 35 **Miyake K**, Miyake N, Kondo S, Tabe Y, Ohsaka A, Miida T. Seasonal variation in liver function tests: a time-series analysis of outpatient data. *Ann Clin Biochem* 2009; **46**: 377-384 [PMID: 19641005 DOI: 10.1258/acb.2009.008203]
 - 36 **Pettersson J**, Hindorf U, Persson P, Bengtsson T, Malmqvist U, Werkström V, Ekelund M. Muscular exercise can cause highly pathological liver function tests in healthy men. *Br J Clin Pharmacol* 2008; **65**: 253-259 [PMID: 17764474 DOI: 10.1111/j.1365-2125.2007.03001.x]
 - 37 **Dufour DR**, Lott JA, Nolte FS, Gretch DR, Koff RS, Seeff LB. Diagnosis and monitoring of hepatic injury. I. Performance characteristics of laboratory tests. *Clin Chem* 2000; **46**: 2027-2049 [PMID: 11106349 DOI: 10.1093/clinchem/46.12.2027]
 - 38 **Preuss HG**, Kaats GR, Mrvichin N, Bagchi D, Preuss JM. Circulating ALT Levels in Healthy Volunteers Over Life-Span: Assessing Aging Paradox and Nutritional Implications. *J Am Coll Nutr* 2019; **38**: 661-669 [PMID: 31075051 DOI: 10.1080/07315724.2019.1580169]
 - 39 **Dong MH**, Bettencourt R, Brenner DA, Barrett-Connor E, Loomba R. Serum levels of alanine aminotransferase decrease with age in longitudinal analysis. *Clin Gastroenterol Hepatol* 2012; **10**: 285-90.e1 [PMID: 22020064 DOI: 10.1016/j.cgh.2011.10.014]
 - 40 **Moon SS**, Lee YS, Kim SW. Association of nonalcoholic fatty liver disease with low bone mass in postmenopausal women. *Endocrine* 2012; **42**: 423-429 [PMID: 22407492 DOI: 10.1007/s12020-012-9639-6]
 - 41 **Cui R**, Sheng H, Rui XF, Cheng XY, Sheng CJ, Wang JY, Qu S. Low bone mineral density in chinese adults with nonalcoholic Fatty liver disease. *Int J Endocrinol* 2013; **2013**: 396545 [PMID: 23983685 DOI: 10.1155/2013/396545]
 - 42 **Xia MF**, Lin HD, Yan HM, Bian H, Chang XX, Zhang LS, He WY, Gao X. The association of liver fat content and serum alanine aminotransferase with bone mineral density in middle-aged and elderly Chinese men and postmenopausal women. *J Transl Med* 2016; **14**: 11 [PMID: 26791408 DOI: 10.1186/s12967-016-0766-3]
 - 43 **Zoncapè M**, Liguori A, Tsochatzis EA. Non-invasive testing and risk-stratification in patients with MASLD. *Eur J Intern Med* 2024; **122**: 11-19 [PMID: 38246813 DOI: 10.1016/j.ejim.2024.01.013]
 - 44 **Afdhal NH**. Fibroscan (transient elastography) for the measurement of liver fibrosis. *Gastroenterol Hepatol (N Y)* 2012; **8**: 605-607 [PMID: 23483859]
 - 45 **Dulai PS**, Sirlin CB, Loomba R. MRI and MRE for non-invasive quantitative assessment of hepatic steatosis and fibrosis in NAFLD and NASH: Clinical trials to clinical practice. *J Hepatol* 2016; **65**: 1006-1016 [PMID: 27312947 DOI: 10.1016/j.jhep.2016.06.005]
 - 46 **Rangwani S**, Ardeshta DR, Mumtaz K, Kelly SG, Han SY, Krishna SG. Update on endoscopic ultrasound-guided liver biopsy. *World J Gastroenterol* 2022; **28**: 3586-3594 [PMID: 36161047 DOI: 10.3748/wjg.v28.i28.3586]
 - 47 **Diehl DL**. Top tips regarding EUS-guided liver biopsy. *Gastrointest Endosc* 2022; **95**: 368-371 [PMID: 34678299 DOI: 10.1016/j.gie.2021.10.016]
 - 48 **Han AL**. Validation of fatty liver index as a marker for metabolic dysfunction-associated fatty liver disease. *Diabetol Metab Syndr* 2022; **14**: 44 [PMID: 35317824 DOI: 10.1186/s13098-022-00811-2]
 - 49 **Reinshagen M**, Kabisch S, Pfeiffer AFH, Spranger J. Liver Fat Scores for Noninvasive Diagnosis and Monitoring of Nonalcoholic Fatty Liver Disease in Epidemiological and Clinical Studies. *J Clin Transl Hepatol* 2023; **11**: 1212-1227 [PMID: 37577225 DOI: 10.14218/JCTH.2022.00019]
 - 50 **Castellana M**, Donghia R, Guerra V, Procino F, Lampignano L, Castellana F, Zupo R, Sardone R, De Pergola G, Romanelli F, Trimboli P, Giannelli G. Performance of Fatty Liver Index in Identifying Non-Alcoholic Fatty Liver Disease in Population Studies. A Meta-Analysis. *J Clin Med* 2021; **10** [PMID: 33925992 DOI: 10.3390/jcm10091877]
 - 51 **Lee JH**, Kim D, Kim HJ, Lee CH, Yang JI, Kim W, Kim YJ, Yoon JH, Cho SH, Sung MW, Lee HS. Hepatic steatosis index: a simple screening tool reflecting nonalcoholic fatty liver disease. *Dig Liver Dis* 2010; **42**: 503-508 [PMID: 19766548 DOI: 10.1016/j.dld.2009.08.002]

- 52 **Buzzetti E**, Lombardi R, De Luca L, Tsochatzis EA. Noninvasive Assessment of Fibrosis in Patients with Nonalcoholic Fatty Liver Disease. *Int J Endocrinol* 2015; **2015**: 343828 [PMID: 26064107 DOI: 10.1155/2015/343828]
- 53 **Pan JJ**, Fallon MB. Gender and racial differences in nonalcoholic fatty liver disease. *World J Hepatol* 2014; **6**: 274-283 [PMID: 24868321 DOI: 10.4254/wjh.v6.i5.274]
- 54 **Zhang X**, Wong GL, Wong VW. Application of transient elastography in nonalcoholic fatty liver disease. *Clin Mol Hepatol* 2020; **26**: 128-141 [PMID: 31696690 DOI: 10.3350/cmh.2019.0001n]
- 55 **Mikolasevic I**, Orlic L, Franjic N, Hauser G, Stimac D, Milic S. Transient elastography (FibroScan®) with controlled attenuation parameter in the assessment of liver steatosis and fibrosis in patients with nonalcoholic fatty liver disease - Where do we stand? *World J Gastroenterol* 2016; **22**: 7236-7251 [PMID: 27621571 DOI: 10.3748/wjg.v22.i32.7236]
- 56 **Sanyal AJ**, Foucquier J, Younossi ZM, Harrison SA, Newsome PN, Chan WK, Yilmaz Y, De Ledinghen V, Costentin C, Zheng MH, Wai-Sun Wong V, Elkhashab M, Huss RS, Myers RP, Roux M, Labourdette A, Destro M, Fournier-Poizat C, Miette V, Sandrin L, Boursier J. Enhanced diagnosis of advanced fibrosis and cirrhosis in individuals with NAFLD using FibroScan-based Agile scores. *J Hepatol* 2023; **78**: 247-259 [PMID: 36375686 DOI: 10.1016/j.jhep.2022.10.034]
- 57 **Li D**, Zhang M, Wu S, Tan H, Li N. Risk factors and prediction model for nonalcoholic fatty liver disease in northwest China. *Sci Rep* 2022; **12**: 13877 [PMID: 35974018 DOI: 10.1038/s41598-022-17511-6]
- 58 **Yuan Y**, Xu M, Zhang X, Tang X, Zhang Y, Yang X, Xia G. Development and validation of a nomogram model for predicting the risk of MAFLD in the young population. *Sci Rep* 2024; **14**: 9376 [PMID: 38654043 DOI: 10.1038/s41598-024-60100-y]
- 59 **Qin S**, Hou X, Wen Y, Wang C, Tan X, Tian H, Ao Q, Li J, Chu S. Machine learning classifiers for screening nonalcoholic fatty liver disease in general adults. *Sci Rep* 2023; **13**: 3638 [PMID: 36869105 DOI: 10.1038/s41598-023-30750-5]
- 60 **Mohamed RA**, Nabih MI, Elshobaky MB, Khattab HM. The value of noninvasive scoring systems for the diagnosis of advanced fibrosis in Egyptian patients with nonalcoholic fatty liver disease. *Egypt J Intern Med* 2014; **26**: 162-169 [DOI: 10.4103/1110-7782.148151]
- 61 **Alqahtani SA**, Golabi P, Paik JM, Lam B, Moazez AH, Elariny HA, Goodman Z, Younossi ZM. Performance of Noninvasive Liver Fibrosis Tests in Morbidly Obese Patients with Nonalcoholic Fatty Liver Disease. *Obes Surg* 2021; **31**: 2002-2010 [PMID: 33616848 DOI: 10.1007/s11695-020-04996-1]
- 62 **Bertot LC**, Jeffrey GP, de Boer B, Wang Z, Huang Y, Garas G, MacQuillan G, Wallace M, Smith BW, Adams LA. Comparative Accuracy of Clinical Fibrosis Markers, Hepascore and Fibroscan® to Detect Advanced Fibrosis in Patients with Nonalcoholic Fatty Liver Disease. *Dig Dis Sci* 2023; **68**: 2757-2767 [PMID: 36947289 DOI: 10.1007/s10620-023-07896-3]
- 63 **Chen Y**, Du X, Kuppa A, Feitosa MF, Bielak LF, O'Connell JR, Musani SK, Guo X, Kahali B, Chen VL, Smith AV, Ryan KA, Eirksdottir G, Allison MA, Bowden DW, Budoff MJ, Carr JJ, Chen YI, Taylor KD, Oliveri A, Correa A, Crudup BF, Kardia SLR, Mosley TH Jr, Norris JM, Terry JG, Rotter JI, Wagenknecht LE, Halligan BD, Young KA, Hokanson JE, Washko GR, Gudnason V, Province MA, Peyser PA, Palmer ND, Speliotes EK. Genome-wide association meta-analysis identifies 17 loci associated with nonalcoholic fatty liver disease. *Nat Genet* 2023; **55**: 1640-1650 [PMID: 37709864 DOI: 10.1038/s41588-023-01497-6]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

