

**Name of journal: World Journal of Hepatology**

**ESPS Manuscript NO: 24060**

**Manuscript Type: ORIGINAL ARTICLE**

Observational Study

**Combined ARFI, APRI and forns index assessment for hepatic fibrosis  
grading in hepatitis B**

**Chang-Feng Dong, Jia Xiao, Ling-Bo Shan, Han-Ying Li, Yong-Jia Xiong,  
Gui-Lin Yang, Jing Liu, Si-Min Yao, Sha-Xi Li, Xiao-Hua Le, Jing Yuan,  
Bo-Ping Zhou, George L Tipoe, Ying-Xia Liu**

### Match Overview

**1** **CrossCheck** 76 words  
Liu, Yingxia, Chang-feng Dong, Guilin Yang, Jing Liu, Simi  
n Yao, Han-ying Li, Jing Yuan, Shaxi Li, Xiaohua Le, Yimin

2%



Combined ARFI and APRI assessment shows enhanced diagnostic accur



全部 新闻 图片 视频 更多 搜索工具

找到约 3,610 条结果 (用时 0.87 秒)

### Non-invasive diagnosis of liver fibrosis in chronic hepatitis C

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > PubMed Central (PMC) ▼ 翻译此页

作者: L de Lucca Schiavon - 2014 - 被引用次数: 18 - 相关文章

2014年3月21日 - Noninvasive markers of liver fibrosis can be divided into two groups: serum ... significant fibrosis as compared to APRI and the Forns index[50]. ... the accuracy of TE in patients with chronic hepatitis B virus (HBV) infection[83,91-94]. ... of fibrosis between ARFI results and histological assessment in 31.7% ...

### Non-invasive assessment of liver fibrosis

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > PubMed Central (PMC) ▼ 翻译此页

作者: V Papastergiou - 2012 - 被引用次数: 34 - 相关文章

However, noninvasive tests used to detect significant fibrosis and cirrhosis, ... The above data show only a moderate degree of accuracy of APRI for ... More recently this marker has been assessed in patients with chronic hepatitis B (CHB) [49,50] ... with CHC, ARFI was more accurate than TE for the noninvasive staging of ...

### Non-Invasive markers for hepatic fibrosis | BMC ...

[bmcgastroenterol.biomedcentral.com/.../10.../1471-230X-11-9...](http://bmcgastroenterol.biomedcentral.com/.../10.../1471-230X-11-9...) ▼ 翻译此页

作者: A Baranova - 2011 - 被引用次数: 140 - 相关文章

Current non-invasive methods range from serum biomarker assays to ... The rate of adoption of non-invasive diagnostic tests for liver fibrosis differs from country ... Other

Combined ARFI, APRI and forns index assessment for hepatic fibrosis grading



全部 图片 视频 新闻 购物 地图 图书

找到约 5,410 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Non-invasive assessment of liver fibrosis](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC3959378/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3959378/)

In parallel with this, noninvasive **assessment of liver fibrosis** has experienced .... this marker has been **assessed** in patients with chronic **hepatitis B** (CHB) [49,50] with .... Sebastiani et al proposed a sequential algorithm based on **combining APRI** .... **combination** of the Lok **Index** and **Forn's score** was efficient to exclude ( NPV > ...

[Noninvasive assessment of liver fibrosis in patients with chronic ...](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC4161791/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4161791/)

14 Sep 2014 ... Infection with **hepatitis B** virus is an important health problem worldwide: it .... **assessment of liver fibrosis**: serum biomarkers, **combined** indices or scores, ... **hepatitis B** were better for the Fibrotest than for the **APRI and Forns index**. .... accuracy of **ARFI** for the histological **staging of liver fibrosis** in patients with ...

[Acoustic radiation force impulse elastography, FibroScan®, Forns ...](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC4394958/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4394958/)

4 Feb 2015 ... Eighty-one patients who had been diagnosed with **hepatitis B** were recruited and ... The **combination** of FI with **ARFI/FibroScan** increased the predictive accuracy with a ... **ARFI** was shown to be as effective in the diagnosis of **liver fibrosis** as ... The **Forns' index** (FI) is based on the platelet count,  $\gamma$ -glutamyl ...

[Non-invasive diagnosis of liver fibrosis in chronic hepatitis C - NCBI](#)

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 5,600 条结果 (用时 0.69 秒)

### Noninvasive assessment of liver fibrosis in patients with ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: M Enomoto - 2014 - 被引用次数: 8 - [相关文章](#)

2014年9月14日 - Infection with **hepatitis B** virus is an important health problem worldwide: it ... **assessment of liver fibrosis**: serum biomarkers, **combined** indices or scores, ... **hepatitis B** were better for the Fibrotest than for the **APRI and Forns index**. .... accuracy of **ARFI** for the histological **staging of liver fibrosis** in patients with ...

### Non-invasive assessment of liver fibrosis

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: V Papastergiou - 2012 - 被引用次数: 33 - [相关文章](#)

**Indices combining** indirect and direct markers of **liver fibrosis** ... The ALT to platelets ratio **index (APRI)** is calculated as (AST/ upper limit of ... More recently this marker has been **assessed** in patients with chronic **hepatitis B** (CHB) [49,50] with .... **combination** of the Lok **Index** and **Forn's score** was efficient to exclude (NPV > ...

### Acoustic radiation force impulse elastography, FibroScan ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: DAORAN DONG - 2015 - [相关文章](#)

2015年2月4日 - Eighty-one patients who had been diagnosed with **hepatitis B** were recruited and ... The **combination** of FI with **ARFI/FibroScan** increased the predictive accuracy with a ... **ARFI** was shown to be as effective in the diagnosis of **liver fibrosis** as ... The **Forns' index** (FI) is based on the platelet count,  $\gamma$ -glutamyl ...

### Non-Invasive markers for hepatic fibrosis | BMC ...

[bmcgastroenterol.biomedcentral.com/.../10.../1471-230X-11-9...](http://bmcgastroenterol.biomedcentral.com/.../10.../1471-230X-11-9...) ▾ [翻译此页](#)

作者: A Baranova - 2011 - 被引用次数: 149 - [相关文章](#)

The rate of adoption of non-invasive diagnostic tests for **liver fibrosis** differs from ....