



Supplementary Figure 1 Selection of Cut-off values for the number of cleared lymph nodes.

Supplementary Table 1 Clinicopathological information of the study population

Variables	Level	Before PSM			After PSM		
		≤ 21 (n = 664)	> 21 (n = 378)	P value	≤ 21 (n = 203)	> 21 (n = 203)	P value
Sex (%)	Male	517 (77.86)	303 (80.16)	0.43	159 (78.33)	158	1
	Female	147 (22.14)	75 (19.84)		44 (21.67)	45 (22.17)	
Age (yr), median (IQR)		60.0 (55.0, 66.0)	60.0 (54.0, 66.0)	0.45	60.0 (55.0, 66.0)	60.0 (54.0, 66.0)	0.404
T (%)	1	77 (11.60)	21 (5.56)	< 0.05	13 (6.40)	16 (7.88)	0.951
	2	148 (22.29)	71 (18.78)		42 (20.69)	41 (20.20)	
	3	307 (46.23)	180 (47.62)		89 (43.84)	87 (42.86)	
	4	132 (19.88)	106 (28.04)		59 (29.06)	59 (29.06)	
N (%)	0	414 (62.35)	198 (52.38)	< 0.05	195 (96.06)	198	0.545
	1	149 (22.44)	85 (22.49)		5 (2.46)	4 (1.97)	
	2	67 (10.09)	67 (17.72)		2 (0.99)	0 (0.00)	
	3	34 (5.12)	28 (7.41)		1 (0.49)	1 (0.49)	
G (%)	Well-differentiated	79 (11.90)	45 (11.90)	0.42	29 (14.29)	31 (15.27)	0.788
	Moderately differentiated	464 (69.88)	276 (73.02)		148 (72.91)	142	
	Poorly differentiated	121 (18.22)	57 (15.08)		26 (12.81)	30 (14.78)	
Venous thrombus (%)	Negative	560 (84.34)	327 (86.51)	0.39	196 (96.55)	196	1
	Positive	104 (15.66)	51 (13.49)		7 (3.45)	7 (3.45)	
Perineurial invasion (%)	Negative	364 (77.45)	145 (69.38)	< 0.05	98 (80.33)	78 (78.00)	0.795
	Positive	106 (22.55)	64 (30.62)		24 (19.67)	22 (22.00)	
NPLN, mean (SD)		0.91 (1.75)	1.62 (2.67)	< 0.05	0.044	0.020	0.314

				(0.319)	(0.139)	
RPLN, mean (SD)		0.07 (0.13)	0.06 (0.09)	0.17	0.002	0.001 0.32
				(0.015)	(0.007)	
Adjuvant chemotherapy	No	414 (62.35)	185 (48.94)	< 0.05	76 (37.44)	79 (38.92)
	Yes	250 (37.65)	193 (51.06)		127 (62.56)	144 0.092
	(%)					(70.94)

NPLN: Number of positive lymph nodes, RPLN: Ratio of positive lymph nodes.