S W J

World Journal of Gastrointestinal Surgery

Submit a Manuscript: https://www.f6publishing.com

World J Gastrointest Surg 2024 December 27; 16(12): 3862-3869

DOI: 10.4240/wjgs.v16.i12.3862

ISSN 1948-9366 (online)

CASE REPORT

Pathological diagnosis and clinical feature analysis of descending duodenal mucosal adenocarcinoma: A case report

Jin-Yuan Zhang, Liu-Sheng Wu, Jun Yan, Qiang Jiang, Xiao-Qiang Li

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification Scientific Quality: Grade B, Grade B. Grade C Novelty: Grade B, Grade B, Grade B Creativity or Innovation: Grade B, Grade B, Grade B Scientific Significance: Grade A, Grade B, Grade B

P-Reviewer: Ghannam WM; Hameed Y

Received: May 8, 2024 Revised: September 19, 2024 Accepted: October 18, 2024 Published online: December 27, 2024

Processing time: 203 Days and 1.2 Hours



Jin-Yuan Zhang, Xiao-Qiang Li, Department of Thoracic Surgery, Peking University Shenzhen Hospital, Shenzhen 518036, Guangdong Province, China

Liu-Sheng Wu, Jun Yan, School of Medicine, Tsinghua University, Beijing 100084, China

Qiang Jiang, Department of Anorectal, Dazu District People's Hospital, Chongqing 402360, China

Co-first authors: Jin-Yuan Zhang and Liu-Sheng Wu.

Co-corresponding authors: Qiang Jiang and Xiao-Qiang Li.

Corresponding author: Xiao-Qiang Li, PhD, Professor, Department of Thoracic Surgery, Peking University Shenzhen Hospital, No. 1120 Lianhua Road, Futian District, Shenzhen 518036, Guangdong Province, China. dr.lixiaoqiang@gmail.com

Abstract

BACKGROUND

Mucosal adenocarcinoma of the descending duodenum is a very rare gastrointestinal tumor. Due to its low incidence, it has rarely been the focus of clinical and pathological studies. The clinical manifestations of these tumors are usually nonspecific, and they are easily misdiagnosed or missed. Pathological diagnosis is the gold standard for diagnosis, but due to the small number of cases, the relevant pathological characteristics and diagnostic criteria are not completely clear. The purpose of this study was to deepen the understanding of the diagnosis and treatment of this disease and to provide a clinical guidance.

CASE SUMMARY

A 61-year-old woman who was hospitalized with recurrent abdominal pain for more than 20 days. The patient developed epigastric pain with no obvious cause more than 20 days prior, mainly left epigastric pain and middle epigastric pain, and presented persistent dull pain without nausea or vomiting, fever or chills. The patient was treated at a local hospital, gastroscopy revealed a new lesion in the circum-intestinal cavity in the descending part of the duodenum, and pathological biopsy revealed mucous adenocarcinoma in the descending part of the duodenum. Currently, for further diagnosis and treatment, the patient is admitted to our hospital for surgical treatment for "malignant tumor of the duodenum" in the outpatient department.

WJGS https://www.wjgnet.com

CONCLUSION

Mucosal adenocarcinoma of the descending duodenum has a high misdiagnosis rate and missed diagnosis rate, clinical manifestations lack specificity, and pathological diagnosis is the main basis for diagnosis.

Key Words: Descending duodenal mucosal adenocarcinoma; Pathological diagnosis; Clinical features; Rare tumor; Case report

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The pathological diagnosis and clinical features of a rare case of descending duodenal mucosa adenocarcinoma were analyzed. Through detailed pathological examination, we confirmed the histological type of the tumor, combined with the patient's clinical manifestations, imaging findings and treatment response, and comprehensively analyzed the pathogenesis, progression and prognosis of the case. The research focuses on revealing the diagnostic difficulties and treatment challenges of such rare tumors, aiming to provide references for clinicians and improve the level of diagnosis and treatment of similar cases.

Citation: Zhang JY, Wu LS, Yan J, Jiang Q, Li XQ. Pathological diagnosis and clinical feature analysis of descending duodenal mucosal adenocarcinoma: A case report. World J Gastrointest Surg 2024; 16(12): 3862-3869 URL: https://www.wjgnet.com/1948-9366/full/v16/i12/3862.htm DOI: https://dx.doi.org/10.4240/wjgs.v16.i12.3862

INTRODUCTION

Mucosal adenocarcinoma of the descending duodenum is a rare malignant tumor of the digestive system, and its incidence is low; therefore, it is often neglected in clinical and pathological studies[1-3]. At present, the pathological features, clinical manifestations and treatment strategies are not completely clear. The diagnosis of this rare patient is often challenging because of its lack of clinical specificity, and this patient can easily be misdiagnosed or missed[4]. Given the relatively limited number of reports of this case, it is particularly important to study the pathological features and clinical manifestations of this rare tumor[5-7].

Through detailed pathological analysis and clinical evaluation of a patient with descending duodenal mucosal adenocarcinoma, this study comprehensively elucidated the characteristics of the disease and defined its diagnosis and treatment strategy. We will focus on the pathological features of the disease, including histological morphology, immunohistochemical expression and molecular biological features, to provide a more reliable basis for its diagnosis[8-10]. At the same time, we will conduct a comprehensive analysis of the clinical manifestations of the patients, including symptoms, signs, imaging and laboratory examination results, to explore the clinical characteristics and diagnostic difficulties of the disease[11]. Through the in-depth study of this case, we hope to provide clinicians with a more comprehensive and accurate diagnostic basis and a more effective treatment strategy. It is highly important to improve the diagnosis and treatment of rare diseases and improve the prognosis of patients[12]. Through in-depth study of this rare disease, we can also better understand its pathogenesis and tumor biological characteristics and provide new ideas and directions for future disease treatment and research. Through detailed analysis of the pathological features and clinical manifestations of descending duodenal mucosal adenocarcinoma, this study comprehensively elucidated the characteristics of this disease, clarified its diagnosis and treatment strategies, and provided a more accurate diagnostic basis and treatment plan for clinical treatment to improve the prognosis and quality of life of patients.

CASE PRESENTATION

Chief complaints

There was recurrent abdominal pain for more than 20 days.

History of present illness

The patient developed epigastric pain with no obvious cause more than 20 days ago, mainly in the left epigastric and middle epigastric pain, presenting persistent dull pain, no nausea, no fever, chills, no yellow all over the body, dark urine color, black stool and other discomfort. The patient was treated in a local hospital, and gastroscopic biopsy revealed mucous adenocarcinoma of the descending part of the duodenum. Currently, the patient is seeking further treatment at our hospital. The outpatient department plans to admit "malignant tumor of the duodenum" for surgical treatment.

History of past illness

The patient had a history of hypertension, regular oral antihypertensive medication treatment, and well-controlled blood pressure.



WJGS | https://www.wjgnet.com

Personal and family history

Nothing special.

Physical examination

There was upper abdominal tenderness; no rebound pain or muscle tension; no obvious mass in the abdomen; no liver, spleen or subcostal region; Murphy's sign (-); and no hepatic buckle pain (-).

Laboratory examinations

Histology revealed that the tumor cells exhibited a glandular tubular structure, abnormal enlargement of the nucleus, and nuclear division. Hematoxylin and eosin staining revealed deep staining of cell nuclei and uneven superficial staining of the cytoplasm. Immunohistochemical staining revealed that CK20 and CDX2 were positively expressed in tumor tissues, while CK7, CK19, carcinoembryonic antigen (CEA), carbohydrate antigen (CA) 19-9 and other markers were weakly or negatively expressed. In addition, the Ki-67 index was high, indicating that the tumor had strong proliferative activity. Combining the pathological features and immunohistochemical results, we concluded that the patient was diagnosed with adenocarcinoma of the descending duodenum mucosa. These features help to further understand the biological characteristics of this rare tumor and provide important guidance for its clinical diagnosis and treatment (Figure 1).

Imaging examinations

Abdominal computed tomography examination revealed that the intestinal wall was slightly thickened at the junction of the descending and horizontal parts of the duodenum, and the surrounding fat space was slightly blurred (Figure 2). Abdominal magnetic resonance imaging revealed that the intestinal wall of the descending duodenum was significantly thickened and that the signal was abnormal and tended to indicate neoplastic lesions; additionally, medical examination revealed secondary intrahepatic and external bile duct dilatation, hyperdilatation of the gallbladder, and slight dilatation of the pancreatic duct (Figure 3). Electronic proctoscopy revealed that new organisms were growing in the descending part of the duodenum, and the biopsy elasticity was poor (Figure 4).

FINAL DIAGNOSIS

The pathological diagnosis was descending duodenal mucosal adenocarcinoma.

TREATMENT

We recommend that patients continue to receive treatment after surgery and undergo regular follow-up observations.

OUTCOME AND FOLLOW-UP

The patient recovered after the operation, and no recurrence was found after 3 months.

DISCUSSION

Mucosal adenocarcinoma of the descending part of the duodenum is a rare malignant tumor of the digestive system, and its pathological features are somewhat heterogeneous compared with those of other digestive system tumors[13-15]. In this study, we observed that the histological morphology of the tumor showed a glandular tubular structure, increased nuclear atypia, and obvious mitotic features. These features are similar to those of other adenocarcinomas of the digestive system but also have their own characteristics[16]. We noted that the clinical manifestations of this patient lacked specificity, and this patient was easily misdiagnosed or missed. This is similar to the clinical manifestations of other intestinal tumors, including abdominal pain, indigestion, and weight loss. Therefore, clinicians should be vigilant in the face of such cases, combined with clinical manifestations, imaging examination, and timely pathological examination, to make a clear diagnosis[17-19].

In terms of diagnosis, we emphasized the importance of pathological diagnosis in this disease. Immunohistochemical staining revealed positive expression of CK20 and CDX2, while the expression of CK7, CK19, CEA, CA19-9 and other markers was weak or negative, which helped to rule out the possibility of other tumors and further confirmed the diagnosis of mucosal adenocarcinoma in the descending part of the duodenum[20-22]. Because this patient had a certain degree of invasion and metastasis, active treatment was needed. Surgical resection is the preferred treatment, but for advanced cases, chemoradiotherapy is also necessary[23]. In addition, we emphasize the importance of a comprehensive treatment strategy, including a combination of surgery, chemoradiotherapy, and targeted therapy, to improve treatment outcomes and patient survival[24].



Figure 1 Pathological features of descending duodenal mucosal adenocarcinoma.



Figure 2 Results of plain computed tomography scan of the patient's lower abdomen.

In terms of clinical features, we observed a lack of specificity in the symptoms of this patient, including abdominal pain, indigestion, weight loss, *etc.*, which are similar to those of other intestinal tumors and can easily lead to misdiagnosis or missed diagnosis. Epidemiologically, adenocarcinoma of the descending duodenum is a rare disease with a low incidence and has rarely become the focus of clinical and pathological studies[25-27]. Pathological analysis revealed that the tumor tissue of this patient presented a glandular tubular structure, increased nuclear atypia, and obvious mitosis, similar to other digestive system adenocarcinomas but also had its own characteristics. In terms of immunohistochemistry, positive CK20 and CDX2 expression was detected, while weak or negative CK7, CK19, CEA, and CA19-9 expression was detected, which helped to further clarify the diagnosis. Because this patient had a certain degree of invasion and metastasis, active treatment measures were taken[28]. Surgical resection is the preferred treatment, but for Zhang JY et al. Diagnosis of descending duodenal mucosal adenocarcinoma



Figure 3 Magnetic resonance imaging images of the patient's lower abdomen. A: Cross section and side view; B: Anterior coronal plane; C: Posterior coronal plane.

Baisbideng® WJGS | https://www.wjgnet.com

December 27, 2024 Volume 16 Issue 12



Figure 4 Electronic proctoscopy image results. Arrow: Lesion of coronal plane.

advanced cases, chemoradiotherapy is also necessary. The importance of comprehensive treatment strategies, including surgery, chemoradiotherapy and targeted therapy, has been highlighted to improve the therapeutic efficacy and survival rate of patients.

CONCLUSION

This study is highly important for further understanding the pathological features and clinical manifestations of rare adenocarcinomas of the descending duodenum mucosa. Through pathological analysis, clinical evaluation and discussion of treatment strategies, we provide clinicians with more accurate diagnosis and treatment options to improve patient prognosis and quality of life.

FOOTNOTES

Author contributions: Wu LS wrote the manuscript; Zhang JY analysed the manuscript; Yan J collected the data; Li XQ and Jiang Q guided the study. Wu LS is responsible for the design of the entire research scheme, the conception of the research scheme and the writing of the manuscript; Zhang JY is responsible for the collation and analysis of the case literature and the follow-up of the case patients under the guidance of Wu LS's research ideas. Therefore, Wu LS and Zhang JY have the same contribution. Jiang Q and Li XQ are the main persons in charge of this study. Li XQ organizes the discussion and analysis of this case and guides the relevant work; Jiang Q is responsible for supervising the progress of the whole research process and providing guidance and financial support. Li XQ and Jiang Q have made significant contributions to this study. All authors reviewed, edited, and approved the final manuscript and revised it critically for important intellectual content, gave final approval of the version to be published, and agreed to be accountable for all aspects of the work.

Supported by China Scholarship Council, No. 202406210298; the Scientific Research Project of the Education Department of Anhui Province, No. YJS20210324; the Research and Development of Intelligent Surgical Navigation and Operating System for Precise Liver Resection, No. 2022ZLA006; and the National Natural Science Foundation of China, No. 81972829.



Baishideng® WJGS https://www.wjgnet.com

Informed consent statement: This study has obtained informed consent and signed treatment consent from patients and their families.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country of origin: China

ORCID number: Liu-Sheng Wu 0000-0001-9580-2257; Xiao-Qiang Li 0000-0002-9692-1458.

S-Editor: Wang IJ L-Editor: A P-Editor: Yuan YY

REFERENCES

- Abbass R, Rigaux J, Al-Kawas FH. Nonampullary duodenal polyps: characteristics and endoscopic management. Gastrointest Endosc 2010; 71: 754-759 [PMID: 20363416 DOI: 10.1016/j.gie.2009.11.043]
- 2 Matsumoto S, Yoshida Y. Selection of appropriate endoscopic therapies for duodenal tumors: an open-label study, single-center experience. World J Gastroenterol 2014; 20: 8624-8630 [PMID: 25024618 DOI: 10.3748/wjg.v20.i26.8624]
- 3 Goldner B, Stabile BE. Duodenal adenocarcinoma: why the extreme rarity of duodenal bulb primary tumors? Am Surg 2014; 80: 956-959 [PMID: 25264638]
- Nakayama A, Kato M, Masunaga T, Kubosawa Y, Hayashi Y, Mizutani M, Kiguchi Y, Sasaki M, Takatori Y, Matsuura N, Mutaguchi M, 4 Takabayashi K, Yahagi N. Differential diagnosis of superficial duodenal epithelial tumor and non-neoplastic lesion in duodenum by magnified endoscopic examination with image-enhanced endoscopy. J Gastroenterol 2022; 57: 164-173 [PMID: 35091754 DOI: 10.1007/s00535-021-01844-6
- 5 Wu L, Zheng Y, Liu J, Luo R, Wu D, Xu P, Wu D, Li X. Comprehensive evaluation of the efficacy and safety of LPV/r drugs in the treatment of SARS and MERS to provide potential treatment options for COVID-19. Aging (Albany NY) 2021; 13: 10833-10852 [PMID: 33879634 DOI: 10.18632/aging.202860
- Wu L, Zhong Y, Wu D, Xu P, Ruan X, Yan J, Liu J, Li X. Immunomodulatory Factor TIM3 of Cytolytic Active Genes Affected the Survival 6 and Prognosis of Lung Adenocarcinoma Patients by Multi-Omics Analysis. Biomedicines 2022; 10 [PMID: 36140350 DOI: 10.3390/biomedicines10092248
- Iwamuro M, Tanaka T, Okada H. Review of lymphoma in the duodenum: An update of diagnosis and management. World J Gastroenterol 7 2023; 29: 1852-1862 [PMID: 37032723 DOI: 10.3748/wjg.v29.i12.1852]
- Beger HG, Mayer B, Poch B. Duodenum-Preserving Pancreatic Head Resection for Benign and Premalignant Tumors-a Systematic Review 8 and Meta-analysis of Surgery-Associated Morbidity. J Gastrointest Surg 2023; 27: 2611-2627 [PMID: 37670106 DOI: 10.1007/s11605-023-05789-4]
- Hayashi H, Miyamoto H, Nakagawa S, Matsuno K, Miyata T, Tsukamoto M, Mima K, Nitta H, Tanaka Y, Baba H. Distinct approaches 9 (antecolic and retrocolic) according to tumor location in laparoscopic and endoscopic cooperative surgery for non-ampullary duodenum tumors. Surg Endosc 2023; 37: 6718-6726 [PMID: 37217687 DOI: 10.1007/s00464-023-10114-w]
- Wu L, Liu Q, Ruan X, Luan X, Zhong Y, Liu J, Yan J, Li X. Multiple Omics Analysis of the Role of RBM10 Gene Instability in Immune 10 Regulation and Drug Sensitivity in Patients with Lung Adenocarcinoma (LUAD). Biomedicines 2023; 11 [PMID: 37509501 DOI: 10.3390/biomedicines11071861]
- Zeng Y, Yao H, Jiang X, Tang X, Wang X. GL11-altered Mesenchymal Tumor Involving the Duodenum: Case Report and Literature Review. Int J Surg Pathol 2023; 31: 1538-1547 [PMID: 36855297 DOI: 10.1177/10668969231157782]
- Wu L, Zheng Y, Ruan X, Wu D, Xu P, Liu J, Wu D, Li X. Long-chain noncoding ribonucleic acids affect the survival and prognosis of 12 patients with esophageal adenocarcinoma through the autophagy pathway: construction of a prognostic model. Anticancer Drugs 2022; 33: e590-e603 [PMID: 34338240 DOI: 10.1097/CAD.00000000001189]
- Milione M, Parente P, Grillo F, Zamboni G, Mastracci L, Capella C, Fassan M, Vanoli A. Neuroendocrine neoplasms of the duodenum, 13 ampullary region, jejunum and ileum. Pathologica 2021; 113: 12-18 [PMID: 33686306 DOI: 10.32074/1591-951X-228]
- 14 Okano K, Oshima M, Suto H, Ando Y, Asano E, Kamada H, Kobara H, Masaki T, Suzuki Y. Ampullary carcinoma of the duodenum: current clinical issues and genomic overview. Surg Today 2022; 52: 189-197 [PMID: 33797636 DOI: 10.1007/s00595-021-02270-0]
- 15 Del Toro C, Cabrera-Aguirre A, Casillas J, Ivanovic A, Scortegagna E, Estanga I, Alessandrino F. Imaging spectrum of non-neoplastic and neoplastic conditions of the duodenum: a pictorial review. Abdom Radiol (NY) 2023; 48: 2237-2257 [PMID: 37099183 DOI: 10.1007/s00261-023-03909-x
- Wu L, Zhong Y, Yu X, Wu D, Xu P, Lv L, Ruan X, Liu Q, Feng Y, Liu J, Li X. Selective poly adenylation predicts the efficacy of 16 immunotherapy in patients with lung adenocarcinoma by multiple omics research. Anticancer Drugs 2022; 33: 943-959 [PMID: 35946526 DOI: 10.1097/CAD.00000000001319]
- Osama MA, Dhawan S, Khetan M, Rawat KS. Inflammatory myofibroblastic tumor of colon and duodenum: Metachronous, metastatic or 17 local recurrence? An unusual presentation. J Cancer Res Ther 2023; 19: S917-S920 [PMID: 38384079 DOI: 10.4103/jcrt.jcrt 258 22]



- Beger HG, Mayer B, Poch B. Resection of the duodenum causes long-term endocrine and exocrine dysfunction after Whipple procedure for 18 benign tumors - Results of a systematic review and meta-analysis. HPB (Oxford) 2020; 22: 809-820 [PMID: 31983660 DOI: 10.1016/j.hpb.2019.12.016]
- 19 Wu L, Li H, Liu Y, Fan Z, Xu J, Li N, Qian X, Lin Z, Li X, Yan J. Research progress of 3D-bioprinted functional pancreas and in vitro tumor models. Int J Bioprinting 2024; 10: 1256 [DOI: 10.36922/ijb.1256]
- Chaparro Mirete M, López-López V, Robles Campos R. Lymphoepithelioma-like carcinoma of the duodenum: a very infrequent tumor. Rev 20 Esp Enferm Dig 2020; 112: 239 [PMID: 32022569 DOI: 10.17235/reed.2020.6503/2019]
- Portale G, Mazzeo A, Zuin M, Spolverato Y, Cipollari C, Fiscon V. Fully Laparoscopic Pancreas-Preserving Resection of the Third and 21 Fourth Portion of the Duodenum for Adenocarcinomas and Gastrointestinal Stromal Tumors: Technical Report on a Case Series. J Laparoendosc Adv Surg Tech A 2022; 32: 466-470 [PMID: 34762524 DOI: 10.1089/lap.2021.0480]
- 22 Wu L, Li X, Qian X, Wang S, Liu J, Yan J. Lipid Nanoparticle (LNP) Delivery Carrier-Assisted Targeted Controlled Release mRNA Vaccines in Tumor Immunity. Vaccines (Basel) 2024; 12 [PMID: 38400169 DOI: 10.3390/vaccines12020186]
- Fukusada S, Shimura T, Iwasaki H, Okuda Y, Katano T, Nishigaki R, Ozeki T, Kitagawa M, Nishie H, Tanaka M, Ozeki K, Kubota E, Tanida 23 S, Kataoka H. Relationship between Immunophenotype and Clinicopathological Findings for Superficial Nonampullary Duodenal Epithelial Tumor. *Digestion* 2021; **102**: 870-877 [PMID: 33794536 DOI: 10.1159/000514812]
- 24 Boggi U. Laparoscopic duodenum-preserving total pancreatic head resection for pancreatic tumors: the difficult balance among overtreatment, ideal treatment, and undertreatment. Langenbecks Arch Surg 2022; 407: 3859-3861 [PMID: 35434751 DOI: 10.1007/s00423-022-02512-w]
- 25 Chen Y, Zhang H, Zhou Q, Lu L, Lin J. Metastases to duodenum in cervical squamous cell carcinoma: A case report and review of the literature. Medicine (Baltimore) 2022; 101: e28526 [PMID: 35029209 DOI: 10.1097/MD.00000000028526]
- Wu L, Chen X, Zeng Q, Lai Z, Fan Z, Ruan X, Li X, Yan J. NR5A2 gene affects the overall survival of LUAD patients by regulating the 26 activity of CSCs through SNP pathway by OCLR algorithm and immune score. Heliyon 2024; 10: e28282 [PMID: 38601554 DOI: 10.1016/j.heliyon.2024.e28282
- Köroğlu G, Görür GD, Doğan S, Bayrak BY, Demir H. Incidental Uptake of 68Ga-DOTATATE Gastrointestinal Stromal Tumor of 27 Duodenum. Clin Nucl Med 2022; 47: e289-e290 [PMID: 35025797 DOI: 10.1097/RLU.00000000004036]
- Liu W, Wang J, Ma L, Zhuang A, Xu J, He J, Yang H, Fang Y, Lu W, Zhang Y, Tong H. Which style of duodenojejunostomy is better after 28 resection of distal duodenum. BMC Surg 2022; 22: 409 [PMID: 36434558 DOI: 10.1186/s12893-022-01850-2]



WJGS | https://www.wjgnet.com



Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

