

Association between tourniquet use and blood loss during below-knee amputation

International Orthopaedics

Alden E. Wyland BS, Erik Woelber MD, Liam H. Wong BS, Jordan Arakawa BA, Zachary M. Working MD, James E. Meeker MD

Author information

Affiliations

Oregon Health and Sciences University (OHSU), Department of Orthopaedics and Rehabilitation, Portland, Oregon, USA

Erik Woelber, Zachary M. Working, and James E. Meeker

Oregon Health and Sciences University (OHSU), School of Medicine, Portland, Oregon, USA

Alden E. Wyland, Liam H. Wong, and Jordan Arakawa

Corresponding Author:

Correspondence to James E. Meeker

Email: meekerj@ohsu.edu

Supplemental Table 1. Variables collected during chart review

Category	Variables
Patient characteristics	Age, gender, body mass index (BMI), diabetes, sepsis, respiratory rate, temperature, dementia, psych disorder, bleeding disorder, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), hepatitis C, hepatic cirrhosis, cardiac issues, smoking, IV drug use, alcohol use, homelessness, outside hospital transfer, preoperative ventilator use, steroid use, American Society of Anesthesiologists (ASA) score, white blood cell count, platelets, creatinine
Surgery characteristics	Tourniquet use, indication, intraoperative units of packed red blood cells transfused
Hospital stay	Preoperative and postoperative hemoglobin, length of stay, unplanned reoperations, number of postoperative transfusions