

Supplementary Table 1 Sensitivity, specificity, and accuracy of parameters alone or in combination in predicting endoscopic healing in the ileum

Variables	Sensitivity (95%CI)	Specificity (95%CI)	Accuracy
LRG	0.767 (0.623-0.869)	0.787 (0.669-0.871)	0.779
CRP	0.465 (0.325-0.611)	0.869 (0.762-0.932)	0.702
Albumin	0.744 (0.598-0.851)	0.590 (0.465-0.705)	0.654
HBI	0.326 (0.205-0.475)	0.918 (0.822-0.965)	0.673
Combination			
LRG (+) and CRP (+)	0.442 (0.304-0.589)	0.902 (0.802-0.954)	0.712
LRG (+) or CRP (+)	0.791 (0.648-0.886)	0.754 (0.633-0.845)	0.769
LRG (+) and albumin (+)	0.605 (0.456-0.736)	0.853 (0.743-0.858)	0.750
LRG (+) or albumin (+)	0.907 (0.784-0.963)	0.525 (0.402-0.645)	0.683
LRG (+) and HBI (+)	0.256 (0.149-0.402)	0.984 (0.913-0.999)	0.683
LRG (+) or HBI (+)	0.837 (0.700-0.919)	0.721 (0.598-0.818)	0.769

CI: Confidence interval; LRG: Leucine-rich alpha-2 glycoprotein; LRG: Leucine-rich alpha-2 glycoprotein; CRP: C-reactive protein;

HBI: Harvey-Bradshaw index