

The intestinal epithelium, intraepithelial lymphocytes and the gut microbiota – key

全部 图片 视频 新闻 地图 图书

找到约 326,000 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Google 学术: The intestinal epithelium, intraepithelial lymphocytes and the gut microbiota – key players in the pathogenesis of celiac disease](#)

[... bacteria\) and the mucosal barrier in the pathogenesis ...](#) - Tlaskalová-Hogenová - 被引用次数: 322

[Pathogenesis of IBS: role of inflammation, immunity ...](#) - Öhman - 被引用次数: 422

[... B lymphocytes, microbiota and the intestinal epithelium ...](#) - Shulzhenko - 被引用次数: 194

[Novel players in coeliac disease pathogenesis: role of the gut ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5102016/>

Even though the bacterial load in the colon is markedly higher ... A comprehensive study of the gut microbiota using ... receptor (TCR) $\alpha\beta^+$ intraepithelial lymphocytes (IELs) ... from Paneth cells, $\gamma\delta$ TCR $^+$ IELs and epithelial cells. ... Thus, microbiota–host interactions are key in the ...

[Novel players in coeliac disease pathogenesis: role of the gut ...](#)

www.nature.com/nrgastro/journal/v12/n9/abs/nrgastro.2015.90.html

9 Jun 2015 ... Potential microbial modulation of coeliac disease pathogenesis. ... Umesaki, Y. Proliferative recruitment of intestinal intraepithelial lymphocytes ... The key role of segmented filamentous bacteria in the coordinated maturation of gut ... inhibit the toxic effects induced by wheat gliadin in epithelial cell culture.

[\[PDF\] Overview and Pathogenesis of Celiac Disease - Gastroenterology](#)

[www.gastrojournal.org/article/S0016-5085\(05\)00177-0/pdf](http://www.gastrojournal.org/article/S0016-5085(05)00177-0/pdf)

Celiac disease is characterized by small intestinal mu- ... The pathogenesis of disease involves ... ing * of intraepithelial lymphocytes (IEL) occurring at a ... squamoid surface epithelial cells, an increase in intraepi- ... duced by bacteria can digest these proline-rich gluten ... Genetics clearly play a key role in the pathogen-

[Novel players in coeliac disease pathogenesis: Role of the gut ...](#)

Name of journal: *World Journal of Gastroenterology*

Manuscript NO: 34724

Manuscript Type: REVIEW

Intestinal epithelium, intraepithelial lymphocytes, and the gut microbiota - key players in the pathogenesis of celiac disease

Bożena Cukrowska, Agnieszka Sowińska, Joanna Beata Bierła, Elżbieta Czarnowska, Anna Rybak, Urszula Grzybowska-Chlebowczyk

Abstract

Celiac disease (CD) is a chronic immune-mediated disorder triggered by the ingestion of gluten in genetically predisposed individuals. Before activating the

Match Overview

1	Internet 62 words crawled on 10-Dec-2012 bmlab.no	1%
2	Crossref 51 words Lee, Won-Jae, and Koji Hase. "Gut microbiota-genera ... d metabolites in animal health and disease". Nature Ch	1%
3	Internet 46 words crawled on 26-Nov-2016 pdfs.semanticscholar.org	1%
4	Crossref 36 words Kozakova, Hana, Martin Schwarzer, Ludmila Tuckova, Da gmar Srutkova, Elzbieta Czarnowska, Ilona Rosiak, Tom	1%
5	Internet 30 words crawled on 26-Dec-2016 onlinelibrary.wiley.com	1%
6	Internet 29 words crawled on 08-Jul-2016 mdpi.com	1%
7	Crossref 29 words Harris, Lucinda A., Jason Y. Park, Lysandra Voltaggio, an d Dora Lam-Himlin. "Celiac disease: clinical, endosco ...	1%
8	Internet 28 words crawled on 14-Oct-2015 www.science.gov	1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 109,000 条结果 (用时 0.60 秒)

Google 学术: Intestinal epithelium, intraepithelial lymphocytes, and the gut microbiota – key players in the pathogenesis of celiac disease

Pathogenesis of IBS: role of inflammation, immunity ... - Öhman - 被引用次数: 440

The role of gut microbiota (commensal bacteria) and ... - Tlaskalová-Hogenová - 被引用次数: 341

... B lymphocytes, microbiota and the intestinal epithelium ... - Shulzhenko - 被引用次数: 197

Novel players in coeliac disease pathogenesis: role of the gut microbiota

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5102016/> - 翻译此页

作者: EF Verdu - 2015 - 被引用次数: 46 - 相关文章

2015年6月9日 - Oral tolerance is a key feature of the gut immune system, whereby a state of ... T-cell receptor (TCR) $\alpha\beta^+$ intraepithelial lymphocytes (IELs) compared with colonized mice. ... The gut microbiota induces maturation of the gastrointestinal ... production from epithelial cells via IL-22 release from dendritic cells.

IL-15: a central regulator of celiac disease immunopathology

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4066219/> - 翻译此页

作者: V Abadie - 2014 - 被引用次数: 67 - 相关文章

Interestingly, because it is overexpressed both in the gut epithelium and in the lamina ... Since IL-15 was first proposed to play a key role in CD pathogenesis (4–6), ... In addition, IL-15 triggers other signaling pathways in T lymphocytes by In mice, IL-15 and IL-15R α expression on intestinal epithelial cells (IECs) was ...

Novel players in coeliac disease pathogenesis: role of the gut microbiota

www.nature.com/nrgastro/journal/v12/n9/abs/nrgastro.2015.90.html - 翻译此页

2015年6月9日 - Potential microbial modulation of coeliac disease pathogenesis. Umesaki, Y.

Proliferative recruitment of intestinal intraepithelial lymphocytes ... The key role of segmented filamentous bacteria in the coordinated maturation of gut ... inhibit the toxic effects induced by wheat

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 110,000 条结果 (用时 0.34 秒)

Google 学术 : Intestinal epithelium, intraepithelial lymphocytes, and the gut microbiota – key players in the pathogenesis of celiac disease

Pathogenesis of IBS: role of inflammation, immunity ... - Öhman - 被引用次数 : 436

The role of gut microbiota (commensal bacteria) and ... - Tlaskalová-Hogenová - 被引用次数 : 341

... B lymphocytes, microbiota and the intestinal epithelium ... - Shulzhenko - 被引用次数 : 197

Novel players in coeliac disease pathogenesis: role of the gut microbiota

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5102016/> - 翻译此页

作者 : EF Verdu - 2015 - 被引用次数 : 46 - 相关文章

2015年6月9日 - Oral tolerance is a key feature of the gut immune system, whereby a state of ... T-cell receptor (TCR) $\alpha\beta^+$ intraepithelial lymphocytes (IELs) compared with colonized mice. ... The gut microbiota induces maturation of the gastrointestinal ... production from epithelial cells via IL-22 release from dendritic cells.

IL-15: a central regulator of celiac disease immunopathology

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4066219/> - 翻译此页

作者 : V Abadie - 2014 - 被引用次数 : 67 - 相关文章

Interestingly, because it is overexpressed both in the gut epithelium and in the lamina ... Since IL-15 was first proposed to play a key role in CD pathogenesis (4–6), ... In addition, IL-15 triggers other signaling pathways in T lymphocytes by In mice, IL-15 and IL-15R α expression on intestinal epithelial cells (IECs) was ...

Novel players in coeliac disease pathogenesis: role of the gut microbiota

www.nature.com/nrgastro/journal/v12/n9/abs/nrgastro.2015.90.html - 翻译此页

2015年6月9日 - Potential microbial modulation of coeliac disease pathogenesis. Umesaki, Y.

Proliferative recruitment of intestinal intraepithelial lymphocytes ... The key role of segmented filamentous bacteria in the coordinated maturation of gut inhibit the toxic effects induced by wheat gliadin in epithelial cell culture.

Novel players in coeliac disease pathogenesis: Role of the gut ...

<https://www.researchgate.net/.../277958433> Novel players in coeliac disea... - 翻译此页