

World Journal of *Cardiology*

World J Cardiol 2024 April 26; 16(4): 168-216



EDITORIAL

- 168 Pacemaker post transcatheter aortic valve replacement: A multifactorial risk?
Noble S, Bendjelid K
- 173 Challenging situation of coronary artery anomaly associated with ischemia and/or risk of sudden death
Ito S
- 177 Transcatheter aortic valve replacement in low-risk young population: A double edge sword?
Bhogal S, Batta A
- 181 Hypoxia-inducible factor-1 α in myocardial infarction
Škrlec I, Kolomeichuk SN
- 186 Left bundle branch pacing set to outshine biventricular pacing for cardiac resynchronization therapy?
Batta A, Hatwal J

MINIREVIEWS

- 191 Aspirin interruption before neurosurgical interventions: A controversial problem
Kulikov A, Konovalov A, Pugnaroni PP, Bilotta F

SCIENTOMETRICS

- 199 The cardiovascular system at high altitude: A bibliometric and visualization analysis
Zhao ML, Lu ZJ, Yang L, Ding S, Gao F, Liu YZ, Yang XL, Li X, He SY

CORRECTION

- 215 Correction: Establishment of a prediction model for prehospital return of spontaneous circulation in out-of-hospital patients with cardiac arrest
Wang JJ, Zhou Q, Huang ZH, Han Y, Qin CZ, Chen ZQ, Xiao XY, Deng Z

ABOUT COVER

Peer Reviewer of *World Journal of Cardiology*, Ramachandra Barik, DNB, MD, Full Professor, Department of Cardiology, All India Institute of Medical Sciences, Bhubaneswar 751019, Odisha, India.
cardioramachandra@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Cardiology (WJC, World J Cardiol)* is to provide scholars and readers from various fields of cardiology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJC mainly publishes articles reporting research results and findings obtained in the field of cardiology and covering a wide range of topics including acute coronary syndromes, aneurysm, angina, arrhythmias, atherosclerosis, atrial fibrillation, cardiomyopathy, congenital heart disease, coronary artery disease, heart failure, hypertension, imaging, infection, myocardial infarction, pathology, peripheral vessels, public health, Raynaud's syndrome, stroke, thrombosis, and valvular disease.

INDEXING/ABSTRACTING

The WJC is now abstracted and indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJC as 1.9; IF without journal self cites: 1.8; 5-year IF: 2.3; Journal Citation Indicator: 0.33. The WJC's CiteScore for 2022 is 1.9 and Scopus CiteScore rank 2022: Cardiology and cardiovascular medicine is 226/354.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xiang Li; Cover Editor: Yun-Xiaojiao Wu.

NAME OF JOURNAL

World Journal of Cardiology

ISSN

ISSN 1949-8462 (online)

LAUNCH DATE

December 31, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Ramdas G Pai, Dimitrios Tousoulis, Marco Matteo Ciccone, Pal Pacher

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/1949-8462/editorialboard.htm>

PUBLICATION DATE

April 26, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Pacemaker post transcatheter aortic valve replacement: A multifactorial risk?

Stephane Noble, Karim Bendjelid

Specialty type: Cardiac and cardiovascular systems

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Jha AK, United States

Received: November 21, 2023

Peer-review started: November 21, 2023

First decision: January 24, 2024

Revised: February 1, 2024

Accepted: March 18, 2024

Article in press: March 18, 2024

Published online: April 26, 2024



Stephane Noble, Department of Medicine, Structural Heart Unit, University Hospital of Geneva, Geneva 1211, Switzerland

Karim Bendjelid, Department of Acute Medicine, Geneva University Hospitals, Geneva 1211, Switzerland

Corresponding author: Stephane Noble, MD, Assistant Professor, Department of Medicine, Structural Heart Unit, University Hospital of Geneva, 4 rue Gabrielle Perret Gentil, Geneva 1211, Switzerland. stephane.noble@hcuge.ch

Abstract

Pacemaker post-transcatheter aortic valve replacement is related to multifactorial risk. Nwaedozie *et al* brought to the body of evidence electrocardiogram and clinical findings. However, procedural characteristics have at least as much impact on the final need for a permanent pacemaker and potentially on the pacing rate. In this regard, long-term follow-up and understanding of the impact of long-term stimulation is of utmost importance.

Key Words: Transcatheter aortic valve replacement; Permanent pacemaker implantation; Conduction abnormalities; Right bundle branch block; Left bundle branch block

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Since the first transcatheter aortic valve replacement (TAVR) in 2002, TAVR has become a recognized alternative therapy to symptomatic severe aortic stenosis independently of the surgical risk score. The multiple iterations of the delivery systems and transcatheter heart valves (THV) over time associated with better patient assessment and the growing experience and expertise of the operators improved the procedural and follow-up results. However, despite the possibility of repositioning and partially recapturing some of the self-expanding THV and generally higher implantation targets, the need for a permanent pacemaker remains the most frequent complication post-procedure.

Citation: Noble S, Bendjelid K. Pacemaker post transcatheter aortic valve replacement: A multifactorial risk? *World J Cardiol* 2024; 16(4): 168-172

URL: <https://www.wjgnet.com/1949-8462/full/v16/i4/168.htm>

DOI: <https://dx.doi.org/10.4330/wjc.v16.i4.168>

INTRODUCTION

Since the first transcatheter aortic valve replacement (TAVR) in 2002, TAVR has become a recognized alternative therapy to symptomatic severe aortic stenosis independently of the surgical risk score. The multiple iterations of the delivery systems and transcatheter heart valves (THV) over time associated with better patient assessment and the growing experience and expertise of the operators improved the procedural and follow-up results[1-3]. However, despite the possibility of repositioning and partially recapturing some of the self-expanding THV and generally higher implantation targets, the need for a permanent pacemaker remains the most frequent complication post-procedure. Importantly, the left bundle branch travels commonly 2 to 3 mm below the base of the interleaflet triangle between the noncoronary and right coronary leaflets and is therefore at risk for interaction with the THV[4].

Conductance disturbances and the need for permanent pacemaker implantation post-TAVR are multifactorial and have an important clinical and economic impact (price of the pacemaker implantation, higher length of stay)[5]. In a multicenter European trial using balloon-expandable valves, conductance abnormalities were the second most common reason for prolonged hospitalization after logistic causes[6]. Delayed atrioventricular block can be seen up to 8 d post-TAVR in 7% of the cases, but patients without conduction abnormalities immediately post-TAVR did not present any delayed high-degree conduction disorder in a multicentric report including 1064 patients[7]. The European Society of Cardiology (ESC) guidelines on cardiac pacing gave a class I indication for permanent pacemaker implantation in the context of a complete atrioventricular block or new alternating bundle branch block and a class IIa for right bundle branch block and new conductance disturbance (PR prolongation or axis changes)[8]. Interestingly, a class IIa for ambulatory ECG monitoring or electrophysiology study was given in the setting of persistent new left bundle branch block > 150 ms or PR > 240 ms with no further prolongation during 48 h and a class IIb in the context of pre-existing conduction abnormalities with prolongation of > 20 ms of the QRS and PR interval.

In the present issue of the Journal, Nwaedozi *et al*[9] assessed the effects of baseline nonspecific interventricular conduction delay and supraventricular arrhythmia on post-TAVR permanent pacemaker need and also reported the impact of permanent pacemaker implantation on clinical outcomes at one year. In this regard, they retrospectively analyzed the single-center cohort of a tertiary hospital in central Wisconsin, United States involving 357 patients who underwent a TAVR (95.2% of transfemoral approach) between January 2012 and December 2019 using balloon-expandable and self-expanding THV in 53.8% and 46.2%, respectively.

One of the strengths of the study is that they analyzed the rate of pacemaker dependency at follow-up, which was as high as 78.9% one month post-pacemaker implantation. In addition, board-certified cardiologists reviewed the ECG. They found a permanent pacemaker rate of 16% at one year with no significant differences between self-expanding (17.6%) and balloon-expandable (14.6%) THV. Their pacemaker rates are similar to the Medtronic self-expanding THV in the Evolut Low-Risk trial (17.4% at 30 d and 19.4% at one year) and slightly above what we could expect with the Edwards SAPIEN 3 balloon-expandable valve (PARTNER 3 trial: 6.6% at 30 d and 7.5% at one year)[10,11]. The median time of implantation in the study by Nwaedozi *et al*[9] was 2 d, and half of the patients underwent pacemaker implantation within 48 h post-TAVR. Complete atrioventricular block was the predominant indication (66.7%) and the other indications were as follows: left ventricle dysfunction (10.5%), symptomatic bradycardia (8.8%), and symptomatic second atrioventricular block (1.8%).

The main findings of this trial are that pre-TAVR type II Diabetes Mellitus and QRS duration > 120 ms, regardless of the presence of bundle branch blocks were predictors of permanent pacemaker need post-TAVR. They also demonstrated a linear association between post-TAVR permanent pacemaker rate for every 20 ms prolongation of the QRS duration above 100 ms. Finally, at one year, there were more heart failure hospitalizations (28% *vs* 14%, $P = 0.022$) and myocardial infarction (9% *vs* 2%, $P = 0.031$) in the group requiring a permanent pacemaker.

The limitations of this report are related to the design of the study which is a retrospective analysis of a single center experience with a relatively small number of patients ($n = 357$) treated over 8 years. During this long period, there were multiple THV iterations, implantation technique refinements, and a regular expansion of the indication creating a heterogeneous population. The results have to be brought into perspective. Indeed, not only does the baseline ECG influence the post-procedural risk of permanent pacemaker need (particularly the presence of a right bundle branch block[5]), but also the left ventricular outflow tract (LVOT) anatomy and the calcium burden and repartition as well as the valve type used (*i.e.*, balloon-expandable *vs* self-expanding THV), and finally patient and procedural characteristics (*i.e.*, the height of implantation, percentage of oversizing, the technique of implantation, pre- and post-dilatation, resheathing and recapture).

Recently, the best clinical practices concerning the view of valve deployment have progressively switched from a three-cusp view to a combination of cusp overlap and three-cusp views, particularly for the self-expanding valves[12]. When using the cusp-overlap view, we focus on the non-coronary cusp which is on the left side of the screen, whereas the right and left cusps are superimposed on the right side of the screen. This view allows the elongation of the LVOT and subsequently a more precise height of implantation. It also contributes to eliminating the parallax of the delivery catheter, deploying the valve in a true coplanar view, and better aligning the THV commissures[12]. The cusp overlap technique

has been associated with a lower pacemaker implantation rate at 30 d than the conventional technique in a meta-analysis including 1227 Medtronic Evolut valves (cusp overlap technique: 641 *vs* co-planar view: 586).

In the cusp overlap technique, the implantation height was 1.03 mm higher and the incidence of pacemaker rate was 9.8% compared to 20.6% in the conventional technique[13]. However, the incidence of a left bundle branch block did not defer. In a propensity-matched analysis on a small Spanish cohort (92 patients in each group with no baseline characteristic differences), there was a significant reduction of new onset of left bundle branch block and reduced P wave and QRS widening at one year in the cusp overlap technique group compared to the conventional technique group[14]. There was also a significant reduction in a combined primary endpoint including pacemaker implantation, hospitalization and cardiovascular death at one year[14]. Recently, the interim analysis of the Optimize PRO TAVR study showed again the benefit of the cusp overlap technique in 400 patients[15]. This study reports the absence of moderate or severe paravalvular leak and the lowest pacemaker rate (9.8% at 30 d) in a multicenter prospective study with the Evolut platform using the cusp overlap technique and an “optimized TAVR care pathway”. The pacemaker rate was as low as 5.7% when the 4-step cusp overlap technique was precisely followed.

Importantly, Nwaedozie *et al*[9] did not report data on the procedural depth of THV implantation which is a major predictor of the need for a permanent pacemaker post-TAVR. To emphasize the role of the implantation technique and volume-outcome relationship, in a sub-analysis of the Evolut low-risk trial, there was a substantial variation in the rate of permanent pacemaker implantation from site to site in this study including 699 patients from 84 centers, with a lower rate of pacemaker need in high volume centers[3,16]. The sites with a low pacemaker rate had higher implantation at the non-coronary and left coronary sinus levels and fewer patients with an implantation depth of more than 5 mm[16].

Finally, the long-term data post-pacemaker implantation after TAVR are conflicting. Right ventricular pacing is associated with electromechanical dyssynchrony, left ventricular remodeling, increased risk of atrial fibrillation, and tricuspid regurgitation. In a series of 377 post-TAVR patients at one year, a stimulation rate > 40% of the time was associated in a propensity-matched analysis with a higher risk of cardiovascular mortality and hospitalization for heart failure[17]. More physiological pacing such as cardiac resynchronization in cases of reduced left ventricular ejection fraction is recommended to decrease the adverse outcome. His bundle or conduction system stimulation is also a more physiological pacing, which should be promoted. Of note, in the later series, 6 patients had His stimulation with no event or hospitalization during follow-up. In 2020, a meta-analysis including 30 studies showed a higher risk at one year of all-cause death and heart failure hospitalization after new onset of left bundle branch block or peri-procedural pacemaker implantation[18]. In addition, post TAVR new onset of left bundle branch block was associated with an increased risk of cardiac death and need for pacemaker implantation at one year[18]. In a recent meta-analysis of 31 studies with a mean follow-up of 22 months, new permanent pacemaker implantation after TAVR was associated with a higher risk of all-cause death and heart failure hospitalization[19]. Conversely, data from the nationwide, population cohort study with 3420 TAVR performed between 2008 and 2018 in Sweden, 481 (14.1%) required permanent pacemaker implantation within 30 d post procedure[20]. With a median follow-up of 2.7 years, there was no difference in long-term survival between patients who were or were not implanted with a permanent pacemaker (survival at 5 and 10 years: 52.7% and 10.9% in the pacemaker group and 53.8% and 15.3% in the group without a pacemaker, respectively).

CONCLUSION

In conclusion, pacemaker post-TAVR is related to multifactorial risk. Nwaedozie *et al*[9] brought to the body of evidence ECG and clinical findings, but procedural characteristics have at least as much impact on the final need for a permanent pacemaker and potentially on the pacing rate. Long-term follow-up and understanding of the impact of long-term stimulation is of utmost importance.

FOOTNOTES

Co-corresponding authors: Stephane Noble and Karim Bendjelid.

Author contributions: Noble S wrote the first draft; and both authors interpreted the data, critically revised the manuscript for important intellectual content, gave approval for the final version to be published and affirmed that the manuscript is an honest, accurate, and transparent account of the study being reported.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Switzerland

ORCID number: Stephane Noble 0000-0002-8018-2857; Karim Bendjelid 0000-0002-7921-7207.

Corresponding Author's Membership in Professional Societies: American Society of Physiology; European Society of Cardiology.

S-Editor: Gong ZM

L-Editor: A

P-Editor: Zhao S

REFERENCES

- 1 Perrin N, Frei A, Noble S. Transcatheter aortic valve implantation: Update in 2018. *Eur J Intern Med* 2018; **55**: 12-19 [PMID: 30180946 DOI: 10.1016/j.ejim.2018.07.002]
- 2 Frei A, Adamopoulos D, Müller H, Walder B, Perrin N, Reynaud T, Ho S, Roffi M, Mach F, Licker MJ, Noble S. Determinants of hospital length of stay after transcatheter aortic valve implantation with self-expanding prostheses: a prospective, single centre observational study. *Swiss Med Wkly* 2019; **149**: w20095 [PMID: 31256415 DOI: 10.4414/smww.2019.20095]
- 3 Mauler-Wittwer S, Noble S. Volume–Outcome Relationship in Surgical and Cardiac Transcatheter Interventions with a Focus on Transcatheter Aortic Valve Implantation. *J Clin Med* 2022; **11** [PMID: 35807093 DOI: 10.3390/jcm11133806]
- 4 Piazza N, Onuma Y, Jesserun E, Kint PP, Maugenest AM, Anderson RH, de Jaegere PP, Serruys PW. Early and persistent intraventricular conduction abnormalities and requirements for pacemaking after percutaneous replacement of the aortic valve. *JACC Cardiovasc Interv* 2008; **1**: 310-316 [PMID: 19463319 DOI: 10.1016/j.jcin.2008.04.007]
- 5 Muthu Krishnan A, Kadavath S, Kowlgi G, Goel A, Wu F, Jha A, Correa De Sa D, Lahoud R. Permanent Pacemaker Use in Transcatheter Aortic Valve Replacement: Real World Experience from the National Inpatient Sample. *Int J Clin Res* 2023; **3**: 263-275 [DOI: 10.38179/ijcr.v3i1.219]
- 6 Barbanti M, van Mourik MS, Spence MS, Iacovelli F, Martinelli GL, Muir DF, Saia F, Bortone AS, Densem CG, van der Kley F, Bramlage P, Vis M, Tamburino C. Optimising patient discharge management after transfemoral transcatheter aortic valve implantation: the multicentre European FAST-TAVI trial. *EuroIntervention* 2019; **15**: 147-154 [PMID: 30777842 DOI: 10.4244/EIJ-D-18-01197]
- 7 Toggweiler S, Stortecky S, Holy E, Zuk K, Cuculi F, Nietlispach F, Sabti Z, Suci R, Maier W, Jamshidi P, Maisano F, Windecker S, Kobza R, Wenaweser P, Lüscher TF, Binder RK. The Electrocardiogram After Transcatheter Aortic Valve Replacement Determines the Risk for Post-Procedural High-Degree AV Block and the Need for Telemetry Monitoring. *JACC Cardiovasc Interv* 2016; **9**: 1269-1276 [PMID: 27339844 DOI: 10.1016/j.jcin.2016.03.024]
- 8 Glikson M, Nielsen JC, Kronborg MB, Michowitz Y, Auricchio A, Barbash IM, Barrabés JA, Boriani G, Braunschweig F, Brignole M, Burri H, Coats AJS, Deharo JC, Delgado V, Diller GP, Israel CW, Keren A, Knops RE, Kotecha D, Leclercq C, Merkely B, Starck C, Thylén I, Tolosana JM; ESC Scientific Document Group. 2021 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy. *Eur Heart J* 2021; **42**: 3427-3520 [PMID: 34455430 DOI: 10.1093/eurheartj/ehab364]
- 9 Nwaedozie S, Zhang H, Najjar Mojarrab J, Sharma P, Yeung P, Umukoro P, Soodi D, Gabor R, Anderson K, Garcia-Montilla R. Novel predictors of permanent pacemaker implantation following transcatheter aortic valve replacement. *World J Cardiol* 2023; **15**: 582-598 [PMID: 38058399 DOI: 10.4330/wjcv.15.i11.582]
- 10 Popma JJ, Deeb GM, Yakubov SJ, Mumtaz M, Gada H, O'Hair D, Bajwa T, Heiser JC, Merhi W, Kleiman NS, Askew J, Sorajja P, Rovin J, Chetcuti SJ, Adams DH, Teirstein PS, Zorn GL 3rd, Forrest JK, Tchétché D, Resar J, Walton A, Piazza N, Ramlawi B, Robinson N, Petrossian G, Gleason TG, Oh JK, Boulware MJ, Qiao H, Mugglin AS, Reardon MJ; Evolut Low Risk Trial Investigators. Transcatheter Aortic-Valve Replacement with a Self-Expanding Valve in Low-Risk Patients. *N Engl J Med* 2019; **380**: 1706-1715 [PMID: 30883053 DOI: 10.1056/NEJMoa1816885]
- 11 Mack MJ, Leon MB, Thourani VH, Makkar R, Kodali SK, Russo M, Kapadia SR, Malaisrie SC, Cohen DJ, Pibarot P, Leipsic J, Hahn RT, Blanke P, Williams MR, McCabe JM, Brown DL, Babaliaros V, Goldman S, Szeto WY, Genereux P, Pershad A, Pocock SJ, Alu MC, Webb JG, Smith CR; PARTNER 3 Investigators. Transcatheter Aortic-Valve Replacement with a Balloon-Expandable Valve in Low-Risk Patients. *N Engl J Med* 2019; **380**: 1695-1705 [PMID: 30883058 DOI: 10.1056/NEJMoa1814052]
- 12 Tang GHL, Zaid S, Michev I, Ahmad H, Kaple R, Udemir C, Cohen M, Lansman SL. "Cusp-Overlap" View Simplifies Fluoroscopy-Guided Implantation of Self-Expanding Valve in Transcatheter Aortic Valve Replacement. *JACC Cardiovasc Interv* 2018; **11**: 1663-1665 [PMID: 30139479 DOI: 10.1016/j.jcin.2018.03.018]
- 13 Rawish E, Macherey S, Jurczyk D, Pätz T, Jose J, Stiermaier T, Eitel I, Frerker C, Schmidt T. Reduction of permanent pacemaker implantation by using the cusp overlap technique in transcatheter aortic valve replacement: a meta-analysis. *Clin Res Cardiol* 2023; **112**: 633-644 [PMID: 36656375 DOI: 10.1007/s00392-022-02150-8]
- 14 Persia-Paulino YR, Almendarez Lacayo M, Alperi A, Hernández-Vaquero D, Fernández Asensio R, Cuevas Pérez J, Adeva A, Flórez P, Vigil-Escalera M, Álvarez Velasco R, Renilla A, Del Valle Fernández R, Antuña P, Morís de la Tassa C, Avanzas P, Pascual I. Self-expanding TAVI using the cusp overlap technique versus the traditional technique: electrocardiogram changes and 1-year cardiovascular outcomes. *Rev Esp Cardiol (Engl Ed)* 2024; **77**: 29-38 [PMID: 37437882 DOI: 10.1016/j.rec.2023.04.008]
- 15 Grubb K, Yakubov SJ, Nazif TM, Mittal S, Gada H, Fraser DGW, Rovin JD, Khalil R, Pyo RT, Sharma S, Ahmed M, Huang J, Rodes-Cabau J. Management of Postprocedural Conduction Disturbances Using a Prespecified Algorithm in the Optimize PRO Study. *JSCAI* 2023; **3**: 101066 [DOI: 10.1016/j.jscai.2023.101066]
- 16 Gada H, Vora AN, Tang GHL, Mumtaz M, Forrest JK, Laham RJ, Yakubov SJ, Deeb GM, Rammohan C, Huang J, Reardon MJ. Site-Level Variation and Predictors of Permanent Pacemaker Implantation Following TAVR in the Evolut Low-Risk Trial. *Cardiovasc Revasc Med* 2023; **47**: 48-54 [PMID: 36266154 DOI: 10.1016/j.carrev.2022.10.004]
- 17 Bruno F, Munoz Pousa I, Saia F, Vaira MP, Baldi E, Leone PP, Cabanas-Grandio P, Corcione N, Spinoni EG, Annibali G, Russo C, Ziacchi M, Caruzzo CA, Ferlini M, Lanzillo G, De Filippo O, Dusi V, Gallone G, Castagno D, Patti G, La Torre M, Musumeci G, Giordano A, Stefanini G, Salizzoni S, Conrotto F, Rinaldi M, Rordorf R, Abu-Assi E, Raposeiras-Roubin S, Biffi M, D'Ascenzo F, De Ferrari GM. Impact of Right Ventricular Pacing in Patients With TAVR Undergoing Permanent Pacemaker Implantation. *JACC Cardiovasc Interv* 2023; **16**: 1081-1091 [PMID: 37164607 DOI: 10.1016/j.jcin.2023.02.003]
- 18 Farouq L, Chen S, Muntané-Carol G, Regueiro A, Philippon F, Sondergaard L, Jørgensen TH, Lopez-Aguilera J, Kodali S, Leon M, Nazif T, Rodés-Cabau J. Clinical impact of conduction disturbances in transcatheter aortic valve replacement recipients: a systematic review and meta-analysis. *Eur Heart J* 2020; **41**: 2771-2781 [PMID: 31899484 DOI: 10.1093/eurheartj/ehz924]

- 19 **Zito A**, Princi G, Lombardi M, D'Amario D, Vergallo R, Aurigemma C, Romagnoli E, Pelargonio G, Bruno P, Trani C, Burzotta F, Crea F. Long-term clinical impact of permanent pacemaker implantation in patients undergoing transcatheter aortic valve implantation: a systematic review and meta-analysis. *Europace* 2022; **24**: 1127-1136 [PMID: [35138367](#) DOI: [10.1093/europace/euac008](#)]
- 20 **Rück A**, Saleh N, Glaser N. Outcomes Following Permanent Pacemaker Implantation After Transcatheter Aortic Valve Replacement: SWEDEHEART Observational Study. *JACC Cardiovasc Interv* 2021; **14**: 2173-2181 [PMID: [34620397](#) DOI: [10.1016/j.jcin.2021.07.043](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

