

Supplementary Table 1 The clinicopathological characteristics of the Surveillance, Epidemiology, and End Results cohort and the Chinese cohort

Characteristics	SEER Database No. (%)	Chinese Registry No. (%)	<i>P</i> value
Total	32742	3153	
ELN, median [IQR]	20.00[15.00,26.00]	18.00[15.00,22.00]	<0.001
Chemotherapy			<0.001
No	28228(86.2)	1429(45.3)	
Yes	4514(13.8)	1724(54.7)	
Size, median [IQR]	5.00[3.80,7.00]	5.00[4.00,6.50]	0.003
Age			<0.001
>=65	22829(69.7)	1269(40.2)	
20-64	9913(30.3)	1884(59.8)	
Sex			<0.001
Female	17156(52.4)	1345(42.7)	
Male	15586(47.6)	1808(57.3)	
Grade			<0.001
Grade1/2	26221(80.1)	2758(87.5)	
Grade3/4	6521(19.9)	395(12.5)	
Histology			0.507
Adenocarcinoma	27996(85.5)	2683(85.1)	
Mucinous Adenocarcinoma	4222(12.9)	425(13.5)	
Others	524(1.6)	45(1.4)	
Tstage			<0.001
T3	27920(85.3)	2234(70.9)	
T4	4822(14.7)	919(29.1)	