

# World Journal of *Clinical Cases*

*World J Clin Cases* 2021 February 26; 9(6): 1247-1498



**EDITORIAL**

- 1247 Interactive platform for peer review: A proposal to improve the current peer review system  
*Emile SH*

**MINIREVIEWS**

- 1251 Animal models of cathartic colon  
*Meng YY, Li QD, Feng Y, Liu J, Wang EK, Zhong L, Sun QL, Yuan JY*

**ORIGINAL ARTICLE****Case Control Study**

- 1259 New indicators in evaluation of hemolysis, elevated liver enzymes, and low platelet syndrome: A case-control study  
*Kang SY, Wang Y, Zhou LP, Zhang H*

**Retrospective Study**

- 1271 Analysis of hospitalization costs related to fall injuries in elderly patients  
*Su FY, Fu ML, Zhao QH, Huang HH, Luo D, Xiao MZ*

- 1284 Effect of alprostadil in the treatment of intensive care unit patients with acute renal injury  
*Jia Y, Liu LL, Su JL, Meng XH, Wang WX, Tian C*

**Clinical Trials Study**

- 1293 Etomidate *vs* propofol in coronary heart disease patients undergoing major noncardiac surgery: A randomized clinical trial  
*Dai ZL, Cai XT, Gao WL, Lin M, Lin J, Jiang YX, Jiang X*

**Observational Study**

- 1304 Healthy individuals *vs* patients with bipolar or unipolar depression in gray matter volume  
*Zhang YN, Li H, Shen ZW, Xu C, Huang YJ, Wu RH*
- 1318 Impact of metabolism-related mutations on the heart rate of gastric cancer patients after peritoneal lavage  
*Yuan Y, Yao S, Luo GH, Zhang XY*

**CASE REPORT**

- 1329 Efficacy of afatinib in a patient with rare EGFR (G724S/R776H) mutations and amplification in lung adenocarcinoma: A case report  
*He SY, Lin QF, Chen J, Yu GP, Zhang JL, Shen D*

- 1336** Esophageal superficial adenosquamous carcinoma resected by endoscopic submucosal dissection: A rare case report  
*Liu GY, Zhang JX, Rong L, Nian WD, Nian BX, Tian Y*
- 1343** Do medullary thyroid carcinoma patients with high calcitonin require bilateral neck lymph node clearance? A case report  
*Gan FJ, Zhou T, Wu S, Xu MX, Sun SH*
- 1353** Femoral epithelioid hemangioendothelioma detected with magnetic resonance imaging and positron emission tomography/computed tomography: A case report  
*Zhao HG, Zhang KW, Hou S, Dai YY, Xu SB*
- 1359** Noninvasive tools based on immune biomarkers for the diagnosis of central nervous system graft-vs-host disease: Two case reports and a review of the literature  
*Lyu HR, He XY, Hao HJ, Lu WY, Jin X, Zhao YJ, Zhao MF*
- 1367** Periodontally accelerated osteogenic orthodontics with platelet-rich fibrin in an adult patient with periodontal disease: A case report and review of literature  
*Xu M, Sun XY, Xu JG*
- 1379** Subtalar joint pigmented villonodular synovitis misdiagnosed at the first visit: A case report  
*Zhao WQ, Zhao B, Li WS, Assan I*
- 1386** Wilson disease – the impact of hyperimmunity on disease activity: A case report  
*Stremmel W, Longerich T, Liere R, Vacata V, van Helden J, Weiskirchen R*
- 1394** Unexplained elevation of erythrocyte sedimentation rate in a patient recovering from COVID-19: A case report  
*Pu SL, Zhang XY, Liu DS, Ye BN, Li JQ*
- 1402** Thoracic pyogenic infectious spondylitis presented as pneumothorax: A case report  
*Cho MK, Lee BJ, Chang JH, Kim YM*
- 1408** Unilateral pulmonary hemorrhage caused by negative pressure pulmonary edema: A case report  
*Park HJ, Park SH, Woo UT, Cho SY, Jeon WJ, Shin WJ*
- 1416** Osseous Rosai-Dorfman disease of tibia in children: A case report  
*Vithran DTA, Wang JZ, Xiang F, Wen J, Xiao S, Tang WZ, Chen Q*
- 1424** Abdominopelvic leiomyoma with large ascites: A case report and review of the literature  
*Wang YW, Fan Q, Qian ZX, Wang JJ, Li YH, Wang YD*
- 1433** Unusual presentation of granulomatosis with polyangiitis causing periaortitis and consequent subclavian steal syndrome: A case report  
*Cho U, Kim SK, Ko JM, Yoo J*
- 1439** Postoperative discal pseudocyst and its similarities to discal cyst: A case report  
*Fu CF, Tian ZS, Yao LY, Yao JH, Jin YZ, Liu Y, Wang YY*

- 1446** Treatment of oral lichen planus by surgical excision and acellular dermal matrix grafting: Eleven case reports and review of literature  
*Fu ZZ, Chen LQ, Xu YX, Yue J, Ding Q, Xiao WL*
- 1455** Nonalcoholic fatty liver disease as a risk factor for cytomegalovirus hepatitis in an immunocompetent patient: A case report  
*Khiatah B, Nasrollah L, Covington S, Carlson D*
- 1461** Early reoccurrence of traumatic posterior atlantoaxial dislocation without fracture: A case report  
*Sun YH, Wang L, Ren JT, Wang SX, Jiao ZD, Fang J*
- 1469** Intrahepatic cholangiocarcinoma is more complex than we thought: A case report  
*Zeng JT, Zhang JF, Wang Y, Qing Z, Luo ZH, Zhang YL, Zhang Y, Luo XZ*
- 1475** Congenital hepatic fibrosis in a young boy with congenital hypothyroidism: A case report  
*Xiao FF, Wang YZ, Dong F, Li XL, Zhang T*
- 1483** Polidocanol sclerotherapy for multiple gastrointestinal hemangiomas: A case report  
*Yao H, Xie YX, Guo JY, Wu HC, Xie R, Shi GQ*
- 1490** Gastrointestinal stromal tumor with multisegmental spinal metastases as first presentation: A case report and review of the literature  
*Kong Y, Ma XW, Zhang QQ, Zhao Y, Feng HL*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Dr. Quach is an Associate Professor of Gastroenterology at the University of Medicine and Pharmacy at Hochiminh City, Viet Nam, where he received his MD in 1997 and his PhD in 2011. Dr. Quach has published more than 100 reviews and original papers in local and international journals. He has received several awards, including Outstanding Presentation at the Biannual Scientific Congress of Vietnamese Nationwide Medical Schools, Medal of Creativeness from the Vietnamese Central Youth League, etc. Currently, he serves as a Vice President of the Vietnam Association of Gastroenterology and Secretary General of the Vietnam Federation for Digestive Endoscopy. (L-Editor: Filipodia)

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

*WJCC* mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2020 Edition of Journal Citation Reports® cites the 2019 impact factor (IF) for *WJCC* as 1.013; IF without journal self cites: 0.991; Ranking: 120 among 165 journals in medicine, general and internal; and Quartile category: Q3. The *WJCC*'s CiteScore for 2019 is 0.3 and Scopus CiteScore rank 2019: General Medicine is 394/529.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ji-Hong Liu*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lai Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Dennis A Bloomfield, Sandro Vento, Bao-Gan Peng

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

February 26, 2021

**COPYRIGHT**

© 2021 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Retrospective Study

## Effect of alprostadil in the treatment of intensive care unit patients with acute renal injury

Yan Jia, Li-Li Liu, Ji-Liang Su, Xiao-Hua Meng, Wei-Xin Wang, Cui Tian

**ORCID number:** Yan Jia 0000-0001-5306-5842; Li-Li Liu 0000-0001-8080-2772; Ji-Liang Su 0000-0002-0863-5233; Xiao-Hua Meng 0000-0001-6750-0147; Wei-Xin Wang 0000-0002-8451-3123; Cui Tian 0000-0002-6578-8033.

**Author contributions:** Jia Y and Tian C performed the operation; Liu LL and Wang WX designed this retrospective study; Su JL and Meng XH wrote the paper; Tian C was responsible for sorting the data.

**Institutional review board statement:** The study was reviewed and approved by the Research Ethical Committee of Shandong Taishan Hospital Institutional Review Board, No. 2017BL-045-01.

**Informed consent statement:** Informed written consent was obtained from the patient for publication of this report and any accompanying images.

**Conflict-of-interest statement:** The authors declare that they have no conflict of interest.

**Data sharing statement:** No additional data are available.

**Open-Access:** This article is an open-access article that was

Yan Jia, Li-Li Liu, Ji-Liang Su, Xiao-Hua Meng, Department of Critical Care Medicine, Shandong Taishan Hospital, Tai'an 271000, Shandong Province, China

Wei-Xin Wang, Cui Tian, Department of Critical Care Medicine, The Second Affiliated Hospital of Shandong First Medical University, Tai'an 271000, Shandong Province, China

**Corresponding author:** Cui Tian, MHS, Attending Doctor, Department of Critical Care Medicine, The Second Affiliated Hospital of Shandong First Medical University, No. 706 Taishan Street, Taishan District, Tai'an 271000, Shandong Province, China. [15621589099@163.com](mailto:15621589099@163.com)

## Abstract

## BACKGROUND

Acute kidney injury (AKI) is a sudden or rapid decline in the filtration function of the kidneys which is marked by increased serum creatinine or blood urea nitrogen.

## AIM

To examine the value of alprostadil-assisted continuous venous-venous hemofiltration (CVVH) in the treatment of severe AKI in severely ill patients.

## METHODS

This was a retrospective study and the inclusion criteria were as follows: (1) Age of patients ( $\geq 18$  years); (2) Admission to intensive care unit due to non-renal primary disease, APACHE II score ( $\geq 18$  points); (3) The diagnostic criteria of AKI guidelines were formulated with reference to the Global Organization for the Improvement of Prognosis in Kidney Diseases, with AKI grades of II-III; (4) All patients were treated with CVVH; and (5) Complete basic data were obtained for all patients.

## RESULTS

The clinical effect of alprostadil administered in the treatment group was better than that observed in the control group ( $P < 0.05$ ). The urine output of patients in the alprostadil group returned to normal time ( $9.1 \pm 2.0$  d) and was lower than that in the control group ( $10.6 \pm 2.5$  d), the difference was statistically significant ( $P < 0.05$ ); adverse reactions occurred in the alprostadil group compared with the control group, but the difference was not statistically significant ( $P > 0.05$ ).

selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Unsolicited manuscript

**Specialty type:** Medicine, research and experimental

**Country/Territory of origin:** China

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B, B  
Grade C (Good): C  
Grade D (Fair): 0  
Grade E (Poor): 0

**Received:** November 3, 2020

**Peer-review started:** November 3, 2020

**First decision:** November 20, 2020

**Revised:** December 2, 2020

**Accepted:** December 16, 2020

**Article in press:** December 16, 2020

**Published online:** February 26, 2021

**P-Reviewer:** Dediulia T, Lo G, Morgan E

**S-Editor:** Huang P

**L-Editor:** Webster JR

**P-Editor:** Liu JH



## CONCLUSION

Alprostadil-assisted CVVH in the treatment of severely ill patients with AKI can effectively improve the renal resistance index and partial pressure of urine oxygen, and has a positive effect on improving renal function.

**Key Words:** Alprostadil; Continuous venous-venous hemofiltration; Severe; Acute kidney injury

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Alprostadil-assisted continuous venous-venous hemofiltration in the treatment of severely ill patients with acute kidney injury may provide guidance and the basis for clinical practice.

**Citation:** Jia Y, Liu LL, Su JL, Meng XH, Wang WX, Tian C. Effect of alprostadil in the treatment of intensive care unit patients with acute renal injury. *World J Clin Cases* 2021; 9(6): 1284-1292

**URL:** <https://www.wjgnet.com/2307-8960/full/v9/i6/1284.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v9.i6.1284>

## INTRODUCTION

Acute kidney injury (AKI) is a sudden or rapid decline in the filtration function of the kidneys which is marked by increased serum creatinine (Scr) or blood urea nitrogen (BUN). Treatment of severe AKI is usually relatively complicated.

Continuous venous-venous hemofiltration (CVVH) is an important method for the clinical treatment of kidney diseases, which uses convection and diffusion to remove solutes by mimicking the filtration principle of glomeruli. It introduces arterial or venous blood into semi-permeable membrane filters with good permeability, and ultimately removes water and solute. However, kidney function damage is a common complication of CVVH, which increases the cost of clinical treatment and has a serious impact on the quality of life of patients<sup>[1]</sup>. Alprostadil has been proved to have the effects of dilating vascular smooth muscle, anti-platelet agglutination, protecting cells and inhibiting inflammation. It has positive effects on organ protection in critically ill patients, but relatively few clinical studies have reported its use in CVVH to avoid kidney injury<sup>[2]</sup>.

In order to improve the effect of CVVH in patients with AKI and reduce treatment complications, this study was conducted to observe the effect of alprostadil-assisted CVVH in the treatment of severely ill patients with AKI to provide guidance and the basis for clinical practice.

## MATERIALS AND METHODS

### Patients

This retrospective study was conducted to select 100 intensive care unit (ICU) patients with AKI, of whom 50 patients received CVVH and basic treatment measures (control group), and the remaining 50 patients were also given alprostadil simultaneously (alprostadil group). The study subjects were admitted from January 2017 to March 2019, and the inclusion criteria were as follows: (1) Age of patients ( $\geq 18$  years); (2) Admission to ICU due to non-renal primary disease, Acute Physiological Function and Chronic Health Score II (APACHE II) score ( $\geq 18$  points); (3) The diagnostic criteria of AKI guidelines were formulated with reference to the Global Organization for the Improvement of Prognosis in Kidney Diseases, with AKI grades of II-III; (4) All patients were treated with CVVH; and (5) Complete basic data were obtained for all patients. The exclusion criteria were as follows: (1) Patients with primary kidney disease and renal dysfunction; (2) Patients with advanced malignant tumors; (3) Patients after kidney transplantation and liver transplantation; (4) Contraindications to

CVVH treatment; and (5) Patients who were transferred for treatment.

In the alprostadil group, the age of patients ranged from 42 to 79 years, with an average of  $61.7 \pm 8.0$  years, including 29 males and 21 females, all of whom had basic diseases, including 13 patients admitted to hospital due to trauma, 10 patients with cardiovascular diseases, 7 patients with diabetes, and 20 patients with cerebrovascular diseases. The APACHE II score was  $23.6 \pm 2.7$  points, while the AKI grade was grade II in 29 cases and grade III in 21 cases. In the control group, the age of patients ranged from 38 to 79 years, with an average of  $60.4 \pm 10.3$  years, including 32 males and 18 females, who had basic diseases, including 15 patients admitted to hospital due to trauma, 6 patients with cardiovascular diseases, 5 patients with diabetes, and 23 patients with cerebrovascular diseases. The APACHE II score was  $24.0 \pm 2.5$  points, while AKI grade was grade II in 33 cases and grade III in 17 cases. There was no significant difference in age, gender and other baseline data between the two groups ( $P > 0.05$ ).

### Methods

**Treatment methods:** Control group: Jinbao Prismaflex continuous blood purification equipment was used for treatment, and 4000 mL/bag of replacement fluid was selected to filter the basic replacement fluid. The blood flow rate was set to 150 mL/min, and citrate was used for anticoagulation. The initial pump speed was 2.0%-2.5% of the blood flow rate, and 10% calcium gluconate solution was pumped into the venous end of the treatment pipeline to maintain the whole-body free calcium ion level of 1.0-1.2 mmol/L, and fluid resuscitation, anti-infection and glucocorticoid support were given at the same time for symptomatic treatment.

Alprostadil group: In addition to the treatment given to the control group, patients were also treated with alprostadil (produced by Harbin Pharmaceutical Group Bioengineering Co., Ltd., drug approval No: National drug approval No. H20059787, drug batch No. 20170516), and intravenous injection of 10 UG of alprostadil and 10 mL of normal saline was performed, once every 12 h for 1 wk.

Fluid infusion rate was controlled and vital signs of the patients were strictly monitored during treatment in both groups.

**Data collection:** Comparisons between the two groups of patients, included the APACHE II score, Sequential Organ Failure Score (SOFA), Scr, BUN, urinary kidney injury molecule-1 (KIM-1), urinary neutrophil gelatinase-associated lipocalin (NAGL), urinary interleukin-8 (IL-8), renal resistance index, urinary partial pressure of oxygen and therapeutic effect before and after treatment for 7 d [criteria for the improvement of renal function: (1) Cure: After treatment, the patient's urine volume is more than 1000 mL/24 h, Scr is less than 176.8 mmol/L, and BUN is less than 7.14 mmol/L; (2) Improvement: After treatment, the patient's urine volume is more than 1000 mL/24 h, Scr and urea nitrogen levels significantly decreased, but did not reach the normal level (trouble to confirm the normal range of creatinine, should not be 176.8); and (3) Ineffective: Did not meet the above criteria or patient died]. The fasting venous blood of the patients before and after treatment was sampled and centrifuged at a speed of 3000/rpm for 10 min to separate the serum. Then, using the kit provided by BIO SWAMP Company, and operating according to the kit instructions, an enzyme-linked immunosorbent assay was used to determine the concentration changes in urine and KIM-1 in patients.

### Statistical analysis

SPSS 21.0 software was used for statistical analysis, while Scr, BUN, KIM-1 and other measurement indicators in the two groups were expressed by mean  $\pm$  SD. Independent sample *t*-test was used for comparison between the two groups, and  $\chi^2$  or non-parametric test was used for comparison of enumeration data such as gender and clinical efficacy, with  $P < 0.05$  considered statistically significant.

## RESULTS

### Comparison of APACHE II score and SOFA score between the two groups

Before treatment, there was no significant difference in APACHE II score and SOFA score between the alprostadil group and the control group ( $P > 0.05$ ); while after 7 d of treatment, the APACHE II score and SOFA score of the alprostadil group were lower than those of the control group ( $P < 0.05$ ), as shown in [Table 1](#).

Table 1 Baseline data were compared between the two groups

Group	n	Year	Sex		Basic diseases				APACHE II	AKI		
			Men	Women	Trauma	Cardiovascular disease	Cerebrovascular disease	Other		II	III	
Alprostadil group	50	61.7 ± 8.0	29	21	13	10		20	7	23.6 ± 2.7	29	21
Control group	50	60.4 ± 10.3	32	18	15	6		23	5	24.0 ± 2.5	33	17
<i>t/χ<sup>2</sup></i>		0.705	0.378		1.676					-0.769	0.679	
<i>P</i>		0.483	0.539		0.642					0.444	0.410	

APACHE II: Acute Physiological Function and Chronic Health Score II; AKI: Acute renal injury.

### Comparison of renal function index between the two groups

Before treatment, there were no significant differences in the levels of Scr, BUN, KIM-1, NAGL and IL-8 between the alprostadil group and the control group ( $P > 0.05$ ), while after 7 d of treatment, the levels of Scr, BUN, KIM-1, NAGL and IL-8 in the alprostadil group were lower than those in the control group ( $P < 0.05$ ), as shown in Table 2.

### Comparison of renal resistance index and partial pressure of oxygen in urine between the two groups

Before treatment, there was no significant difference in renal resistance index and partial pressure of oxygen in urine between the alprostadil group and control group ( $P > 0.05$ ), while after 7 d of treatment, the renal resistance index in the alprostadil group was lower than that in the control group ( $P < 0.05$ ), and the partial pressure of oxygen in urine was higher than that in the control group ( $P < 0.05$ ), as shown in Table 3, Figures 1 and 2.

### Comparison of the therapeutic effect between the two groups of patients

After treatment, the cure rate, improvement rate and ineffectiveness in the alprostadil group were 64.00%, 28.00% and 8.00%, respectively, while those in the control group were 42.00%, 44.00% and 14.00%, respectively. The clinical effect in the alprostadil group was superior to that in the control group ( $P < 0.05$ ), as shown in Table 4.

### Comparison of urine volume recovery time and adverse reactions between the two groups

The recovery time of urine volume in the alprostadil group was  $9.1 \pm 2.0$  d, which was lower than that in the control group ( $10.6 \pm 2.5$  d), and the difference was statistically significant ( $P < 0.05$ ), while the incidence of adverse reactions in the alprostadil group was not significantly different to that in the control group ( $P > 0.05$ ), as shown in Table 5.

## DISCUSSION

The ICU is a place to rescue severely ill patients, and most patients suffer from multiple organ failure. It has been reported that the mortality rate of patients with more than 2 failed organs is more than 50%, and that of patients with more than 4 failed organs is 100%, which seriously affects the quality of life and safety of patients<sup>[3,4]</sup>. Studies have shown that the release of proinflammatory mediators is increased in severely ill patients, and then tissue replacement downregulates the release of proinflammatory mediators. Generally, the concentration of local inflammatory mediators is higher than the concentration in the circulation and does not lead to destruction of the body's inflammatory response. However, with aggravation of the disease, the local proinflammatory response cannot resist the invasion and will mobilize the systemic defense mechanism, and the excessive and long-lasting release of the proinflammatory response may destroy the homeostasis of the human body and form a systemic inflammatory cascade reaction, eventually causing cell damage, endothelial cell dysfunction, increased microvascular permeability, microcirculation disorder, and changes in the body's coagulation system, which cannot restore the stability of the body's internal environment and have a

**Table 2 Comparison of serum creatinine, blood urea nitrogen and kidney injury molecule-1 levels between the two groups (mean ± SD)**

Group	n	Scr (μmol/L)		BUN (mmol/L)		KIM-1 (pg/mL)	
		Before treatment	7 d after treatment	Before treatment	7 d after treatment	Before treatment	7 d after treatment
Alprostadil group	50	448.1 ± 95.2	180.6 ± 34.7 <sup>ab</sup>	17.8 ± 3.9	8.1 ± 1.8 <sup>ab</sup>	1338.2 ± 202.1	787.4 ± 144.2 <sup>ab</sup>
Control group	50	420.8 ± 81.7	207.1 ± 42.4 <sup>a</sup>	17.3 ± 3.4	9.3 ± 2.0 <sup>a</sup>	1310.3 ± 231.8	855.8 ± 153.9 <sup>a</sup>
T value		1.539	-3.420	0.683	-3.154	0.642	-2.293
P value		0.127	0.001	0.496	0.002	0.523	0.024
Group	n	NAGL (ng/mL)		IL-8 (ng/mL)			
		Before treatment	7 d after treatment	Before treatment	7 d after treatment		
Alprostadil group	50	126.9 ± 16.4	89.5 ± 11.0 <sup>ab</sup>	58.3 ± 9.1	36.8 ± 7.0 <sup>ab</sup>		
Control group	50	130.0 ± 20.1	96.7 ± 12.4 <sup>a</sup>	56.8 ± 8.3	41.0 ± 8.3 <sup>a</sup>		
T value		-0.845	-3.071	0.861	-2.735		
P value		0.400	0.003	0.391	0.007		

<sup>a</sup>P < 0.05, compared with this group before treatment.

<sup>b</sup>P < 0.05, compared with the control group. Scr: Serum creatinine; BUN: Blood urea nitrogen; KIM-1: Kidney injury molecule-1; NAGL: Neutrophil gelatinase-associated lipocalin; IL-8: Interleukin-8.

**Table 3 Comparison of renal function indices between the two groups (mean ± SD)**

Group	n	Renal resistance index		Partial pressure of oxygen in urine (mmHg)	
		Before treatment	7 d after treatment	Before treatment	7 d after treatment
Alprostadil group	50	0.74 ± 0.06	0.65 ± 0.05 <sup>ab</sup>	51.32 ± 10.32	65.06 ± 9.04 <sup>ab</sup>
Control group	50	0.73 ± 0.07	0.68 ± 0.06 <sup>a</sup>	52.40 ± 9.64	59.25 ± 8.75 <sup>a</sup>
T value		0.767	-2.716	-0.541	3.265
P value		0.445	0.008	0.590	0.002

<sup>a</sup>P < 0.05, compared with this group before treatment.

<sup>b</sup>P < 0.05, compared with the control group.

**Table 4 Comparison of clinical effect between two groups, n (%)**



Group	n	Cure	Improve	Ineffective
Alprostadil group	50	32 (64.00)	14 (28.00)	4 (8.00)
Control group	50	21 (42.00)	22 (44.00)	7 (14.00)
Z value		-2.154		
P value		0.031		

serious impact on the safety of patients<sup>[5,6]</sup>. Therefore, in the treatment of severely ill patients, it is necessary not only to provide effective nutritional support to avoid aggravating the burden of organ function, but also to carry out active treatment of anti-inflammatory mediators in the body. Animal experiments have confirmed that the uncontrolled inflammatory response causes the occurrence of MODS; therefore, it is of great significance to improve the prognosis of patients by effectively removing toxins and inflammatory mediators from the body<sup>[7,8]</sup>.

At present, hemodialysis is an important treatment method in severely ill patients, which can remove endotoxin substances, filter excess water, and prolong the survival time of patients. CVVH treatment has become a commonly used method in clinical practice, which uses a variety of physical techniques to displace inflammatory factors

**Table 5 Comparison of urine volume recovery time and adverse reactions between the two groups**

Group	n	Injection spot ache	Abdominal distention	Incidence of adverse reactions (%)
Alprostadil group	50	2	4	6 (12.00)
Control group	50	0	2	2 (4.00)
$\chi^2$ value				2.174
P value				0.14

**Figure 1 Histogram of renal resistance index in the two groups of patients.** Compared with this group before treatment <sup>a</sup> $P < 0.05$ , compared with the control group <sup>b</sup> $P < 0.05$ .**Figure 2 Histogram of partial pressure of oxygen in urine of the two groups of patients.** Compared with this group before treatment <sup>a</sup> $P < 0.05$ , compared with the control group <sup>b</sup> $P < 0.05$ .

from the body, and improve the lung microcirculation and oxygen carrying capacity of parenchymal cells, as well as tissue oxygen utilization, but can also clear inflammatory mediators, improve endothelial cell function in patients, and have a stabilizing effect on disorders of coagulation and fibrinolysis<sup>[9]</sup>. However, it was found that AKI can occur during the course of CVVH treatment, which is due to abnormal renal microcirculation, reduced peripheral vascular resistance, uneven distribution of blood flow in organs, and reduced capillary blood volume. AKI generally occurs mainly in the renal cortex and medulla, and the reduction of renal blood flow causes kidney injury, as well as the swelling of endothelial cells which causes renal vascular occlusion<sup>[10,11]</sup>. On the other hand, due to the expansion of endothelial cells, the reduction of vascular substances and the enhancement of vasoconstrictor substances after injury, the redistribution of renal blood flow leads to the oxidative stress reaction *in vivo*, and renal tubular cell damage can also cause AKI due to the uneven distribution of renal blood flow caused by microcirculation disorder<sup>[12,13]</sup>.

This study combined alprostadil with CVVH in the treatment of patients

complicated by AKI, which is a prostaglandin substance containing high vasoactive substances, and can improve blood viscosity in patients, and at the same time inhibit the function of monocytes, neutrophils and reduce the platelet adhesion rate, so as to achieve anti-inflammatory and other effects<sup>[14,15]</sup>. This study found that after 7 d of treatment, the levels of Scr, BUN and KIM-1 in the alprostadil group were lower than those in the control group, which indicated that the application of alprostadil improved renal function in patients and reduced the degree of renal injury. This is mainly because alprostadil is not easily inactivated as it is wrapped by lipid microspheres, and contains prostatic E1, which can target and be dispersed to injured blood vessels, and has the effects of dilating blood vessels and anti-platelet aggregation. Alprostadil can also lead to increased intracellular levels of cyclic adenosine monophosphate, inhibit thromboxane formation, reduce calcium influx, and play a role in dilating blood vessels<sup>[16,17]</sup>. In addition, this study also found that after 7 d of treatment, the measured values of renal resistance index in the alprostadil group were lower than those in the control group, and the measured values of partial pressure of oxygen in urine were higher than those in the control group, which indicated that the application of alprostadil reduced renal resistance. This is mainly due to the effect of alprostadil on stabilizing lysosomal membrane, weakening vasoconstrictive response, enhancing erythrocyte deformability, stabilizing renal hemodynamics, and enhancing renal glomerular filtration; thus, renal function in patients can be protected by increasing renal blood flow to reduce the pressure in the glomerulus<sup>[18,19]</sup>. Animal experiments have also demonstrated that alprostadil could alleviate pathological damage in renal tissue and promote the regeneration and recovery of renal tubular epithelial cells in rats with acute renal tubular injury, with a good renal protective effect<sup>[20]</sup>.

This study also found that after 7 d of treatment, the APACHE II score and SOFA score in the alprostadil group were lower than those in the control group, indicating that the application of alprostadil in patients undergoing CVVH complicated by AKI can improve the clinical signs and health status of patients. The therapeutic effect and clinical effect in the alprostadil group were better than those in the control group, which indicated that the application of alprostadil in patients undergoing CVVH complicated by AKI could improve the clinical effect. The recovery time of urine volume in the alprostadil group was  $9.1 \pm 2.0$  d, which was short and did not increase adverse reactions, indicating that the application of alprostadil was relatively safe. The advantage of this study was the identification of the therapeutic effect of alprostadil in patients undergoing CVVH complicated with AKI, further confirming its role in protecting renal function in these patients, which is similar to the results of previous studies, and provides a basis for rational clinical treatment. In addition, this study analyzed changes in the renal resistance index and renal blood flow index, which were rarely measured in previous studies. However, due to the short follow-up time and the small number of patients enrolled in this study, it is necessary to expand the sample size and carry out long-term follow-up for further validation of these findings.

## CONCLUSION

In conclusion, alprostadil combined with CVVH in the treatment of severely ill patients complicated by AKI can effectively improve renal resistance index, partial pressure of oxygen in urine, and has a positive effect on improving renal function.

## ARTICLE HIGHLIGHTS

### **Research background**

Acute kidney injury is a serious disease.

### **Research motivation**

To explore new methods for the treatment of acute kidney injury.

### **Research objectives**

To examine the effect of alprostadil on continuous venous-venous hemofiltration (CVVH) in patients with acute renal injury (AKI).

### Research methods

This was a retrospective study of 100 intensive care unit patients with AKI.

### Research results

The clinical effect in the alprostadil group was better than that in the control group ( $P < 0.05$ ); The urine output of patients in the alprostadil group returned to normal ( $9.1 \pm 2.0$  d) quicker than in the control group ( $10.6 \pm 2.5$  d), the difference was statistically significant ( $P < 0.05$ ); adverse reactions occurred in the alprostadil group compared with the control group, but the difference was not statistically significant ( $P > 0.05$ ).

### Research conclusions

Alprostadil combined with CVVH in the treatment of severely ill patients complicated with AKI can effectively improve renal resistance index, partial pressure of oxygen in urine, and has a positive effect on improving renal function.

### Research perspectives

New treatment methods are required for patients with acute kidney injury.

---

## REFERENCES

- 1 **Yang Y**, Meng L, Wu S, Li Y, Zhong Y, Xu F, Zhou XC, Li GQ, Xu GL, Peng KF. LIGHT deficiency aggravates cisplatin-induced acute kidney injury by upregulating mitochondrial apoptosis. *Int Immunopharmacol* 2020; **89**: 106999 [PMID: 33045563 DOI: 10.1016/j.intimp.2020.106999]
- 2 **Manou E**, Thodis E, Arsos G, Pasadakis P, Panagoutsos S, Papadopoulou D, Papagianni A. Fibroblast Growth Factor 23 and  $\alpha$ -Klotho Protein Are Associated with Adverse Clinical Outcomes in Non-Dialysis CKD Patients. *Kidney Blood Press Res* 2020; **45**: 900-915 [PMID: 33040068 DOI: 10.1159/000510351]
- 3 **Cui R**, Li F, Shao J, Wang Y, Yue C, Zheng Y, Li X. Postoperative anemia is a risk factor for acute kidney injury after open aorta and vena cava surgeries. *PLoS One* 2020; **15**: e0240243 [PMID: 33048948 DOI: 10.1371/journal.pone.0240243]
- 4 **Dunaevich A**, Chen H, Musseri D, Kuzi S, Mazaki-Tovi M, Aroch I, Segev G. Acute on chronic kidney disease in dogs: Etiology, clinical and clinicopathologic findings, prognostic markers, and survival. *J Vet Intern Med* 2020; **34**: 2507-2515 [PMID: 33044036 DOI: 10.1111/jvim.15931]
- 5 **Liu D**, Shu G, Jin F, Qi J, Xu X, Du Y, Yu H, Wang J, Sun M, You Y, Zhu M, Chen M, Zhu L, Shen Q, Ying X, Lou X, Jiang S, Du Y. ROS-responsive chitosan-SS31 prodrug for AKI therapy *via* rapid distribution in the kidney and long-term retention in the renal tubule. *Sci Adv* 2020; **6** [PMID: 33036968 DOI: 10.1126/sciadv.abb7422]
- 6 **Sako K**, Furuichi K, Yamamura Y, Oshima M, Toyama T, Kaneko S, Wada T. Association between the recurrence period of acute kidney injury and mortality: a single-centre retrospective observational study in Japan. *BMJ Open* 2019; **9**: e023259 [PMID: 31209081 DOI: 10.1136/bmjopen-2018-023259]
- 7 **Gadalean F**, Simu M, Parv F, Vorovenci R, Tudor R, Schiller A, Timar R, Petrica L, Velciov S, Gluhovschi C, Bob F, Mihaescu A, Timar B, Spasovski G, Ivan V. The impact of acute kidney injury on in-hospital mortality in acute ischemic stroke patients undergoing intravenous thrombolysis. *PLoS One* 2017; **12**: e0185589 [PMID: 29040276 DOI: 10.1371/journal.pone.0185589]
- 8 **Conroy AL**, Hawkes MT, Elphinstone R, Opoka RO, Namasopo S, Miller C, John CC, Kain KC. Chitinase-3-like 1 is a biomarker of acute kidney injury and mortality in paediatric severe malaria. *Malar J* 2018; **17**: 82 [PMID: 29448936 DOI: 10.1186/s12936-018-2225-5]
- 9 **Wen Y**, Parikh CR. The Aftermath of AKI: Recurrent AKI, Acute Kidney Disease, and CKD Progression. *J Am Soc Nephrol* 2021; **32**: 2-4 [PMID: 33037129 DOI: 10.1681/ASN.2020091317]
- 10 **Leballo G**, Chakane PM. Cardiac surgery-associated acute kidney injury: pathophysiology and diagnostic modalities and management. *Cardiovasc J Afr* 2020; **31**: 205-212 [PMID: 32555928 DOI: 10.5830/CVJA-2019-069]
- 11 **Oweis AO**, Alshelleh SA. Incidence and outcomes of acute kidney injury in octogenarians in Jordan. *BMC Res Notes* 2018; **11**: 279 [PMID: 29739428 DOI: 10.1186/s13104-018-3397-3]
- 12 **Nayak SL**, Kumar M, Bihari C, Rastogi A. Bile Cast Nephropathy in Patients with Acute Kidney Injury Due to Hepatorenal Syndrome: A Postmortem Kidney Biopsy Study. *J Clin Transl Hepatol* 2017; **5**: 92-100 [PMID: 28660146 DOI: 10.14218/JCTH.2016.00063]
- 13 **Vanmassenhove J**, Kielstein J, Jörres A, Biesen WV. Management of patients at risk of acute kidney injury. *Lancet* 2017; **389**: 2139-2151 [PMID: 28561005 DOI: 10.1016/S0140-6736(17)31329-6]
- 14 **Sun P**, Xu L, Zhang Q, Li Q. Impact of Toll-like receptor 4 deficiency on cerebrocardiac syndrome. *J Huazhong Univ Sci Technolog Med Sci* 2014; **34**: 161-164 [PMID: 24710925 DOI: 10.1007/s11596-014-1251-y]
- 15 **Bellomo R**, Ronco C, Mehta RL, Asfar P, Boissramé-Helms J, Darmon M, Diehl JL, Duranteau J, Hoste EAJ, Olivier JB, Legrand M, Lerolle N, Malbrain MLNG, Mårtensson J, Oudemans-van Straaten HM, Parienti JJ, Payen D, Perinel S, Peters E, Pickkers P, Rondeau E, Schetz M, Vinsonneau

- C, Wendon J, Zhang L, Laterre PF. Acute kidney injury in the ICU: from injury to recovery: reports from the 5th Paris International Conference. *Ann Intensive Care* 2017; **7**: 49 [PMID: 28474317 DOI: 10.1186/s13613-017-0260-y]
- 16 **Yu F**, Zhu J, Lei M, Wang CJ, Xie K, Xu F, Lin SH. Exploring the metabolic phenotypes associated with different host inflammation of acute respiratory distress syndrome (ARDS) from lung metabolomics in mice. *Rapid Commun Mass Spectrom* 2021; **35**: e8971 [PMID: 33049802 DOI: 10.1002/rcm.8971]
- 17 **Liu G**, Liu F, Xiao L, Kuang Q, He X, Wang Y, Yu Y. Treatment of hyperlipidemic acute pancreatitis with modified Dachengqi decoction combining with conventional therapy based on "six-hollow-organs to be unblocked" theory. *Ann Palliat Med* 2020; **9**: 2045-2053 [PMID: 32692216 DOI: 10.21037/apm-20-1106]
- 18 **Blinder JJ**, Asaro LA, Wypij D, Selewski DT, Agus MSD, Gaies M, Ferguson MA. Acute Kidney Injury After Pediatric Cardiac Surgery: A Secondary Analysis of the Safe Pediatric Euglycemia After Cardiac Surgery Trial. *Pediatr Crit Care Med* 2017; **18**: 638-646 [PMID: 28492399 DOI: 10.1097/PCC.0000000000001185]
- 19 **Mavrakanas TA**, Aurian-Blajeni DE, Charytan DM. Early *versus* late initiation of renal replacement therapy in patients with acute kidney injury: a meta-analysis of randomised clinical trials. *Swiss Med Wkly* 2017; **147**: w14507 [PMID: 29039628 DOI: 10.4414/smw.2017.14507]
- 20 **Chen Z**, Hua S. Transcription factor-mediated signaling pathways' contribution to the pathology of acute lung injury and acute respiratory distress syndrome. *Am J Transl Res* 2020; **12**: 5608-5618 [PMID: 33042442]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

