



全部 图片 视频 新闻 购物 更多 ▾ 搜索工具

找到约 1,910 条结果 (用时 0.65 秒)

Google 学术: A Novel CagA ELISA Exhibits Enhanced Sensitivity of Helicobacter pylori CagA Antibody

Helicobacter pylori infection and gastric lymphoma - Parsonnet - 被引用次数: 1827

... of SHP-2 complexed with Helicobacter pylori CagA - Tsutsumi - 被引用次数: 185

Helicobacter pylori counteracts the apoptotic action of ... - Oldani - 被引用次数: 106

Serum Helicobacter pylori CagA antibody as a biomarker for gastric ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: S Shiota - 2010 - 被引用次数: 33 - 相关文章

Keywords: anti-CagA antibody, east Asia, gastric cancer, Helicobacter pylori ... the east-Asian-type CagA, containing EPIYA-D segments, exhibits a stronger Serum CagA antibodies were assayed by ELISA using purified recombinant CagA protein. With these criteria, the sensitivity was 100% in both countries, but the ...

Serum Helicobacter pylori CagA antibody titer was a useful marker for ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: S Shiota - 2014 - 被引用次数: 7 - 相关文章

Keywords: Helicobacter pylori, CagA, serum antibody, pepsinogen. Go to: ... The level of the serum PG I and PG II were measured by Pepsinogen ELISA kit (Eiken, Co. However, as PG II exhibits a greater raise relative to PG I, the PG I/II ratio ... protein (CRP) using a high-sensitivity assay (hs-CRP) can reveal subclinical ...

Serum antibodies anti-H. pylori and anti-CagA: a comparison between ...

www.ncbi.nlm.nih.gov/pubmed/10414600 ▾ 翻译此页

作者: D Basso - 1999 - 被引用次数: 26 - 相关文章

Serum antibodies anti-H. pylori and anti-CagA: a comparison between four ... RADIM [RADIM I]) in detecting total anti-H. pylori antibodies, and of two further ELISA ... sensitive (92%) than RADIM 1 (79%) in identifying H. pylori positive patients, ...

缺少字词: novel exhibits enhanced

Relationship between ghrelin and Helicobacter pylori infection in ...

onlinepubs.wiley.com/doi/10.1111/j.1365-2026.2006.02949.x/ft1 翻译此页

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 29465

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Novel CagA ELISA exhibits enhanced sensitivity of *Helicobacter pylori* CagA antibody

Matsuo Y *et al.* A Novel ELISA enhances CagA sensitivity

Yuichi Matsuo, Yasutoshi Kido, Junko Akada, Seiji Shiota, Tran Thanh Binh, Tran Thi Huyen Trang, Ho DQ Dung, Pham Huu Tung, Tran Dinh Tri, Ngo P Minh Thuan, Le Quang Tam, Bui Chi Nam, Vu Van Khiem, Yoshio Yamaoka

Yuichi Matsuo, Yasutoshi Kido, Junko Akada, Seiji Shiota, Tran Thanh Binh, Tran Thi Huyen Trang, Yoshio Yamaoka, Department of Environmental and Preventive Medicine, Oita University Faculty of Medicine, Yufu, Oita 879-5593, Japan

Match Overview

1	Internet 177 words crawled on 16-Jun-2016 www.wjgnet.com	3%
2	Internet 64 words crawled on 08-Jun-2015 www.gutpathogens.com	1%
3	Crossref 34 words Honda, Koichi, Masataka Seike, Junya Oribe, Mizuki Endo, Mie Arakawa, Hiroki Syo, Masao Iwao, Masanori Tokoro, Ju	1%
4	Internet 27 words crawled on 27-Apr-2014 www.ncbi.nlm.nih.gov	<1%



学术搜索

找到约 629 条结果 (用时0.06秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

包括专利

包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Focal adhesion kinase is a substrate and downstream effector of SHP-2 complexed with Helicobacter pylori CagA

[HTML] asm.org

..., A Takahashi, T Azuma, H Higashi... - ... and cellular biology, 2006 - Am Soc Microbiol
... CagA proteins isolated from various H. pylori strains exhibit sequence polymorphism, especially in their C ... with a nonradioactive isotope solid-phase enzyme-linked immunosorbent assay (ELISA) kit using ... Tyr-397 and Tyr-576 were reduced upon expression of WT CagA-HA (Fig. ...
被引用次数: 186 相关文章 所有 13 个版本 引用 保存

[HTML] Helicobacter pylori counteracts the apoptotic action of its VacA toxin by injecting the CagA protein into gastric epithelial cells

[HTML] plos.org

A Oldani, M Cormont, V Hofman, V Chiozzi... - PLoS ... , 2009 - journals.plos.org
... CagA association with VacA is thus a novel, highly ingenious microbial strategy to locally protect ... treated or not with VacA, evaluated by using the ratiometric potential-sensitive dye JC ... Infection of gastric epithelial cells with CagA-defective H. pylori confirms that CagA counteracts ...
被引用次数: 108 相关文章 所有 10 个版本 引用 保存 更多

[HTML] Involvement of Helicobacter pylori infection in neuro-hormonal control of food intake

[HTML] jpp.krakow.pl

PC Konturek, M Czesnikiewicz-Guzik... - Journal of physiology ... , 2006 - jpp.krakow.pl
... positive results were obtained for the UBT and the anti-IgG or anti-CagA ELISA tests ... Anti-CagA antibodies occurred in 58% of Hp positive adults and in 52% of Hp positive ... Ghrelin, a novel growth hormone releasing acylated peptide is synthesized in a distinct endocrine cell type ...
被引用次数: 40 相关文章 所有 11 个版本 引用 保存 更多

Helicobacter pylori detection and antimicrobial susceptibility testing

[HTML] asm.org

F Mégraud, P Lehours - Clinical microbiology reviews, 2007 - Am Soc Microbiol
被引用次数: 528 相关文章 所有 6 个版本 引用 保存

Relationship between ghrelin and Helicobacter pylori infection in Polish adult shepherds and their children

[PDF] researchgate.net

M Plonka, PC Konturek, W Bielanski... - Alimentary ... , 2006 - Wiley Online Library
... the results obtained for the UBT and the anti-IgG or anti-CagA ELISA tests were ... 22,23 As suggested previously, 18,23 increased CagA seropositivity in our shepherds could ... Ghrelin, a novel growth hormone releasing acylated peptide is synthesized in a distinct endocrine cell

全部 图片 视频 新闻 购物 更多 ▾ 搜索工具

找到约 1,320 条结果 (用时 0.74 秒)

Google 学术 : Novel CagA ELISA exhibits enhanced sensitivity of Helicobacter pylori CagA antibody

... of SHP-2 complexed with Helicobacter pylori CagA - Tsutsumi - 被引用次数 : 186

Helicobacter pylori counteracts the apoptotic action of ... - Oldani - 被引用次数 : 108

Involvement of Helicobacter pylori infection in neuro- ... - Konturek - 被引用次数 : 40

Serum Helicobacter pylori CagA antibody titer was a useful marker for ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者 : S Shiota - 2014 - 被引用次数 : 9 - 相关文章

Keywords: Helicobacter pylori, CagA, serum antibody, pepsinogen. Go to: ... The level of the serum PG I and PG II were measured by Pepsinogen ELISA kit (Eiken, Co. However, as PG II exhibits a greater raise relative to PG I, the PG I/II ratio ... protein (CRP) using a high-sensitivity assay (hs-CRP) can reveal subclinical ...

Serum Helicobacter pylori CagA antibody as a biomarker for gastric ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : S Shiota - 2010 - 被引用次数 : 33 - 相关文章

Keywords: anti-CagA antibody, east Asia, gastric cancer, Helicobacter pylori ... the east-Asian-type CagA, containing EPIYA-D segments, exhibits a stronger Serum CagA antibodies were assayed by ELISA using purified recombinant CagA protein. With these criteria, the sensitivity was 100% in both countries, but the ...

Serum Helicobacter pylori CagA antibody titer as a useful marker for ...

www.ncbi.nlm.nih.gov/pubmed/24033876 ▾ 翻译此页

作者 : S Shiota - 2014 - 被引用次数 : 9 - 相关文章

Serum Helicobacter pylori CagA antibody titer as a useful marker for advanced ... Serum CagA antibody titer. PG I and PG II were evaluated by ELISA