World Journal of *Radiology*

World J Radiol 2024 October 28; 16(10): 497-628





Published by Baishideng Publishing Group Inc

JR

World Journal of Radiology

Contents

Monthly Volume 16 Number 10 October 28, 2024

REVIEW

- 497 Quantitative magnetic resonance imaging in prostate cancer: A review of current technology Dhiman A, Kumar V, Das CJ
- 512 Yttrium-90 radioembolization treatment strategies for management of hepatocellular carcinoma Hao K, Paik AJ, Han LH, Makary MS

ORIGINAL ARTICLE

Retrospective Study

528 Breast cancer imaging-clinical experience with two-dimensional-shear wave elastography: A retrospective study

Chervenkov L, Georgiev A, Doykov M, Velikova T

CASE REPORT

537 Ectopic recurrence following treatment of arteriovenous malformations in an adult: A case report and review of literature

Cao WY, Li JP, Guo P, Song LX

- Exertional heat stroke with pronounced presentation of microangiopathic hemolytic anemia: A case report 545 Xiang CH, Zhang XM, Liu J, Xiang J, Li L, Song Q
- High complex anal fistula managed by the modified transanal opening of the intersphincteric space via the 552 inter-sphincteric approach: A case report

Wang YQ, Wang Y, Jia XF, Yan QJ, Zheng XP

- 561 Hypoparathyroidism with situs inversus totalis: A case report Yang M, Pu SL, Li L, Ma Y, Qin Q, Wang YX, Huang WL, Hu HY, Zhu MF, Li CZ
- 569 Mesenteric venous thrombosis in a young adult: A case report and review of the literature Yuan JJ, Zhang HF, Zhang J, Li JZ
- 579 Successful management of infection and macrophage activation syndrome patient using low-dose etoposide: A case report Gao SP, Luo XF, Kosari M, Li WJ, Yang L, Tu W, Zhong JX
- 586 Portal venous gas complication following coronary angiography: A case report Yu ZX, Bin Z, Lun ZK, Jiang XJ
- 593 High-resolution magnetic resonance imaging in the diagnosis and management of vertebral artery dissection: A case report

Zhang HB, Duan YH, Zhou M, Liang RC



Contor	World Journal of Radiology							
Conter	Monthly Volume 16 Number 10 October 28, 2024							
600	Epstein-Barr virus positive post-transplant lymphoproliferative disorder with significantly decrease cell chimerism early after transplantation: A case report							
	Guo QN, Liu HS, Li L, Jin DG, Shi JM, Lai XY, Liu LZ, Zhao YM, Yu J, Li YY, Yu FQ, Gao Z, Yan J, Huang H, Luo Y, Ye YS							
608	Asymmetric outcomes in bilateral maxillary impacted tooth extractions: A case report							
	Liu H, Wang F, Tang YL, Yan X							
616	Cryoablation for intrapulmonary bronchial cyst: A case report							
	Li ZH, Ma YY, Niu LZ, Xu KC							
621	Cystic ductal adenocarcinoma of pancreas complicated with neuroendocrine tumor: A case report and review of literature							
	Zou DM, Shu ZY, Cao X							



Contents

Monthly Volume 16 Number 10 October 28, 2024

ABOUT COVER

Editorial Board Member of World Journal of Radiology, Dusan Dj Popovic, MD, PhD, FESBGH, Assistant Professor of Internal Medicine and Gastroenterohepatology, Research Assistant Professor, University of Belgrade, Faculty of Medicine, Department of Gastroenterology and Hepatology, Clinic for Internal Medicine, University Clinical Hospital Center "Dr Dragisa Misovic-Dedinje", Belgrade 11000, Serbia. dusan.popovic@med.bg.ac.rs

AIMS AND SCOPE

The primary aim of World Journal of Radiology (WJR, World J Radiol) is to provide scholars and readers from various fields of radiology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WIR mainly publishes articles reporting research results and findings obtained in the field of radiology and covering a wide range of topics including state of the art information on cardiopulmonary imaging, gastrointestinal imaging, genitourinary imaging, musculoskeletal imaging, neuroradiology/head and neck imaging, nuclear medicine and molecular imaging, pediatric imaging, vascular and interventional radiology, and women's imaging.

INDEXING/ABSTRACTING

The WJR is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for WJR as 1.4; JIF without journal self cites: 1.4; 5-year JIF: 1.8; JIF Rank: 132/204 in radiology, nuclear medicine and medical imaging; JIF Quartile: Q3; and 5-year JIF Quartile: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Wen-Bo Wang; Production Department Director: Xu Guo; Cover Editor: Jia-Ping Yan.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Radiology	https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 1949-8470 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
January 31, 2009	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Monthly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Thomas J Vogl	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/1949-8470/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
October 28, 2024	https://www.wjgnet.com/bpg/Gerlnfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



WJR

World Journal of Radiology

Submit a Manuscript: https://www.f6publishing.com

World J Radiol 2024 October 28; 16(10): 537-544

DOI: 10.4329/wjr.v16.i10.537

ISSN 1949-8470 (online)

CASE REPORT

Ectopic recurrence following treatment of arteriovenous malformations in an adult: A case report and review of literature

Wen-Yu Cao, Jin-Ping Li, Peng Guo, Ling-Xie Song

Specialty type: Radiology, nuclear medicine and medical imaging

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade C Novelty: Grade B Creativity or Innovation: Grade B Scientific Significance: Grade B

P-Reviewer: Goenka S

Received: February 3, 2024 Revised: September 11, 2024 Accepted: September 23, 2024 Published online: October 28, 2024 Processing time: 267 Days and 17.3 Hours



Wen-Yu Cao, Jin-Ping Li, Peng Guo, Department of Neurosurgery, Beijing Chao-Yang Hospital, Capital Medical University, Beijing 100020, China

Ling-Xie Song, Department of Pathology, Beijing Chao-Yang Hospital, Capital Medical University, Beijing 100020, China

Corresponding author: Jin-Ping Li, MD, Chief Physician, Department of Neurosurgery, Beijing Chao-Yang Hospital, Capital Medical University, No. 8 Gongti South Road, Chaoyang District, Beijing 100020, China. ljp6855@aliyun.com

Abstract

BACKGROUND

Digital subtraction angiography (DSA), the gold standard for the diagnosis of intracranial arteriovenous malformations (AVMs), can show clean nidus resection, leading to a perceived cure. Most cases of intracranial AVM recurrence have been reported in pediatric patients. The conventional understanding indicates that AVMs arise when abnormal blood vessels develop between the fourth and eighth weeks of embryonic development, which coincides with the typical period of blood vessel formation in the brain. As such, recurrent ectopic AVM are rare in adults.

CASE SUMMARY

Herein, we present the case of a 31-year-old adult with a history of an intracranial AVM originally diagnosed with a symptomatic de novo cerebellar AVM formation. Recurrence was observed five years following angiographically-confirmed excision of the initial AVM. DSA performed prior to initial AVM resection indicated no cerebellar abnormalities. Moreover, the recurrent arteries exhibited differences in arteries and draining veins. In addition to reporting this case, we analyzed six previously-reported adult patients with similar ectopic recurrent AVMs. These cases are summarized to review and explore the potential causes of ectopic AVM recurrence in adults, which increase the likelihood of acquired AVM.

CONCLUSION

The clinical course of the reported patients demonstrated the possibility of ectopic AVM recurrence in adults. The median time between the diagnosis of the initial AVM and the occurrence of ectopic recurrent AVM in adults was 11 years (range: 5-20 years). Magnetic resonance imaging follow-up for more than 10 years may be required in adult AVM-treated patients.



WJR https://www.wjgnet.com

Key Words: Arteriovenous malformation; Intracranial; Recurrence; De novo; Adults; Ectopic; Case report

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This study highlights an exceptionally rare case of ectopic recurrent intracranial arteriovenous malformation (AVM) in a 31-year-old adult, diagnosed five years following initial AVM resection. this de novo cerebellar AVM demonstrated arterial and venous configurations distinct from typical pediatric cases. Reviewing six similar adult cases, this study explored the potential causes of such rare recurrences. This experience suggests that prolonged magnetic resonance imaging follow-up beyond 10 years may be required to detect new AVMs in previously treated patients.

Citation: Cao WY, Li JP, Guo P, Song LX. Ectopic recurrence following treatment of arteriovenous malformations in an adult: A case report and review of literature. *World J Radiol* 2024; 16(10): 537-544 **URL:** https://www.wjgnet.com/1949-8470/full/v16/i10/537.htm **DOI:** https://dx.doi.org/10.4329/wjr.v16.i10.537

INTRODUCTION

Arteriovenous malformation (AVM) is a deformity defined by the direct communication between arteries and veins, without the presence of a capillary bed[1]. The conventional understanding suggests that AVMs arise when abnormal blood vessels develop between the fourth and eighth weeks of embryonic development, coinciding with the typical period of blood vessel formation in the brain[2]. However, the mechanisms underlying AVM remain unclear. AVM has traditionally been considered a congenital disorder[2–4]; however, numerous case reports have indicated that AVM may also be acquired[5,6].

Most AVMs can be successfully treated with surgical resection, Gamma Knife radiosurgery, or embolization. However, these treatments carry a risk of recurrence. Recurrence is more commonly observed at the original site in pediatric cases, whereas ectopic recurrence in adults is exceedingly rare. Hereon, we present a case of ectopic recurrence of a cranial AVM at a different anatomical site in an adult patient, which is an exceptionally rare form of recurrence.

CASE PRESENTATION

Chief complaints

A 31-year-old man presented with sudden onset vertigo, headache, nausea, vomiting, and urinary and fecal incontinence lasting 6 hours.

History of present illness

On January 31, 2018, the patient presented to our neurology outpatient clinic with acute onset of vertigo, headache, nausea, vomiting, and urinary and fecal incontinence without any apparent precipitating factors. The patient denied experiencing unilateral limb numbness, weakness, or altered mental status.

History of past illness

The patient had a history of seizures, and was initially admitted to the hospital after experiencing a cerebral hemorrhage in 2013, during which time a right frontal lobe AVM was identified through computed tomography (CT) and digital subtraction angiography (DSA) (Figure 1 and Figure 2). AVM resection was performed and confirmed by intraoperative pathology. Angiography performed one year postoperatively (Figure 2) revealed no residual AVM. Additionally, DSA performed prior to the initial AVM resection indicated no cerebellar abnormalities (Figure 3).

Personal and family history

The patient's personal and family history were unremarkable.

Physical examination

A neurological examination was performed on admission, which yielded normal findings.

Laboratory examinations

All laboratory examinations were normal.

Zaishideng® WJR | https://www.wjgnet.com



Figure 1 Preoperative computed tomography showing a lamellar hyperdense area in the right frontal lobe; lamellar hyperdense shadows can be seen in the ventricles, while the right lateral ventricle was narrowed by compression, with a slightly leftward deviation of the midline structure.



Figure 2 Right internal carotid artery angiography. A: Preoperative internal carotid artery angiography lateral cut and; B: An orthogonal slice showing 1 arteriovenous malformation (AVM) in the right frontal lobe, with the supplying artery originating from the anterior cerebral artery; C: Digital subtraction angiography performed one-year postoperatively showing complete AVM resection.

Imaging examinations

CT revealed cerebellar hemorrhage secondary to lateral ventricular hemorrhage (Figure 3), while DSA demonstrated a left cerebellar AVM fed by the left superior cerebellar artery, draining into the transverse sinus *via* two branches of the draining veins (Figure 4).

FINAL DIAGNOSIS

Hypertensive intracerebral hemorrhage (ICH) is a form of acute bleeding within the brain parenchyma or ventricles which develops in patients with a history of hypertension[7]. This condition arises from chronic hypertension, which results in arteriosclerosis and subsequent rupture of the small arteries. The most common sites of hemorrhage are the basal ganglia, pons, and cerebellum[8]. Clinically, it presents with acute onset of severe headache, altered consciousness, and focal neurological deficits. CT typically reveals a hyperdense hemorrhagic lesion, with pronounced surrounding edema. The diagnosis of hypertensive ICH was ruled out given the patient's lack of any history of hypertension.

Cerebral cavernous malformations (CCM) are composed of clusters of abnormal thin-walled blood vessels. Pathologically, it is characterized by multiple lumen formations and vascular leakage at the brain capillary level, leading to disruption of the blood-brain barrier[9]. Clinically, the symptoms are mild and primarily involve recurrent minor hemorrhages. However, in some cases, it can cause severe neurological symptoms, including seizures, focal neurological deficits, and hemorrhagic stroke[10]. magnetic resonance imaging (MRI) findings typically show "popcorn-like" or "honeycomb" lesions, often with evidence of chronic bleeding. CCM was excluded as the patient's MRI did not reveal a "popcorn" appearance, and the hemorrhage was located within the lesion.

WJR https://www.wjgnet.com

Cao WY et al. Ectopic recurrence of arteriovenous malformations



Figure 3 Right vertebral artery angiography and Preoperative computed tomography. A: Right vertebral artery angiography performed before the first arteriovenous malformation resection showing no cerebellar abnormalities; B: Preoperative computed tomography showing multiple patchy hyperdense shadows in the 3rd and 4th ventricles, indicating bilateral cerebellar hemorrhage.



Figure 4 Left vertebral artery angiography. A: Image in the preoperative lateral view showing a cerebellar arteriovenous malformation fed by the left superior cerebellar artery; B: The venous phase of the angiogram showing the draining veins into the transverse sinus; C: Follow-up imaging at 7 months showing no evidence of a residual malformation.

Cerebral developmental venous anomalies (DVAs) are among the most common cerebral vascular anomalies, and are currently viewed as normal variations of the cerebral parenchymal venous angioarchitecture, rather than true malformations[11]. Most DVAs are asymptomatic, although a few patients may present with symptoms related to hemorrhage or local mass effects. On imaging, DSA reveals the characteristic "caput medusae" appearance, in which multiple small veins converge into a single large draining vein[11]. Cerebellar AVM was definitively diagnosed based on the results of DSA. Additionally, the absence of the "caput medusae" sign in the venous phase of the DSA excludes the possibility of a venous malformation.

A final diagnosis of ectopic recurrent intracranial AVM was made.

TREATMENT

Following discussion, the patient underwent a transoccipital submedial craniotomy to address the intracranial hematoma and allow excision of the malformed vascular mass. The hematoma cavity was surgically accessed through the left superior vermis of the cerebellum, allowing removal of 15 mL of the chronic hematoma. Upon exploration, a vascular malformation measuring 2 mm × 1.5 mm × 1.5 mm was identified. Intraoperatively, the cerebellar AVM was fed by the left superior cerebellar artery, which drained into the transverse sinus via two branches of the draining veins. The procedure was completed successfully.

OUTCOME AND FOLLOW-UP

Postoperatively, the patient recovered satisfactorily, with no new neurological dysfunction. Postoperative pathological



Baishidena® WJR | https://www.wjgnet.com

examination confirmed the presence of an AVM (Figure 5), while immunohistochemistry indicated the presence of CD34, and a final diagnosis of ectopic recurrent intracranial AVM was made.

A follow-up examination seven months postoperatively verified complete resection of the lesion (Figure 4), and the patient had recovered completely. The six-year postoperative follow-up confirmed that the patient had fully recovered neurologically, with no signs of recurrence.

DISCUSSION

A search of the PubMed and Web of Science databases covering the period from 2000 to October 2023 was conducted to identify all English-language articles describing adult patients with ectopic recurrent AVM (Table 1). The final search was completed on October 30, 2023. The following subject headings and keywords were used: AVM, recurrence, ectopic, and de novo. The first author assessed 20 potentially eligible articles, of which 14 were excluded, including cases with recurrence occurring in situ, those with no difference in the supply artery or draining vein between the two AVMs, and those in which the patient was underaged at the time of developing recurrent ectopic AVMs.

Our literature search finally revealed six previously-published case reports[6,12–16]; after including the present case report, seven patients were included in the analysis. The patients' demographic characteristics, AVM features, and recurrence times were extracted. Among the seven patients, three were male and four were female. Of the new occurrences, five (71%) were located adjacent to the primary AVM, while two (29%) were located at non-adjacent sites. Additionally, five (71%) patients had the initial AVM detected in adulthood. The median time between the diagnosis of the first AVM and the occurrence of ectopic recurrent AVM in adults was 11 years (range, 5–20 years).

Studies have shown that AVM recurrence predominantly occurs in pediatric patients. Specifically, 82% of AVM recurrences have been reported in this age group[17]. Analysis has further revealed that the average age at initial AVM diagnosis is 13.8 years, with a mean recurrence interval of 4.2 years[17]. Additionally, research has shown that the risk of AVM recurrence decreases by 14% for every annual increase in age at initial surgical resection[18]. These data indicate that the incidence of AVM recurrence is substantially higher among children than adults.

This case study and literature review identified seven adult patients with ectopic recurrent AVM, all of whom were supported by clinical evidence. Although intracranial AVM are relatively rare, they often lead to hemorrhagic stroke in children and young adults^[4]. However, the pathogenesis of this malformation remains unclear^[2]. AVM recurrence is more common in pediatric patients, and some authors have suggested that abnormal vascular growth may continue into childhood, resulting in the eventual development of an AVM^[6,16,19]. Studies have further indicated that genetic and hereditary factors associated with cerebrovascular malformations may play a role in angiogenesis^[3,4,20]. For example, the association between AVM and systemic genetic disorders, such as Osler-Weber-Rendu disease, Sturge-Weber disease, and Wyburn-Mason syndrome, has been reported^[20,21].

This case report describes the case of a man who developed ectopic recurrence at a different anatomical site following initially AVM resection in adulthood, a rarely reported scenario. Six similar cases were further analyzed. Some authors have suggested that postoperative complications such as hematoma, edema, or thrombus in the residual AVM may result in imaging opacities, creating a compartment adjacent to the primary lesion[13,15,16,22,23]. However, this hypothesis is less applicable in cases in which imaging does not reveal any abnormalities following the initial surgery[24-25].

Moreover, in four of the seven analyzed cases, the new AVM was located close to the primary site, with differences in arteries or draining veins, challenging the idea of a hardened compartment[15]. In the present case, the recurrent AVM did not develop adjacent to the initial AVM, thereby reducing the applicability of such an inference. Harris *et al*[26] demonstrated the role of environmental factors and angiogenic processes in AVM formation, supporting the hypothesis that abnormal genes or vasculature coupled with environmental factors activate dormant angiogenesis and form a new AVM mass at a different site[3,24-26].

Some researchers have suggested that the existence of a single genetic abnormality or vascular abnormality increases the risk of AVMs[4]. The presence of external stimuli may further induce neovascularization in a phenomenon known as the "second strike" theory [2,3,20,27]. These external stimuli may include systemic diseases, seizures, radiation exposure, and surgical procedures [2,5,6,14,24,25], and can upregulate environmental proangiogenic factors through multiple pathways, thereby activating otherwise quiescent angiogenic mechanisms [5,14,25].

At the genetic level, research has shown that both germline and somatic mutations can play key roles in the development of AVMs. Indeed, a 2018 study by Nikolaev *et al*[28] revealed that AVM development may be closely associated with KRAS-induced activation of the MAPK-ERK signaling pathway in brain endothelial cells[28,29]. However, a recent study noted that somatic mutations in KRAS are not detected in all patients, and that a single mutation may not be sufficient to trigger AVM, which could require multiple mutations in the same signaling pathway[30]. In addition, signaling pathways such as BMP/TGF- β , PDGF-B/PDGFR- β , and VEGF/VEGFR, as well as microRNA processing, have been indicated to play important roles in the pathogenesis of AVM[29]. Notably, KRAS was the only gene consistently associated with recurrent somatic mutations[30]. Gao *et al*[30] further identified *G12V* and *G12D* mutations in KRAS as common recurrent mutations by identifying 24 relevant somatic mutations in the MAPK pathway [30].

In addition, external stimuli may induce AVM formation by altering the local environment *via* other mechanisms. For example, seizures may upregulate angiogenic factors *via* a hypoxic mechanism, while radiation exposure may promote neoangiogenesis by damaging vascular endothelial cells[3,5]. Related studies have further shown that the expression level of Ki-67 is significantly elevated in pathological sections with recurrence following radiation therapy[13]. These findings indicate that AVM may be characterized by acquired characteristics, and that recurrence may occur as a result of the

Zaishidena® WJR | https://www.wjgnet.com

Table 1 Summary of previously reported cases of ectopic recurrent arteriovenous malformations in adult patients												
	Sex	Initial AVM			Recurrent AVM in adult							
Ref.		Age	Location	Feeding vessels	Age	Location	Feeding vessels	Duration				
Nagm et al[15], 2015	Female	5	Right frontal	MCA, ACA	24	Cortex of the right frontal	MCA, ACA (different branch)	19				
Akimoto <i>et al</i> [12] , 2003	Female	10	Splenium, left occipital		27	Cingulate gyrus, corpus callosum	Bilateral pericallosal arteries	17				
MusLuman <i>et al</i> [<mark>16</mark>], 2010	Female	30	Left occipital	MCA	35	Left parietal-occipital	MCA, PCA	5				
Kawashima et al[13], 2019	Female	8	Cerebellum	SCA, PICA, PCA	28	Cerebellum	SCA, PCA	20				
Shi <i>et al</i> [6], 2018	Male	61	Left temporal		72	Right temporal-occipital	MCA	11				
Torres-Quinones <i>et al</i> [14], 2019	Male	70	Cerebellum	SCA	80	Left posterior thalamus	SCA, PCA	10				
Present case	Male	26	Right frontal	ACA	31	Cerebellum	SCA	5				

AVM: Arteriovenous malformation; SCA: Superior Cerebellar Artery; PICA: Posterior Inferior Cerebellar Artery; PCA: Posterior Cerebral Artery; MCA: Middle Cerebral Artery; ACA: Anterior Cerebral Artery.



Figure 5 Postoperative pathological examination. Large amounts of vascular tissue with crooked and tortuous lumens can be observed in the resected tissue, consistent with the presentation of arteriovenous malformation.

combined influence of genetic, environmental, and other factors.

The interval between the first AVM treatment and the rediscovery of AVMs in seven patients ranged from 5 to 20 years, with a median of 11 years. Approximately 70% of patients experienced ectopic recurrence more than 10 years after treatment, suggesting a significant potential for ectopic AVM recurrence in adulthood. These results suggest that extended MRI follow-up beyond 10 years may be significant for patients who have undergone complete AVM treatment in both childhood and adulthood.

CONCLUSION

Herein, we report a case of ectopic recurrent intracranial AVM in an adult at a different anatomical site, and summarize similar cases to speculate on its pathogenesis, thereby increasing the likelihood of acquired AVM occurrence. The clinical course of these patients demonstrates the possibility of ectopic intracranial AVM recurrence in adults. MRI follow-up for more than 10 years in adult AVM-treated patients may be significant.

Baisbidena® WJR | https://www.wjgnet.com

FOOTNOTES

Author contributions: Cao WY is the first author who performed the literature search and wrote the paper; Li JP provided the medical records; Guo P revised the paper; and Song LX provided the pathological data. All the authors reviewed the results and approved the final version of the manuscript.

Supported by Beijing Municipal Science and Technology Project, No. Z201100005520095.

Informed consent statement: The informed consent form has been signed.

Conflict-of-interest statement: All the authors declare that they have no competing interests.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country of origin: China

ORCID number: Jin-Ping Li 0009-0003-5926-2929; Peng Guo 0009-0000-7250-4855.

S-Editor: Liu H L-Editor: A P-Editor: Zhao YQ

REFERENCES

- 1 Boone CE, Caplan JM, Yang W, Ye X, Colby GP, Coon AL, Tamargo RJ, Huang J. Hemorrhage risk and clinical features of multiple intracranial arteriovenous malformations. J Clin Neurosci 2016; 23: 51-57 [PMID: 26461910 DOI: 10.1016/j.jocn.2015.07.008]
- 2 Nagai Y, Anan M, Fujiki M. Cerebral Arteriovenous Malformations as Acquired Lesions: Case Reports and Review of the Literature. J Stroke Cerebrovasc Dis 2020; 29: 105157 [PMID: 32912502 DOI: 10.1016/j.jstrokecerebrovasdis.2020.105157]
- Rodrigues de Oliveira LF, Castro-Afonso LH, Freitas RK, Colli BO, Abud DG. De Novo Intracranial Arteriovenous Malformation-Case 3 Report and Literature Review. World Neurosurg 2020; 138: 349-351 [PMID: 32229303 DOI: 10.1016/j.wneu.2020.03.109]
- Tasiou A, Tzerefos C, Alleyne CH Jr, Boccardi E, Karlsson B, Kitchen N, Spetzler RF, Tolias CM, Fountas KN. Arteriovenous 4 Malformations: Congenital or Acquired Lesions? World Neurosurg 2020; 134: e799-e807 [PMID: 31715403 DOI: 10.1016/j.wneu.2019.11.001]
- Dogan SN, Bagcilar O, Mammadov T, Kizilkilic O, Islak C, Kocer N. De Novo Development of a Cerebral Arteriovenous Malformation: Case 5 Report and Review of the Literature. World Neurosurg 2019; 126: 257-260 [PMID: 30885864 DOI: 10.1016/j.wneu.2019.02.226]
- Shi S, Gupta R, Moore JM, Griessenauer CJ, Adeeb N, Motiei-Langroudi R, Thomas AJ, Ogilvy CS. De novo AVM formation following 6 venous sinus thrombosis and prior AVM resection in adults: report of 2 cases. J Neurosurg 2018; 128: 506-510 [PMID: 28186446 DOI: 10.3171/2016.9.JNS161710]
- 7 Yu Z, Tao C, Xiao A, Wu C, Fu M, Dong W, Liu M, Yu X, You C. Chinese multidisciplinary guideline for management of hypertensive intracerebral hemorrhage. Chin Med J (Engl) 2022; 135: 2269-2271 [PMID: 36315009 DOI: 10.1097/CM9.00000000001976]
- Sessa M. Intracerebral hemorrhage and hypertension. Neurol Sci 2008; 29 Suppl 2: S258-S259 [PMID: 18690511 DOI: 8 10.1007/s10072-008-0956-8]
- Kim J. Introduction to cerebral cavernous malformation: a brief review. BMB Rep 2016; 49: 255-262 [PMID: 26923303 DOI: 9 10.5483/bmbrep.2016.49.5.036]
- Eren Gozel H, Kök K, Ozlen F, Isler C, Pence S. A novel insight into differential expression profiles of sporadic cerebral cavernous 10 malformation patients with different symptoms. Sci Rep 2021; 11: 19351 [PMID: 34588521 DOI: 10.1038/s41598-021-98647-9]
- San Millán Ruíz D, Gailloud P. Cerebral developmental venous anomalies. Childs Nerv Syst 2010; 26: 1395-1406 [PMID: 20703485 DOI: 10.1007/s00381-010-1253-4]
- 12 Akimoto H, Komatsu K, Kubota Y. Symptomatic de novo arteriovenous malformation appearing 17 years after the resection of two other arteriovenous malformations in childhood: case report. Neurosurgery 2003; 52: 228-31; discussion 231 [PMID: 12493123 DOI: 10.1097/00006123-200301000-00030]
- Kawashima M, Hasegawa H, Kurita H, Suzuki K, Shin M, Ikemura M, Saito N. Ectopic Recurrence of Arteriovenous Malformation After 13 Radiosurgery: Case Report and Insight Regarding Pathogenesis. World Neurosurg 2020; 135: 63-67 [PMID: 31841722 DOI: 10.1016/j.wneu.2019.11.156]
- Torres-Quinones C, Koch MJ, Raymond SB, Patel A. Left Thalamus Arteriovenous Malformation Secondary to Radiation Therapy of 14 Original Vermian Arteriovenous Malformation: Case Report. J Stroke Cerebrovasc Dis 2019; 28: e53-e59 [PMID: 30975463 DOI: 10.1016/j.jstrokecerebrovasdis.2019.03.033]
- 15 Nagm A, Horiuchi T, Ichinose S, Hongo K. Unique double recurrence of cerebral arteriovenous malformation. Acta Neurochir (Wien) 2015; 157: 1461-1466 [PMID: 26153777 DOI: 10.1007/s00701-015-2461-1]
- Musluman AM, Cavusoglu H, Yilmaz A, Aydin Y. Recurrent cerebral arteriovenous malformation with a posterior inferior cerebellar artery 16 aneurysm. Turk Neurosurg 2011; 21: 674-679 [PMID: 22194138 DOI: 10.5137/1019-5149.jtn.3059-10.2]



- Sorenson TJ, Brinjikji W, Bortolotti C, Kaufmann G, Lanzino G. Recurrent Brain Arteriovenous Malformations (AVMs): A Systematic 17 Review. World Neurosurg 2018; 116: e856-e866 [PMID: 29807175 DOI: 10.1016/j.wneu.2018.05.117]
- Copelan A, Drocton G, Caton MT, Smith ER, Cooke DL, Nelson J, Abla AA, Fox C, Amans MR, Dowd CF, Halbach VV, Higashida RT, 18 Lawton MT, Kim H, Fullerton HJ, Gupta N, Hetts SW; UCSF Center For Cerebrovascular Research and UCSF Pediatric Brain Center. Brain Arteriovenous Malformation Recurrence After Apparent Microsurgical Cure: Increased Risk in Children Who Present With Arteriovenous Malformation Rupture. Stroke 2020; 51: 2990-2996 [PMID: 32912090 DOI: 10.1161/STROKEAHA.120.030135]
- Robert T, Blanc R, Botta D, Ciccio G, Smajda S, Redjem H, Fahed R, Piotin M. Management of multiple cerebral arteriovenous 19 malformations in a non-pediatric population. Acta Neurochir (Wien) 2016; 158: 1019-1025 [PMID: 27052512 DOI: 10.1007/s00701-016-2785-5]
- 20 Leblanc GG, Golanov E, Awad IA, Young WL; Biology of Vascular Malformations of the Brain NINDS Workshop Collaborators. Biology of vascular malformations of the brain. Stroke 2009; 40: e694-e702 [PMID: 19834013 DOI: 10.1161/STROKEAHA.109.563692]
- 21 Achrol AS, Guzman R, Varga M, Adler JR, Steinberg GK, Chang SD. Pathogenesis and radiobiology of brain arteriovenous malformations: implications for risk stratification in natural history and posttreatment course. Neurosurg Focus 2009; 26: E9 [PMID: 19409010 DOI: 10.3171/2009.2.FOCUS0926
- Pellettieri L, Svendsen P, Wikholm G, Carlsson CA. Hidden compartments in AVMs--a new concept. Acta Radiol 1997; 38: 2-7 [PMID: 22 9059393 DOI: 10.1080/02841859709171233]
- Utsuki S, Kurata A, Miyasaka Y, Takano M, Ootaka H, Fujii K. Multiple arteriovenous malformations with hemorrhage. Acta Neurochir 23 (Wien) 2002; 144: 97-101 [PMID: 11807652 DOI: 10.1007/s701-002-8279-0]
- Jiang W, Zuo Q, Liu J. Developmental venous anomaly with radiation-induced arteriovenous malformation. Neurol Sci 2022; 43: 6601-6603 24 [PMID: 35790589 DOI: 10.1007/s10072-022-06209-2]
- Lo Presti A, Rogers JM, Assaad NNA, Rodriguez ML, Stoodley MA, Morgan MK. De novo brain arteriovenous malformation after tumor 25 resection: case report and literature review. Acta Neurochir (Wien) 2018; 160: 2191-2197 [PMID: 30206686 DOI: 10.1007/s00701-018-3668-8]
- Harris OA, Chang SD, Harris BT, Adler JR. Acquired cerebral arteriovenous malformation induced by an anaplastic astrocytoma: an 26 interesting case. Neurol Res 2000; 22: 473-477 [PMID: 10935219 DOI: 10.1080/01616412.2000.11740703]
- Florian IA, Beni L, Moisoiu V, Timis TL, Florian IS, Balaşa A, Berindan-Neagoe I. 'De Novo' Brain AVMs-Hypotheses for Development and 27 a Systematic Review of Reported Cases. Medicina (Kaunas) 2021; 57 [PMID: 33652628 DOI: 10.3390/medicina57030201]
- Nikolaev SI, Vetiska S, Bonilla X, Boudreau E, Jauhiainen S, Rezai Jahromi B, Khyzha N, DiStefano PV, Suutarinen S, Kiehl TR, Mendes 28 Pereira V, Herman AM, Krings T, Andrade-Barazarte H, Tung T, Valiante T, Zadeh G, Tymianski M, Rauramaa T, Ylä-Herttuala S, Wythe JD, Antonarakis SE, Frösen J, Fish JE, Radovanovic I. Somatic Activating KRAS Mutations in Arteriovenous Malformations of the Brain. N Engl J Med 2018; 378: 250-261 [PMID: 29298116 DOI: 10.1056/NEJMoa1709449]
- 29 Barbosa Do Prado L, Han C, Oh SP, Su H. Recent Advances in Basic Research for Brain Arteriovenous Malformation. Int J Mol Sci 2019; 20 [PMID: 31731545 DOI: 10.3390/ijms20215324]
- 30 Gao S, Nelson J, Weinsheimer S, Winkler EA, Rutledge C, Abla AA, Gupta N, Shieh JT, Cooke DL, Hetts SW, Tihan T, Hess CP, Ko N, Walcott BP, McCulloch CE, Lawton MT, Su H, Pawlikowska L, Kim H. Somatic mosaicism in the MAPK pathway in sporadic brain arteriovenous malformation and association with phenotype. J Neurosurg 2022; 136: 148-155 [PMID: 34214981 DOI: 10.3171/2020.11.JNS202031]



WJR | https://www.wjgnet.com



Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

