World Journal of *Stem Cells*

World J Stem Cells 2024 August 26; 16(8): 773-826





Published by Baishideng Publishing Group Inc

W J S C World Journal of Stem Cells

Contents

Monthly Volume 16 Number 8 August 26, 2024

EDITORIAL

- Pretreatment can alleviate programmed cell death in mesenchymal stem cells 773 Wan XX, Hu XM, Zhang Q, Xiong K
- 780 Mesenchymal stem cells: A promising therapeutic avenue for non-alcoholic fatty liver disease Cheng CH, Hao WR, Cheng TH

MINIREVIEWS

- 784 Understanding and controlling the variables for advanced cell therapy Jeyaraman N, Shrivastava S, Ravi V, Nallakumarasamy A, Pundkar A, Jeyaraman M
- 799 Bone marrow mesenchymal stem cells in treatment of peripheral nerve injury Zou XF, Zhang BZ, Qian WW, Cheng FM

ORIGINAL ARTICLE

Basic Study

811 Microvesicles derived from mesenchymal stem cells inhibit acute respiratory distress syndrome-related pulmonary fibrosis in mouse partly through hepatocyte growth factor

Chen QH, Zhang Y, Gu X, Yang PL, Yuan J, Yu LN, Chen JM

LETTER TO THE EDITOR

824 Potential plausible role of Wharton's jelly mesenchymal stem cells for diabetic bone regeneration Zheng S, Hu GY, Li JH, Li YK



Contents

Monthly Volume 16 Number 8 August 26, 2024

ABOUT COVER

Editorial Board Member of World Journal of Stem Cells, Mohammed E Grawish, PhD, Professor, Department of Oral Biology, Faculty of Dentistry, Mansoura University, Mansoura 740005, Egypt. Grawish2005@yahoo.com

AIMS AND SCOPE

The primary aim of World Journal of Stem Cells (WJSC, World J Stem Cells) is to provide scholars and readers from various fields of stem cells with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJSC publishes articles reporting research results obtained in the field of stem cell biology and regenerative medicine, related to the wide range of stem cells including embryonic stem cells, germline stem cells, tissue-specific stem cells, adult stem cells, mesenchymal stromal cells, induced pluripotent stem cells, embryonal carcinoma stem cells, hemangioblasts, lymphoid progenitor cells, etc.

INDEXING/ABSTRACTING

The WJSC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, PubMed, PubMed Central, Scopus, Biological Abstracts, BIOSIS Previews, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for WJSC as 3.6; JIF without journal self cites: 3.5; 5-year JIF: 4.2; JIF Rank: 105/205 in cell biology; JIF Quartile: Q3; and 5-year JIF Quartile: Q2. The WJSC's CiteScore for 2023 is 7.8 and Scopus CiteScore rank 2023: Histology is 11/62; Genetics is 78/347; Genetics (clinical) is 19/99; Molecular Biology is 131/410; Cell Biology is 104/285.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Xiang-Di Zhang.; Production Department Director: Xu Guo; Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Stem Cells	https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 1948-0210 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
December 31, 2009	https://www.wignet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Monthly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Shengwen Calvin Li, Carlo Ventura	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/1948-0210/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
August 26, 2024	https://www.wignet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



W J S C World Journal of Stem Cells

Submit a Manuscript: https://www.f6publishing.com

World J Stem Cells 2024 August 26; 16(8): 811-823

DOI: 10.4252/wjsc.v16.i8.811

Basic Study

ISSN 1948-0210 (online)

ORIGINAL ARTICLE

Microvesicles derived from mesenchymal stem cells inhibit acute respiratory distress syndrome-related pulmonary fibrosis in mouse partly through hepatocyte growth factor

Qi-Hong Chen, Ying Zhang, Xue Gu, Peng-Lei Yang, Jun Yuan, Li-Na Yu, Jian-Mei Chen

Specialty type: Cell and tissue engineering Provenance and peer review: Unsolicited article; Externally peer reviewed

Peer-review model: Single blind

Peer-review report's classification Scientific Quality: Grade A, Grade B, Grade C, Grade C, Grade E Novelty: Grade A, Grade B, Grade B, Grade C Creativity or Innovation: Grade A,

Grade B, Grade B, Grade D Scientific Significance: Grade A, Grade A, Grade C, Grade C

P-Reviewer: Brody AR; Cui W; Li SC; Ricci AD

Received: March 19, 2024 Revised: July 4, 2024 Accepted: August 15, 2024 Published online: August 26, 2024 Processing time: 159 Days and 21.3 Hours



Qi-Hong Chen, Department of Critical Care Medicine, Jiangdu People's Hospital of Yangzhou, Jiangdu People's Hospital Affiliated to Yangzhou University, Yangzhou 225200, Jiangsu Province, China

Ying Zhang, Xue Gu, Peng-Lei Yang, Jun Yuan, Li-Na Yu, Department of Critical Care Medicine, Jiangdu People's Hospital of Yangzhou, Yangzhou 225200, Jiangsu Province, China

Jian-Mei Chen, Institute of Translational Medicine, Medical College, Yangzhou University, Yangzhou 225000, Jiangsu Province, China

Corresponding author: Jian-Mei Chen, PhD, Researcher, Institute of Translational Medicine, Medical College, Yangzhou University, No. 136 Jiangyang Middle Road, Hanjiang District, Yangzhou 225000, Jiangsu Province, China. 18051063567@163.com

Abstract

BACKGROUND

Pulmonary fibrosis is one of the main reasons for the high mortality rate among acute respiratory distress syndrome (ARDS) patients. Mesenchymal stromal cellderived microvesicles (MSC-MVs) have been shown to exert antifibrotic effects in lung diseases.

AIM

To investigate the effects and mechanisms of MSC-MVs on pulmonary fibrosis in ARDS mouse models.

METHODS

MSC-MVs with low hepatocyte growth factor (HGF) expression (siHGF-MSC-MVs) were obtained via lentivirus transfection and used to establish the ARDS pulmonary fibrosis mouse model. Following intubation, respiratory mechanicsrelated indicators were measured via an experimental small animal lung function tester. Homing of MSC-MVs in lung tissues was investigated by near-infrared live imaging. Immunohistochemical, western blotting, ELISA and other methods were used to detect expression of pulmonary fibrosis-related proteins and to compare effects on pulmonary fibrosis and fibrosis-related indicators.

RESULTS

The MSC-MVs gradually migrated and homed to damaged lung tissues in the ARDS model mice. Treatment with MSC-MVs significantly reduced lung injury and pulmonary fibrosis scores. However, low expression of HGF (siHGF-MSC-MVs) significantly inhibited the effects of MSC-MVs (P < 0.05). Compared with the ARDS pulmonary fibrosis group, the MSC-MVs group exhibited suppressed expression of type I collagen antigen, type III collagen antigen, and the proteins transforming growth factor- β and α -smooth muscle actin, whereas the siHGF-MVs group exhibited significantly increased expression of these proteins. In addition, pulmonary compliance and the pressure of oxygen/oxygen inhalation ratio were significantly lower in the MSC-MVs group, and the effects of the MSC-MVs were significantly inhibited by low HGF expression (all P < 0.05).

CONCLUSION

MSC-MVs improved lung ventilation functions and inhibited pulmonary fibrosis in ARDS mice partly *via* HGF mRNA transfer.

Key Words: Microvesicles derived from mesenchymal stem cells; Acute respiratory distress syndrome; Pulmonary fibrosis; Hepatocyte growth factor; Mesenchymal stromal cells

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Pulmonary fibrosis is strongly associated with poor outcomes in acute respiratory distress syndrome (ARDS) patients. Currently, there are no effective measures for treating pulmonary fibrosis. Microvesicles derived from mesenchymal stromal cells (MSC-MVs) have anti-pulmonary fibrosis effects; however, their specific effects and mechanisms in ARDS-related pulmonary fibrosis have not been fully established. This study revealed that MSC-MVs inhibited ARDS-related pulmonary fibrosis and fibrosis-related factors partly through hepatocyte growth factor, improved lung compliance and ventilation functions, and increased oxygenation indices. This study provides a new direction for the treatment of ARDS-related pulmonary fibrosis.

Citation: Chen QH, Zhang Y, Gu X, Yang PL, Yuan J, Yu LN, Chen JM. Microvesicles derived from mesenchymal stem cells inhibit acute respiratory distress syndrome-related pulmonary fibrosis in mouse partly through hepatocyte growth factor. *World J Stem Cells* 2024; 16(8): 811-823

URL: https://www.wjgnet.com/1948-0210/full/v16/i8/811.htm DOI: https://dx.doi.org/10.4252/wjsc.v16.i8.811

INTRODUCTION

Acute respiratory distress syndrome (ARDS) is an important cause of respiratory failure in critically ill patients, with a mortality rate of 30%-40% [1,2]. The course of pulmonary fibrosis is also associated with the occurrence of ARDS, exhibiting lung injury, inflammation, and fibrosis. Pulmonary fibrosis is one of the main reasons for the high mortality rate among ARDS patients; however, there are no effective treatment options for pulmonary fibrosis[3-5]. There is an urgent need to identify safer and more effective pulmonary fibrosis treatment options to improve the outcomes of patients with ARDS.

Bone marrow mesenchymal stromal cells (MSCs) have antifibrotic effects; however, their clinical applications are limited by tumorigenicity and ethical issues[6-8]. MSC-derived microvesicles (MSC-MVs) are heterogeneous subcellular structures secreted by MSCs that play important roles in tissue and organ damage repair. Research has shown that MSC-MVs partially inhibit the apoptosis of renal tubular epithelial cells and alleviate renal ischemia-reperfusion injury by transferring miR-21[9]. Du *et al*[10] reported that MSC-MVs alleviate renal fibrosis after partial nephrectomy through M₂ macrophage polarization. MSC-MVs not only have strong tissue repair capabilities but also exert antitumor effects[11]. MSC-MVs derived from adipose tissue inhibit ovarian cancer cell proliferation, apoptosis, and necrosis through different signaling pathways[12]. MSC-MVs have been shown to exert antifibrotic effects in idiopathic pulmonary fibrosis mouse models[13]. Moreover, MSC-MVs have been shown to inhibit ARDS-related pulmonary fibrosis[14]; however, the specific effects and mechanisms have yet to be fully established.

Hepatocyte growth factor (HGF) is an important antifibrotic factor[15-17]. In an ARDS animal model, HGF significantly inhibited lung fibrous tissue proliferation in a dose-dependent manner, reduced hydroxyproline (PcNA) levels in lung tissues, and reversed pulmonary fibrosis[18]. Previously, we reported that MSC-MVSs inhibited pulmonary vascular endothelial fibrous tissue proliferation through HGF, thereby maintaining pulmonary vascular endothelial barrier functions[19]. Therefore, HGF may be an important factor in the inhibition of pulmonary fibrosis by MSC-MVs.

In this study, we investigated the effects of MSC-MVs on pulmonary fibrosis and lung functions in an ARDS model mice. Moreover, we investigated the significance of HGF in the MSC-MV-mediated inhibition of ARDS-related pulmonary fibrosis. Our findings elucidate the role and mechanisms of MSC-MVs in ARDS-induced pulmonary fibrosis

Baishideng® WJSC | https://www.wjgnet.com

and provide insight into new treatment strategies.

MATERIALS AND METHODS

Culture of MSCs and gene interference

Mouse bone marrow-derived MSCs were purchased from Guangzhou Saiye Company (Guangzhou, China), cultured in a specialized medium (Guangzhou Saiye Company) and incubated at 37 °C in a 5% CO₂ atmosphere. The cell culture medium was changed every 3 days, and the cells were used in experiments after they had passed through 3-7 generations. On the basis of our previous method^[19], we used lentivirus-mediated gene transfection technology to obtain MSCs with low expression of the HGF gene (siHGF-MSCs).

Isolation and identification of MSC-MVs

In accordance with our previous method^[19], the MSC-MVs were separated from the MSCs by ultracentrifugation. The culture bottles containing 1×10^6 MSCs were incubated overnight with serum-free DMEM, followed by 0.5% fetal bovine serum. MSC-MVs were obtained by centrifugation at 2000 \times g for 20 minutes, followed by centrifugation at 100000 \times g and 4 °C for 1 hour. The pellets were washed with serum-free DMEM and ultracentrifuged to obtain the MSC-MVs, which were stored at -80 °C for further assays. In accordance with Teresa's method[20], flow cytometry was performed to detect the MSC-MVs markers [positive expression of surface molecules (CD29, CD44, α5 integrins, HLA-I antigen) and negative expression of CD34 and CD73 (Figure 1)]. The morphologies of the MSC-MVs were analyzed with electron microscopy and imaging.

Study ethics declarations

Animal care and interventions were performed in accordance with the National Institute of Health Guide for the Care and Use of Experimental Animals, and the animal protocol (YXYLL-2022-103) was approved by the Institutional Animal Care and Use Committees of the Yangzhou University School of Medicine. This study was carried out in compliance with the ARRIVE guidelines.

Animals and groups

Male C57BL/6 mice (aged 6-8 weeks and weighing 18-20 g) were purchased from Changzhou Cavens Experimental Animal Co., Ltd (Changzhou, China). The C57BL/6 mice were subjected to adaptive care for 2 days under controlled relative humidity ($55\% \pm 10\%$) and temperature (25 ± 2 °C) conditions. Great care was taken to minimize their suffering. This protocol was prepared before the study. Briefly, 36 mice were randomly divided into six groups (6 mice in each group), as follows: Normal saline control group; ARDS group; ARDS pulmonary fibrosis group; MSC group; MSC-MVs group; and siHGF-MSC-MVs group. Lipopolysaccharide (LPS) (2 mg/kg) was instilled into the airways of all groups except the control group. The blank control group was injected with the same volume of normal saline. The ARDS mouse fibrosis model was generated via three LPS injections. Briefly, after weighing the 6-8-week-old C57BL/6 mice, they were intraperitoneally injected with 2% pentobarbital sodium (50 mg/kg) for anesthesia and fixed in the supine position; their airways were opened and LPS (1.5 mg/kg) was dropped into the airway. Twenty-four hours later, they were intraperitoneally injected with LPS (3 mg/kg). After 48 hours, the ARDS pulmonary fibrosis models were finally established by injecting 3 mg/kg LPS into the airways of the mice in the ARDS pulmonary fibrosis, MSC, MSC-MVs and siHGF-MSC groups. After 24 hours, MSCs (5 × 10⁵/100 μ L), MSC-MVs (100 μ g/100 μ L), or siHGF-MSC-MVs (100 μ g/100 μ L), respectively, were injected into the tail vein in the MSC group, MSC-MVs group and siHGF-MSC group.

MSC-MVs homing detection

The MSCs and MSC-MVs were dyed with a near-infrared dye (NIR815; Thermo Fisher Scientific, Waltham, MA, United States), as instructed by the manufacturer. After the ARDS pulmonary fibrosis models were established, they were injected with labeled MSCs or MSC-MVs via the tail vein. In vivo imaging was performed at 24 hours and 72 hours using the CRi Maestro Fluorescence Imaging System (Cambridge Research & Instrumentation, Inc, Woburn, MA, United States). The distributions of MSC-MVs and MSCs in the mouse bodies were determined at wavelengths ranging from 640-660 nm.

Assessment of pulmonary fibrosis and lung injury

Lung tissues were paraffin-embedded, sliced, stained with hematoxylin and eosin (referred to as HE), and evaluated for lung injury using the Smith scoring method to assess the pathological damage and repair status. The degree of pulmonary fibrosis was analyzed via Masson-stained sections. After the animals were euthanized, their left lungs were resected, weighed (wet weight), and placed in a 70 °C constant temperature oven for 72 hours to achieve a constant weight. After reweighing (dry weight), net weight of the two lungs was calculated as the wet/dry (W/D) ratio. Each group of mice was administered 20 g/L Evans blue through the tail vein. After the mice were sacrificed, their left atria were sliced open and uniformly injected with 10 mL of normal saline from the right ventricle to flush the lung tissues thoroughly. The lungs were resected, and the surface water of the lungs was removed. The wet weights were measured. The lung tissues were homogenized with 4 mL formamide and the Evans blue was extracted and centrifuged. The absorbance values of the supernatants were read at 620 nm, after which the amounts of extracted dye were determined using a standard curve. The permeability of the alveolar vascular barriers was determined by the amount of extracted dye



WJSC https://www.wjgnet.com



Figure 1 Detection of mesenchymal stromal cell-derived microvesicle markers by flow cytometry and transmission and scanning electron microscopy. A: The expression of surface molecules (CD29, CD44, a5 integrins and the HLA-I antigen) was positive, whereas the expression of CD34 and CD73 was negative; B: Transmission and scanning electron microscopy were performed on purified mesenchymal stromal cell-derived microvesicles to reveal their spheroid morphologies and confirm their sizes.

per unit of tissue.

Respiratory mechanics and functional detection

After the intervention in each group, the mice were intubated via the trachea, and respiratory mechanics-related indicators of the mice in each group were measured via an experimental small animal lung function tester. The total expiratory resistance, lung compliance, forced vital capacity (FVC) and forced expiratory volume at 0.1 seconds (Fev0.1) were measured in each group. After treatment, arterial blood samples were obtained via left ventricular puncture, and the arterial partial pressure of oxygen (PO₂) was measured using an automated blood gas analyzer. The oxygenation indices [PO₂/oxygen inhalation (FiO₂)] were calculated on the basis of the concentration of FiO₂.

Immunohistochemical detection of pulmonary fibrosis-related indicators

After the experiments, the mice were euthanized and their right lower lung tissues were excised and fixed in 10% neutral formaldehyde. The tissues were paraffin-embedded and subjected to immunohistochemistry to assess type I collagen, type III collagen, transforming growth factor- β (TGF- β) and α -smooth muscle actin (α -SMA) protein expression. Under low magnification, the cells were classified as strongly positive (brown), moderately positive (brown yellow), or weakly



Baishidena® WJSC | https://www.wjgnet.com

positive (light yellow) on the basis of the degree of positive staining. The protein expression levels of type I and III collagens and TGF- β and α -SMA were quantified *via* the use of an immunohistochemical average absorbance tester.

Western blot analysis of pulmonary fibrosis-related indicators

The mice were euthanized, and their right lower lungs were excised for protein extraction. The proteins were incubated with antibodies (against HGF, α-SMA, type I collagen, type III collagen, and GADPH) and secondary antibodies. Finally, the target protein bands were visualized with a chemiluminescence imager for quantitative image analysis.

Determination of hydroxyproline and fibrosis-related inflammatory factor levels

Lung tissues were obtained from the middle lobe of the right lung to measure hydroxyproline levels using an alkaline hydrolysis kit. Briefly, the lungs were mixed with 2 mL of buffered saline (PBS) supplemented with 1 mL of 6 mol/L HCl for hydrolysis. Then, 25 µL of normal saline was added to 1 mL of chloramine (1.4%). After 20 minutes at room temperature, 1 mL of Erlich solution was added to the mixture and incubated at 65 °C for 15 minutes. The concentrations of hydroxyproline in the lung tissues were measured via a spectrophotometer. Mouse serum samples were removed from the -80 °C freezer for analysis of serum TGF- β , HGF, type III procollagen N-terminal aminopeptide (PIIIP), interleukin (IL)-1 β and IL-10 expression via ELISA.

Statistical analysis

The data are presented as mean ± standard deviation. Statistical analyses were performed using the SPSS 21.0 software package (IBM Corp., Armonk, NY, United States). For group comparisons, one-way analysis of variance (i.e. ANOVA) followed by Bonferroni's post hoc correction was used. Statistical significance was considered at P < 0.05.

RESULTS

MSC-MVs homed to the lungs of ARDS pulmonary fibrosis model mice

Near-infrared live imaging was performed to detect homing in the ARDS model mice after injection of MSCs or MSC-MVs. MSCs and MSC-MVs gradually migrated and homed to damaged lung tissues after tail vein injection. After 24 hours, the MSCs and MSC-MVs initially migrated into the damaged lung tissues. After 72 hours, the MSC-MVs distinctly homed to the lung tissues, with significantly increased fluorescence intensity (Figure 2). In summary, the MSC-MVs that were injected into the tail veins of ARDS pulmonary fibrosis model mice were mainly associated with injured lung tissues.

Effects of MSC-MVs on lung injury and pulmonary fibrosis in ARDS pulmonary fibrosis model mice

We compared the effects of MSC-MVs on lung injury and fibrosis in ARDS pulmonary fibrosis model mice via HE and Masson staining, respectively. Compared with those in the control group, the ARDS group showed significantly greater lung injury and pulmonary fibrosis scores. After the injection of the MSC-MVs, there were significant reductions in lung injury and pulmonary fibrosis scores. However, suppression of HGF expression markedly inhibited the above effects of MSC-MVs (Figure 3). These findings suggest that MSC-MVs inhibit ARDS-related pulmonary fibrosis partly through HGFs.

Effects of MSC-MVs on pulmonary fibrosis-related indicators in ARDS pulmonary fibrosis model mice

We further performed immunohistochemistry and western blot assays to investigate the effects of MSC-MVs on expression of fibrosis-related protein markers in the ARDS pulmonary fibrosis model mice. Compared with the ARDS pulmonary fibrosis group, the MSC-MVs group exhibited suppressed expression of type I collagen antigen, type III collagen antigen, and the proteins TGF- β and α -SMA, whereas the siHGF-MVs group exhibited significantly elevated expression of those proteins (Figures 4 and 5). Thus, MSC-MVs inhibit the expression of pulmonary fibrosis-related proteins in ARDS pulmonary fibrosis model mice partly through HGFs.

Effects of MSC-MVs on hydroxyproline in ARDS pulmonary fibrosis model mice

Hydroxyproline levels in lung tissues are correlated with pulmonary fibrosis severity. In this study, we found that the hydroxyproline levels in the pulmonary fibrosis group were significantly greater than those in the control group; those in the MSC-MVs treatment group were significantly lower than those in the pulmonary fibrosis group; whereas those in the siHGF-MSC-MVs group were significantly greater than those in the MSC-MVs group. Therefore, MSC-MVs suppressed hydroxyproline levels in the lung tissues of ARDS pulmonary fibrosis model mice partly through HGFs (Figure 6A).

Effects of MSC-MVs on pulmonary vascular permeability and inflammatory indicators in ARDS pulmonary fibrosis model mice

Lung vascular permeability was assessed by measuring W/D ratios and Evans blue leakage. The lung W/D ratio and Evans blue leakage in the ARDS pulmonary fibrosis group were greater than those in the control group and were significantly lower after MSC-MVs injection. However, the above effects of MSC-MVs were significantly inhibited by HGF suppression (Figure 6B and C). These findings indicate that MSC-MVs inhibited pulmonary vascular permeability in ARDS pulmonary fibrosis partly through HGFs. ELISA was subsequently performed to detect the expression of inflam-



WJSC | https://www.wjgnet.com



Figure 2 Microvesicles derived from mesenchymal stromal cells homing detection by the near-infrared dye NIR815. At 24 hours after injection with mesenchymal stromal cell-derived microvesicles (MSC-MVs), the fluorescence appears mainly yellow in lung, and at 72 hours after injection, the fluorescence appears mainly yellow and red with significantly increased fluorescence intensity.

matory factors related to pulmonary fibrosis. Compared with those in the control group, the expression levels of TGF- β , PIIIP, IL-1 β , and IL-10 in the pulmonary fibrosis group were significantly greater, whereas the HGF levels were lower. However, TGF- β , PIIIP, IL-1 β , and IL-10 expression in the MSC-MVs group were significantly lower while HGF levels were higher than those of the pulmonary fibrosis group. The above effects of MSC-MVs were significantly inhibited by the suppression of HGF expression (Figure 6D). Therefore, MSC-MVs inhibited the expression of fibrosis-related inflammatory factors through HGFs.

Effects of MSC-MVs on respiratory mechanics and lung functions in ARDS pulmonary fibrosis model mice

Pulmonary compliance, Fev0.1, FVC and PO_2/FiO_2 in the ARDS pulmonary fibrosis model group were significantly lower than those in the control group, and MSC-MVs significantly improved these indicators after tail vein injection. However, the above effects of MSC-MVs were significantly inhibited by HGF suppression. There were no significant differences in respiratory resistance or the Fev0.1/FVC among the groups (Figure 7). Therefore, MSC-MVs improved lung ventilation functions and oxygenation indices in ARDS pulmonary fibrosis mouse models partly through HGFs.

DISCUSSION

Pulmonary fibrosis is strongly associated with poor prognostic outcomes in ARDS patients[21-23]. Currently, there are no effective measures for treating pulmonary fibrosis[24,25]. MSC-MVs have anti-pulmonary fibrosis effects; however, their specific effects and mechanisms on ARDS-related pulmonary fibrosis have not been fully established[26]. We found that after MSC-MVs were injected into the tail vein of ARDS pulmonary fibrosis model mice, they first migrated and homed to damaged lung tissues, inhibited the expression of ARDS pulmonary fibrosis-related factors and fibrosis-related factors partly through HGFs, improved lung compliance and ventilation functions, and increased oxygenation indices. The findings of this study provide a new direction for the treatment of ARDS-related pulmonary fibrosis.

Pulmonary fibrosis develops in the early stages of ARDS and occurs throughout the entire process of ARDS[27]. We established ARDS model mice by instilling LPS into their airways. On the 7th day of ARDS model establishment, the lung fibrosis score and fibrosis-related protein expression in the mice, including type I collagen antigen, type III collagen antigen, TGF- β and α -SMA, were significantly increased; therefore, significant pulmonary fibrosis occurred on the 7th day of ARDS. Previous studies on pulmonary fibrosis have focused mostly on bleomycin-induced pulmonary fibrosis models [28,29]. On the basis of a previous study[30], we successfully established ARDS pulmonary fibrosis models *via* three LPS injections. After pulmonary fibrosis was induced, pulmonary fibrosis scores and the expression of type I collagen antigen, type III collagen antigen, TGF- β and α -SMA significantly increased, whereas lung compliance Fev0.1, FVC and ventilation functions significantly decreased, which was consistent with the pathological and pathophysiological changes in pulmonary fibrosis.

Zaishideng® WJSC | https://www.wjgnet.com



Figure 3 Immunohistochemical detection of the effects of mesenchymal stromal cell-derived microvesicles on pulmonary and fibrosisrelated indicators in acute respiratory distress syndrome pulmonary fibrosis mouse models. A: Hematoxylin and eosin staining of lung tissues from each group; B: Lung injury scores of pathological sections from each group; C: Masson staining of lung tissues from each group; D: Lung fibrosis scores of pathological sections from each group: $^{a}P < 0.05$, vs the control group; $^{b}P < 0.05$, vs the pulmonary fibrosis group; $^{c}P < 0.05$, vs the mesenchymal stromal cell-derived microvesicles (MSC-MVs) group (n = 6). Control, acute respiratory distress syndrome (ARDS), pulmonary fibrosis, MSC, MSC-MVs and low hepatocyte growth factor (HGF)-MSC-MVs represent the control group, ARDS group, pulmonary fibrosis group, MSC group, MSC group and low HGF-MSC-MVs group, respectively.

MSCs have good anti-pulmonary fibrosis effects[31], but their applications are limited by their tumorigenicity and ethical considerations[32]. MSC-MVs can avoid these challenges and are convenient for storage, which is conducive to large-scale production[33]. MSC-MVs exhibited good anti-pulmonary fibrosis effects in bleomycin-induced pulmonary fibrosis mouse models; however, their specific effects and the mechanisms involved in alleviating ARDS-related pulmonary fibrosis have yet to be established[34,35]. In this study, we found that most of the MSC-MVs and MSCs homed to the injured lung tissues and significantly inhibited the expression of type I collagen antigen, type III collagen antigen, TGF- β and α -SMA in the ARDS pulmonary fibrosis model mice, with reduced pulmonary fibrosis scores, improved lung compliance, Fev0.1 and FVC, and increased oxygenation indices. Like MSCs, MSC-MVs homed toward the site of lung injury, and there were no significant differences in the inhibition of pulmonary fibrosis between the two. Therefore, MSC-MVs can replace MSCs for anti-ARDS pulmonary fibrosis treatment.

HGF is a cell repair factor that promotes epithelial cell proliferation and collagen degradation and inhibits epithelial cell apoptosis. It is one of the few growth factors that can inhibit fibrosis[36-38]. MSC-MVs contain abundant mRNAs, such as *HGF* mRNA[39]; RNA transfer is an important way by which MSC-MVs exert damage-repair effects[40]. Previously, we reported that MSC-MVs alleviate pulmonary vascular endothelial injury and reduce pulmonary vascular endothelial permeability through *HGF* mRNA transfer[19]. In this study, lentivirus transfection was used to suppress *HGF* mRNA expression in MSC-MVs. The ability of MSC-MVs to inhibit ARDS-related pulmonary fibrosis score, affect the expression of type I collagen antigen, type III collagen antigen, TGF- β and α -SMA, and improve lung compliance and lung ventilation functions such as Fev0.1, FVC, and oxygenation indices, which were significantly inhibited after suppression of *HGF* mRNA in MSC-MVs. These findings imply that *HGF* mRNA transfer is one of the main mechanisms through which MSC-MVs exert their anti-ARDS pulmonary fibrosis effects. During ARDS-induced pulmonary fibrosis, the MSC-MVs administered through the tail vein first homed to the injured lung, after which the HGF transcribed from





Jaisbideng® WJSC | https://www.wjgnet.com



Figure 4 Effects of mesenchymal stromal cell-derived microvesicles on acute respiratory distress syndrome-related pulmonary fibrosis and lung injury. A: Immunohistochemical images of pathological sections of lung tissues from each group; B: α -smooth muscle actin (α -SMA) protein expression; C: Type I collagen protein expression; D: Type III collagen protein expression; E: Transforming growth factor- β 1 (TGF- β 1) protein expression. ^a*P* < 0.05, *vs* the control group; ^b*P* < 0.05 *vs* the pulmonary fibrosis group; ^c*P* < 0.05 *vs* the mesenchymal stromal cell-derived microvesicles (MSC-MVs) group (*n* = 6). Control, acute respiratory distress syndrome (ARDS), pulmonary fibrosis, MSC, MSC-MVs and low hepatocyte growth factor (HGF)-MSC-MVs represent the control group, ARDS group, pulmonary fibrosis group, MSC group, MSC-group and low HGF-MSC-MVs group, respectively.



Figure 5 Western blot analysis of the effects of mesenchymal stromal cell-derived microvesicles on pulmonary fibrosis-related proteins in acute respiratory distress syndrome pulmonary fibrosis model mice. A: Western blot bands of fibrosis-related proteins in the lung tissues of each group; B: Relative expression of fibrosis-related proteins. ${}^{a}P < 0.05$, vs the control group; ${}^{b}P < 0.05$, vs the pulmonary fibrosis group; ${}^{c}P < 0.05$, vs the mesenchymal stromal cell-derived microvesicles (MSC-MVs) group (n = 6). Control, acute respiratory distress syndrome (ARDS), pulmonary fibrosis, MSC, MSC-MVs and low hepatocyte growth factor (HGF)-MSC-MVs represent the control group, ARDS group, pulmonary fibrosis group, MSC group, MSC-MVs group and low HGF-MSC-MVs group, respectively. α -SMA: α -smooth muscle actin; COL: Collagen.

Raishideng® WJSC | https://www.wjgnet.com



Figure 6 Effects of mesenchymal stromal cell-derived microvesicles on pulmonary vascular endothelial permeability and related proteins in acute respiratory distress syndrome pulmonary fibrosis mouse models. A: Lung wet/dry weight ratios for each group; B: Evans blue detection of vascular endothelial permeability in each group; C: Hydroxyproline levels in each group; D: Expression of inflammatory factors in each group detected by ELISA. ^a*P* < 0.05, vs the control group; ^b*P* < 0.05, vs the pulmonary fibrosis group; ^c*P* < 0.05, vs the mesenchymal stromal cell-derived microvesicles (MSC-MVs) group (*n* = 6). Control, acute respiratory distress syndrome (ARDS), pulmonary fibrosis, MSC, MSC-MVs and low hepatocyte growth factor (HGF)-MSC-MVs represent the control group, ARDS group, pulmonary fibrosis group, MSC group, MSC-MVs group and low HGF-MSC-MVs group, respectively. IL: Interleukin; PIIIP: Type III procollagen N-terminal aminopeptide; TGF- β 1: Transforming growth factor- β 1.

the mRNA in the MSC-MVs bound to the c-Met receptors in pulmonary vascular epithelial cells, thereby exerting antipulmonary fibrosis effects. Therefore, *HGF* mRNA transfer is one of the mechanisms by which MSC-MVs resist ARDSinduced pulmonary fibrosis.

MSC-MVs have advantages over MSCs, such as easy mass production, easy storage, no immunogenicity, and no ethical issues[41]. Therefore, MSC-MVs have broader therapeutic prospects than do MSCs in the treatment of ARDS-related pulmonary fibrosis. However, the treatment of pulmonary fibrosis caused by ARDS with MSC-MVs is still in the preliminary research stage, and the following problems remain: (1) There is still a lack of standard methods for the extraction, purification, and identification of MSC-MVs; (2) The timing, dosage, and administration methods of MSC-MV treatment need to be clarified; and (3) The mechanism by which MSC-MVs combat pulmonary fibrosis is not very clear. Future research should focus on how to obtain MSC-MVs in a standardized manner, clarify the timing, dosage and administration of MSC-MVs treatment, explore the mechanism of MSC-MVs treatment for pulmonary fibrosis, and carry out further clinical research[42,43].

This study has several limitations. First, researchers have not determined whether the same or better effects can be obtained by administering MSC-MVs intratracheally instead of intravenously. Second, MSC-MVs were administered 24 hours after the establishment of the pulmonary fibrosis model. However, the antifibrotic effects of MSC-MVs administered at different times after pulmonary fibrosis may vary. In addition, finally, only a single dose of MSC-MVs was administered, and two or more doses after the development of pulmonary fibrosis may have improved the anti-fibrosis effects. These limitations should be addressed in future studies.

CONCLUSION

Intravenously administered MSC-MVs homed to the site of lung injury and inhibited ARDS-related pulmonary fibrosis. In addition, *HGF* mRNA transfer is an important mechanism by which MSC-MVs inhibit ARDS-related pulmonary fibrosis. This study provides insight into new treatment approaches for ARDS-related pulmonary fibrosis.

Zaishideng® WJSC | https://www.wjgnet.com



Figure 7 Effects of mesenchymal stromal cell-derived microvesicles on respiratory mechanics and lung functions in acute respiratory distress syndrome pulmonary fibrosis mouse models. A: Pulmonary expiratory resistance; B: Pulmonary compliance; C: Forced expiratory volume in 0.1 seconds (Fev0.1); D: Forced vital capacity (FVC), E: Fev0.1/FVC; F: Pressure of oxygen (PO₂)/oxygen inhalation (FiO₂); G: Transmission and scanning electron microscopy were performed on purified mesenchymal stromal cell-derived microvesicles (MSC-MVs) to reveal their spheroid morphologies and confirm their sizes. ^aP < 0.05, vs the control group; ^bP < 0.05, vs the pulmonary fibrosis group; ^cP < 0.05, vs the MSC-MVs group (n = 6). Control, acute respiratory distress syndrome (ARDS), pulmonary fibrosis, MSC, MSC-MVs and low hepatocyte growth factor (HGF)-MSC-MVs represent the control group, ARDS group, pulmonary fibrosis group, MSC group, MSC-MVs group and low HGF-MSC-MVs group, respectively.

FOOTNOTES

Author contributions: Chen QH designed the study and wrote the manuscript; Zhang Y participated in the cell culture and molecular biology experiments; Gu X performed the animal experiments and data acquisition; Yang PL contributed to the data acquisition and statistical analysis; Yuan J helped with the animal experiments and data acquisition; Yu LN performed the molecular biology experiments and statistical analyses; Chen JM contributed to research quality control and manuscript revision; All authors approved the final manuscript.

Supported by Research Project of Jiangsu Provincial Health Commission, No. Z2022008; and Research Project of Yangzhou Health Commission, No. 2023-2-27.

Institutional animal care and use committee statement: Animal care and interventions were performed in accordance with the National Institute of Health Guide for the Care and Use of Experimental Animals, and the animal protocol (YXYLL-2022-103) was approved by the Institutional Animal Care and Use Committees of the Yangzhou University School of Medicine.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: Technical appendix, statistical code, and dataset available upon reasonable request from the corresponding author at 18051063567@163.com.

ARRIVE guidelines statement: The authors have read the ARRIVE guidelines, and the manuscript was prepared and revised according to the ARRIVE guidelines.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/



WJSC | https://www.wjgnet.com

Country of origin: China

ORCID number: Jian-Mei Chen 0000-0003-0178-4954.

S-Editor: Wang JJ L-Editor: Filipodia P-Editor: Zheng XM

REFERENCES

- 1 Gorman EA, O'Kane CM, McAuley DF. Acute respiratory distress syndrome in adults: diagnosis, outcomes, long-term sequelae, and management. Lancet 2022; 400: 1157-1170 [PMID: 36070788 DOI: 10.1016/S0140-6736(22)01439-8]
- 2 Margaria JP, Moretta L, Alves-Filho JC, Hirsch E. PI3K Signaling in Mechanisms and Treatments of Pulmonary Fibrosis Following Sepsis and Acute Lung Injury. Biomedicines 2022; 10 [PMID: 35453505 DOI: 10.3390/biomedicines10040756]
- Menk M, Estenssoro E, Sahetya SK, Neto AS, Sinha P, Slutsky AS, Summers C, Yoshida T, Bein T, Ferguson ND. Current and evolving 3 standards of care for patients with ARDS. Intensive Care Med 2020; 46: 2157-2167 [PMID: 33156382 DOI: 10.1007/s00134-020-06299-6]
- 4 Liu H, Shi Q, Tang L, Wang H, Wang D. Apelin-13 ameliorates LPS-induced endothelial-to-mesenchymal transition and post-acute lung injury pulmonary fibrosis by suppressing transforming growth factor-β1 signaling. Shock 2023; 59: 108-117 [PMID: 36377383 DOI: 10.1097/SHK.000000000002046
- Raghu G, Remy-Jardin M, Richeldi L, Thomson CC, Inoue Y, Johkoh T, Kreuter M, Lynch DA, Maher TM, Martinez FJ, Molina-Molina M, 5 Myers JL, Nicholson AG, Ryerson CJ, Strek ME, Troy LK, Wijsenbeek M, Mammen MJ, Hossain T, Bissell BD, Herman DD, Hon SM, Kheir F, Khor YH, Macrea M, Antoniou KM, Bouros D, Buendia-Roldan I, Caro F, Crestani B, Ho L, Morisset J, Olson AL, Podolanczuk A, Poletti V, Selman M, Ewing T, Jones S, Knight SL, Ghazipura M, Wilson KC. Idiopathic Pulmonary Fibrosis (an Update) and Progressive Pulmonary Fibrosis in Adults: An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline. Am J Respir Crit Care Med 2022; 205: e18-e47 [PMID: 35486072 DOI: 10.1164/rccm.202202-0399ST]
- Hou L, Zhu Z, Jiang F, Zhao J, Jia Q, Jiang Q, Wang H, Xue W, Wang Y, Tian L. Human umbilical cord mesenchymal stem cell-derived 6 extracellular vesicles alleviated silica induced lung inflammation and fibrosis in mice via circPWWP2A/miR-223-3p/NLRP3 axis. Ecotoxicol Environ Saf 2023; 251: 114537 [PMID: 36646008 DOI: 10.1016/j.ecoenv.2023.114537]
- 7 Li M, Jiang Y, Hou Q, Zhao Y, Zhong L, Fu X. Potential pre-activation strategies for improving therapeutic efficacy of mesenchymal stem cells: current status and future prospects. Stem Cell Res Ther 2022; 13: 146 [PMID: 35379361 DOI: 10.1186/s13287-022-02822-2]
- Guo Z, Zhang Y, Yan F. Potential of Mesenchymal Stem Cell-Based Therapies for Pulmonary Fibrosis. DNA Cell Biol 2022; 41: 951-965 8 [PMID: 36301251 DOI: 10.1089/dna.2022.0327]
- Du T, Zhou J, Chen WX, Zhang XL, Ji TY, Liu J, Rong L, Wang LD, Zhou RJ, Ding DG. Microvesicles derived from human umbilical cord 9 mesenchymal stem cells ameliorate renal ischemia-reperfusion injury via delivery of miR-21. Cell Cycle 2020; 19: 1285-1297 [PMID: 32329653 DOI: 10.1080/15384101.2020.1748940]
- Du T, Ju G, Zhou J, Zhong L, Rong L, Chen W, Zhang X, Zhou R, Ding D, Ji T. Microvesicles derived from human umbilical cord 10 mesenchyme promote M2 macrophage polarization and ameliorate renal fibrosis following partial nephrectomy via hepatocyte growth factor. Hum Cell 2021; 34: 1103-1113 [PMID: 33860459 DOI: 10.1007/s13577-021-00525-z]
- 11 Guven DC, Sahin TK, Erul E, Rizzo A, Ricci AD, Aksoy S, Yalcin S. The association between albumin levels and survival in patients treated with immune checkpoint inhibitors: A systematic review and meta-analysis. Front Mol Biosci 2022; 9: 1039121 [PMID: 36533070 DOI: 10.3389/fmolb.2022.1039121
- Szyposzynska A, Bielawska-Pohl A, Krawczenko A, Doszyn O, Paprocka M, Klimczak A. Suppression of Ovarian Cancer Cell Growth by 12 AT-MSC Microvesicles. Int J Mol Sci 2020; 21 [PMID: 33266317 DOI: 10.3390/ijms21239143]
- 13 Abreu SC, Lopes-Pacheco M, Weiss DJ, Rocco PRM. Mesenchymal Stromal Cell-Derived Extracellular Vesicles in Lung Diseases: Current Status and Perspectives. Front Cell Dev Biol 2021; 9: 600711 [PMID: 33659247 DOI: 10.3389/fcell.2021.600711]
- 14 Zhang X, Ye L, Tang W, Ji Y, Zheng L, Chen Y, Ge O, Huang C. Wnt/β-Catenin Participates in the Repair of Acute Respiratory Distress Syndrome-Associated Early Pulmonary Fibrosis via Mesenchymal Stem Cell Microvesicles. Drug Des Devel Ther 2022; 16: 237-247 [PMID: 35082486 DOI: 10.2147/DDDT.S344309]
- 15 Zhao Y, Ye W, Wang YD, Chen WD. HGF/c-Met: A Key Promoter in Liver Regeneration. Front Pharmacol 2022; 13: 808855 [PMID: 35370682 DOI: 10.3389/fphar.2022.808855]
- Usunier B, Brossard C, L'Homme B, Linard C, Benderitter M, Milliat F, Chapel A. HGF and TSG-6 Released by Mesenchymal Stem Cells 16 Attenuate Colon Radiation-Induced Fibrosis. Int J Mol Sci 2021; 22 [PMID: 33670243 DOI: 10.3390/ijms22041790]
- 17 Yunus J, Salman M, Lintin GBR, Muchtar M, Sari DCR, Arfian N, Romi MM. Chlorogenic acid attenuates kidney fibrosis via antifibrotic action of BMP-7 and HGF. Med J Malaysia 2020; 75: 5-9 [PMID: 32471962]
- Quesnel C, Marchand-Adam S, Fabre A, Marchal-Somme J, Philip I, Lasocki S, Leçon V, Crestani B, Dehoux M. Regulation of hepatocyte 18 growth factor secretion by fibroblasts in patients with acute lung injury. Am J Physiol Lung Cell Mol Physiol 2008; 294: L334-L343 [PMID: 18065658 DOI: 10.1152/ajplung.00096.2007]
- Wang H, Zheng R, Chen Q, Shao J, Yu J, Hu S. Mesenchymal stem cells microvesicles stabilize endothelial barrier function partly mediated 19 by hepatocyte growth factor (HGF). Stem Cell Res Ther 2017; 8: 211 [PMID: 28969681 DOI: 10.1186/s13287-017-0662-7]
- L Ramos T, Sánchez-Abarca LI, Muntión S, Preciado S, Puig N, López-Ruano G, Hernández-Hernández Á, Redondo A, Ortega R, Rodríguez 20 C, Sánchez-Guijo F, del Cañizo C. MSC surface markers (CD44, CD73, and CD90) can identify human MSC-derived extracellular vesicles by conventional flow cytometry. Cell Commun Signal 2016; 14: 2 [PMID: 26754424 DOI: 10.1186/s12964-015-0124-8]
- 21 Bos LDJ, Ware LB. Acute respiratory distress syndrome: causes, pathophysiology, and phenotypes. Lancet 2022; 400: 1145-1156 [PMID: 36070787 DOI: 10.1016/S0140-6736(22)01485-4]
- Sturgill JL, Mayer KP, Kalema AG, Dave K, Mora S, Kalantar A, Carter DJ, Montgomery-Yates AA, Morris PE. Post-intensive care 22



syndrome and pulmonary fibrosis in patients surviving ARDS-pneumonia of COVID-19 and non-COVID-19 etiologies. *Sci Rep* 2023; **13**: 6554 [PMID: 37085548 DOI: 10.1038/s41598-023-32699-x]

- Fang XZ, Li M, Wang YX, Zhang P, Sun MM, Xu JX, Yang YY, He YJ, Yu Y, Li RT, Zhou T, Reng LH, Sun DY, Shu HQ, Yuan SY, Xu JQ, Shang Y. Mechanosensitive ion channel Piezo1 mediates mechanical ventilation-exacerbated ARDS-associated pulmonary fibrosis. *J Adv Res* 2023; 53: 175-186 [PMID: 36526145 DOI: 10.1016/j.jare.2022.12.006]
- Wendisch D, Dietrich O, Mari T, von Stillfried S, Ibarra IL, Mittermaier M, Mache C, Chua RL, Knoll R, Timm S, Brumhard S, Krammer T, Zauber H, Hiller AL, Pascual-Reguant A, Mothes R, Bülow RD, Schulze J, Leipold AM, Djudjaj S, Erhard F, Geffers R, Pott F, Kazmierski J, Radke J, Pergantis P, Baßler K, Conrad C, Aschenbrenner AC, Sawitzki B, Landthaler M, Wyler E, Horst D; Deutsche COVID-19 OMICS Initiative (DeCOI), Hippenstiel S, Hocke A, Heppner FL, Uhrig A, Garcia C, Machleidt F, Herold S, Elezkurtaj S, Thibeault C, Witzenrath M, Cochain C, Suttorp N, Drosten C, Goffinet C, Kurth F, Schultze JL, Radbruch H, Ochs M, Eils R, Müller-Redetzky H, Hauser AE, Luecken MD, Theis FJ, Conrad C, Wolff T, Boor P, Selbach M, Saliba AE, Sander LE. SARS-CoV-2 infection triggers profibrotic macrophage responses and lung fibrosis. *Cell* 2021; **184**: 6243-6261.e27 [PMID: 34914922 DOI: 10.1016/j.cell.2021.11.033]
- Liang Y, Xu Y, Lu B, Huang Y, Xu S, Xie J, Liu M, Che D, Ma L, Tao J, Hong J, Zhang J, Situ X, Ou X, Chen L, Li Y, Zhang L, Wu Z. Inositol Alleviates Pulmonary Fibrosis by Promoting Autophagy *via* Inhibiting the HIF-1α-SLUG Axis in Acute Respiratory Distress Syndrome. *Oxid Med Cell Longev* 2022; 2022: 1030238 [PMID: 36589681 DOI: 10.1155/2022/1030238]
- 26 Sang L, Guo X, Fan H, Shi J, Hou S, Lv Q. Mesenchymal Stem Cell-Derived Extracellular Vesicles as Idiopathic Pulmonary Fibrosis Microenvironment Targeted Delivery. Cells 2022; 11 [PMID: 35954166 DOI: 10.3390/cells11152322]
- 27 Aschner Y, Correll KA, Beke KM, Foster DG, Roybal HM, Nelson MR, Meador CL, Strand M, Anderson KC, Moore CM, Reynolds PR, Kopf KW, Burnham EL, Downey GP. PTPα promotes fibroproliferative responses after acute lung injury. *Am J Physiol Lung Cell Mol Physiol* 2022; **323**: L69-L83 [PMID: 35670474 DOI: 10.1152/ajplung.00436.2021]
- 28 Alvarez-Palomo B, Sanchez-Lopez LI, Moodley Y, Edel MJ, Serrano-Mollar A. Induced pluripotent stem cell-derived lung alveolar epithelial type II cells reduce damage in bleomycin-induced lung fibrosis. *Stem Cell Res Ther* 2020; 11: 213 [PMID: 32493487 DOI: 10.1186/s13287-020-01726-3]
- 29 Periera-Simon S, Xia X, Catanuto P, Coronado R, Kurtzberg J, Bellio M, Lee YS, Khan A, Smith R, Elliot SJ, Glassberg MK. Anti-fibrotic effects of different sources of MSC in bleomycin-induced lung fibrosis in C57BL6 male mice. *Respirology* 2021; 26: 161-170 [PMID: 32851725 DOI: 10.1111/resp.13928]
- 30 Hu X, Xu Q, Wan H, Hu Y, Xing S, Yang H, Gao Y, He Z. PI3K-Akt-mTOR/PFKFB3 pathway mediated lung fibroblast aerobic glycolysis and collagen synthesis in lipopolysaccharide-induced pulmonary fibrosis. *Lab Invest* 2020; 100: 801-811 [PMID: 32051533 DOI: 10.1038/s41374-020-0404-9]
- 31 Escarrer-Garau G, Martín-Medina A, Truyols-Vives J, Gómez-Bellvert C, Elowsson L, Westergren-Thorsson G, Molina-Molina M, Mercader-Barceló J, Sala-Llinàs E. In Vivo and In Vitro Pro-Fibrotic Response of Lung-Resident Mesenchymal Stem Cells from Patients with Idiopathic Pulmonary Fibrosis. Cells 2024; 13 [PMID: 38247851 DOI: 10.3390/cells13020160]
- Hu C, Li L. Melatonin plays critical role in mesenchymal stem cell-based regenerative medicine in vitro and in vivo. *Stem Cell Res Ther* 2019;
 10: 13 [PMID: 30635065 DOI: 10.1186/s13287-018-1114-8]
- Gemayel J, Chaker D, El Hachem G, Mhanna M, Salemeh R, Hanna C, Harb F, Ibrahim A, Chebly A, Khalil C. Mesenchymal stem cellsderived secretome and extracellular vesicles: perspective and challenges in cancer therapy and clinical applications. *Clin Transl Oncol* 2023; 25: 2056-2068 [PMID: 36808392 DOI: 10.1007/s12094-023-03115-7]
- 34 Gad ES, Salama AAA, El-Shafie MF, Arafa HMM, Abdelsalam RM, Khattab M. The Anti-fibrotic and Anti-inflammatory Potential of Bone Marrow-Derived Mesenchymal Stem Cells and Nintedanib in Bleomycin-Induced Lung Fibrosis in Rats. *Inflammation* 2020; 43: 123-134 [PMID: 31646446 DOI: 10.1007/s10753-019-01101-2]
- 35 Li S, Zhang J, Feng G, Jiang L, Chen Z, Xin W, Zhang X. The Emerging Role of Extracellular Vesicles from Mesenchymal Stem Cells and Macrophages in Pulmonary Fibrosis: Insights into miRNA Delivery. *Pharmaceuticals (Basel)* 2022; 15 [PMID: 36297388 DOI: 10.3390/ph15101276]
- 36 Guo Q, Lu Y, Cheng X, Xiao F, Zhang Q, Gao P, Du L. Hepatocyte Growth Factor Delivered by Nanocomposites for Gene Therapy of Bleomycin-Induced Pulmonary Fibrosis in Rats. *Curr Drug Deliv* 2023; 20: 1368-1379 [PMID: 35702802 DOI: 10.2174/1567201819666220613145417]
- 37 Chen H, Luo Y, Zhu Y, Ye Y, Chen D, Song X, Xiao Z, Liu M, Li S. Enhanced secretion of hepatocyte growth factor in human umbilical cord mesenchymal stem cells ameliorates pulmonary fibrosis induced by bleomycin in rats. *Front Pharmacol* 2022; 13: 1070736 [PMID: 36726784 DOI: 10.3389/fphar.2022.1070736]
- 38 Gharbia S, Nazarie SR, Dinescu S, Balta C, Herman H, Peteu VE, Gherghiceanu M, Hermenean A, Costache M. Adipose-Derived Stem Cells (ADSCs) Supplemented with Hepatocyte Growth Factor (HGF) Attenuate Hepatic Stellate Cell Activation and Liver Fibrosis by Inhibiting the TGF-β/Smad Signaling Pathway in Chemical-Induced Liver Fibrosis Associated with Diabetes. *Cells* 2022; **11** [PMID: 36359733 DOI: 10.3390/cells11213338]
- 39 Chen WX, Zhou J, Zhou SS, Zhang YD, Ji TY, Zhang XL, Wang SM, Du T, Ding DG. Microvesicles derived from human Wharton's jelly mesenchymal stem cells enhance autophagy and ameliorate acute lung injury *via* delivery of miR-100. *Stem Cell Res Ther* 2020; 11: 113 [PMID: 32169098 DOI: 10.1186/s13287-020-01617-7]
- 40 Keshtkar S, Azarpira N, Ghahremani MH. Mesenchymal stem cell-derived extracellular vesicles: novel frontiers in regenerative medicine. Stem Cell Res Ther 2018; 9: 63 [PMID: 29523213 DOI: 10.1186/s13287-018-0791-7]
- 41 Hormozi A, Hasanzadeh S, Ebrahimi F, Daei N, Hajimortezayi Z, Mehdizadeh A, Zamani M. Treatment with Exosomes Derived from Mesenchymal Stem Cells: A New Window of Healing Science in Regenerative Medicine. *Curr Stem Cell Res Ther* 2024; 19: 879-893 [PMID: 37622719 DOI: 10.2174/1574888X18666230824165014]
- 42 Giovannelli L, Bari E, Jommi C, Tartara F, Armocida D, Garbossa D, Cofano F, Torre ML, Segale L. Mesenchymal stem cell secretome and extracellular vesicles for neurodegenerative diseases: Risk-benefit profile and next steps for the market access. *Bioact Mater* 2023; 29: 16-35 [PMID: 37456581 DOI: 10.1016/j.bioactmat.2023.06.013]
- 43 Bari E, Ferrarotti I, Saracino L, Perteghella S, Torre ML, Richeldi L, Corsico AG. Mesenchymal Stromal Cell Secretome for Post-COVID-19 Pulmonary Fibrosis: A New Therapy to Treat the Long-Term Lung Sequelae? *Cells* 2021; 10 [PMID: 34068958 DOI: 10.3390/cells10051203]

Raisbidena® WJSC | https://www.wjgnet.com



Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

