World Journal of *Diabetes*

World J Diabetes 2024 September 15; 15(9): 1829-2000





Published by Baishideng Publishing Group Inc

World Journal of Diabetes

Conten	Monthly Volume 15 Number 9 September 15, 2024
	EDITORIAL
1829	Exploring the genetic basis of childhood monogenic diabetes
	Sanyal D
1833	New therapy for metabolic syndrome: Gut microbiome supplementation
	Qureshi W, Dar MA, Rather MY
1837	MicroRNA-630: A potential guardian against inflammation in diabetic kidney disease
	Al Madhoun A
1842	Link between periodontitis and diabetic retinopathy: Inflammatory pathways and clinical implications
	Zhao Y, Shen QQ
1847	Macrophage modulation with dipeptidyl peptidase-4 inhibitors: A new frontier for treating diabetic cardiomyopathy?
	Mohammadi S, Al-Harrasi A
1853	Inflammatory markers, oxidative stress, and mitochondrial dynamics: Repercussions on coronary artery disease in diabetes
	Tatmatsu-Rocha JC, Mendes-Costa LS
	FIELD OF VISION
1858	Rapid correction of chronic hyperglycemia and bone remodeling, warning against overdoing
	Dardari D, Segurence B
	REVIEW
1862	Selection of dialysis methods for end-stage kidney disease patients with diabetes
	Hu YH, Liu YL, Meng LF, Zhang YX, Cui WP
	MINIREVIEWS
1874	Gut microbiome: A revolution in type II diabetes mellitus
	Jeyaraman M, Mariappan T, Jeyaraman N, Muthu S, Ramasubramanian S, Santos GS, da Fonseca LF, Lana JF
	ORIGINAL ARTICLE
	Case Control Study
1889	Platelet indices as predictors of poor glucoregulation in type 2 diabetes mellitus adults at Bishoftu General Hospital, Ethiopia
	Regassa DA, Berihun GA, Habtu BF, Haile WB, Nagaash RS, Kiya GT

Contents

Retrospective Study

1903 Non-linear relationship between age and subfoveal choroidal thickness in Chinese patients with proliferative diabetic retinopathy

Lei CY, Xie JY, Ran QB, Zhang MX

Basic Study

1916 Corilagin alleviates podocyte injury in diabetic nephropathy by regulating autophagy via the SIRT1-AMPK pathway

Lou Y, Luan YT, Rong WQ, Gai Y

1932 cNPAS2 induced β cell dysfunction by regulating KANK1 expression in type 2 diabetes Yin YB, Ji W, Liu YL, Gao QH, He DD, Xu SL, Fan JX, Zhang LH

1942 Molecular mechanisms of Buqing granule for the treatment of diabetic retinopathy: Network pharmacology analysis and experimental validation

Yang YF, Yuan L, Li XY, Liu Q, Jiang WJ, Jiao TQ, Li JQ, Ye MY, Niu Y, Nan Y

1962 Dexmedetomidine ameliorates diabetic intestinal injury by promoting the polarization of M2 macrophages through the MMP23B pathway

Lu M, Guo XW, Zhang FF, Wu DH, Xie D, Luo FQ

Bone marrow-derived mesenchymal stem cell-derived exosome-loaded miR-129-5p targets high-mobility 1978 group box 1 attenuates neurological-impairment after diabetic cerebral hemorrhage

Wang YY, Li K, Wang JJ, Hua W, Liu Q, Sun YL, Qi JP, Song YJ



Contents

Monthly Volume 15 Number 9 September 15, 2024

ABOUT COVER

Peer Review of World Journal of Diabetes, Tao-Hsin Tung, PhD, Researcher, Director, Epidemiologist, Evidencebased Medicine Center, Taizhou Hospital of Zhejiang Province Affiliated to Wenzhou Medical University, Taizhou 317000, Zhejiang Province, China. dongdx@enzemed.com .

AIMS AND SCOPE

The primary aim of World Journal of Diabetes (WJD, World J Diabetes) is to provide scholars and readers from various fields of diabetes with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJD mainly publishes articles reporting research results and findings obtained in the field of diabetes and covering a wide range of topics including risk factors for diabetes, diabetes complications, experimental diabetes mellitus, type 1 diabetes mellitus, type 2 diabetes mellitus, gestational diabetes, diabetic angiopathies, diabetic cardiomyopathies, diabetic coma, diabetic ketoacidosis, diabetic nephropathies, diabetic neuropathies, Donohue syndrome, fetal macrosomia, and prediabetic state.

INDEXING/ABSTRACTING

The WID is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for WJD as 4.2; JIF without journal self cites: 4.1; 5-year JIF: 4.2; JIF Rank: 40/186 in endocrinology and metabolism; JIF Quartile: Q1; and 5year JIF Quartile: Q2.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yu-Xi Chen.; Production Department Director: Xu Guo; Cover Editor: Jia-Ping Yan.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Diabetes	https://www.wignet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 1948-9358 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
June 15, 2010	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Monthly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Lu Cai, Md. Shahidul Islam, Michael Horowitz	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/1948-9358/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
September 15, 2024	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



WJD

World Journal of Diabetes

Submit a Manuscript: https://www.f6publishing.com

World J Diabetes 2024 September 15; 15(9): 1874-1888

DOI: 10.4239/wjd.v15.i9.1874

ISSN 1948-9358 (online)

MINIREVIEWS

Gut microbiome: A revolution in type II diabetes mellitus

Madhan Jeyaraman, Tejaswin Mariappan, Naveen Jeyaraman, Sathish Muthu, Swaminathan Ramasubramanian, Gabriel Silva Santos, Lucas Furtado da Fonseca, José Fábio Lana

Specialty type: Endocrinology and metabolism

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification Scientific Quality: Grade B, Grade C, Grade C, Grade C, Grade D Novelty: Grade B, Grade C, Grade C Creativity or Innovation: Grade B, Grade C, Grade C

Scientific Significance: Grade B, Grade B, Grade C

P-Reviewer: Bortolotti M; Han J; Horowitz M; Lie Z; Papazafiropoulou A

Received: April 18, 2024 Revised: June 11, 2024 Accepted: July 18, 2024 Published online: September 15, 2024 Processing time: 130 Days and 18.9 Hours



Madhan Jeyaraman, Naveen Jeyaraman, Department of Orthopaedics, ACS Medical College and Hospital, Dr MGR Educational and Research Institute, Chennai 600077, Tamil Nadu, India

Madhan Jeyaraman, Gabriel Silva Santos, Lucas Furtado da Fonseca, José Fábio Lana, Department of Orthopaedics, Brazilian Institute of Regenerative Medicine, Indaiatuba 13334-170, São Paulo, Brazil

Madhan Jeyaraman, Sathish Muthu, Department of Orthopaedics, Orthopaedic Research Group, Coimbatore 641045, Tamil Nadu, India

Tejaswin Mariappan, Department of Community Medicine, Government Stanley Medical College and Hospital, Chennai 600001, Tamil Nadu, India

Sathish Muthu, Department of Orthopaedics, Government Medical College, Karur 639004, Tamil Nadu, India

Sathish Muthu, Department of Biotechnology, Faculty of Engineering, Karpagam Academy of Higher Education, Coimbatore 641021, Tamil Nadu, India

Swaminathan Ramasubramanian, Department of Orthopaedics, Government Medical College, Omandurar Government Estate, Chennai 600002, Tamil Nadu, India

Corresponding author: Madhan Jeyaraman, MS, PhD, Assistant Professor, Research Associate, Department of Orthopaedics, ACS Medical College and Hospital, Dr MGR Educational and Research Institute, Velappanchavadi, Chennai 600077, Tamil Nadu, India. madhanjeyaraman@gmail.com

Abstract

Type II diabetes mellitus (T2DM) has experienced a dramatic increase globally across countries of various income levels over the past three decades. The persistent prevalence of T2DM is attributed to a complex interplay of genetic and environmental factors. While numerous pharmaceutical therapies have been developed, there remains an urgent need for innovative treatment approaches that offer effectiveness without significant adverse effects. In this context, the exploration of the gut microbiome presents a promising avenue. Research has increasingly shown that the gut microbiome of individuals with T2DM exhibits distinct differences compared to healthy individuals, suggesting its potential role in the disease's pathogenesis and progression. This emerging field offers diverse applications, particularly in modifying the gut environment through the administration of prebiotics, probiotics, and fecal microbiome transfer. These inter-



ventions aim to restore a healthy microbiome balance, which could potentially alleviate or even reverse the metabolic dysfunctions associated with T2DM. Although current results from clinical trials have not yet shown dramatic effects on diabetes management, the groundwork has been laid for deeper investigation. Ongoing and future clinical trials are critical to advancing our understanding of the microbiome's impact on diabetes. By further elucidating the mechanisms through which microbiome alterations influence insulin resistance and glucose metabolism, researchers can develop more targeted interventions. The potential to harness the gut microbiome in developing new therapeutic strategies offers a compelling prospect to transform the treatment landscape of T2DM, potentially reducing the disease's burden significantly with approaches that are less reliant on traditional pharmaceuticals and more focused on holistic, systemic health improvements.

Key Words: Type II diabetes; Gut microbiome; Probiotics; Prebiotics; Fecal microbiota transplantation

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Type II diabetes mellitus (T2DM) has surged globally, driven by genetic and environmental factors. Amidst pharmaceutical options, exploring the gut microbiome stands out. Research reveals distinct microbiome differences in T2DM, suggesting its role in pathogenesis. Interventions such as prebiotics, probiotics, and fecal transfers aim to restore balance. While clinical trials have not shown dramatic effects yet, ongoing research holds promise. Understanding microbiome mechanisms could revolutionize T2DM treatment, emphasizing holistic health approaches.

Citation: Jeyaraman M, Mariappan T, Jeyaraman N, Muthu S, Ramasubramanian S, Santos GS, da Fonseca LF, Lana JF. Gut microbiome: A revolution in type II diabetes mellitus. World J Diabetes 2024; 15(9): 1874-1888 URL: https://www.wjgnet.com/1948-9358/full/v15/i9/1874.htm DOI: https://dx.doi.org/10.4239/wjd.v15.i9.1874

INTRODUCTION

The human gut microbiome is a complex ecosystem, hosting thousands of bacterial species, each exerting a unique influence on host metabolism. This intricate interplay involves a variety of signaling molecules derived from dietary components that are metabolized by these microbiota. The role of the gut microbiome extends beyond digestion, as it actively engages with multiple bodily systems to maintain physiological homeostasis[1]. In the context of type II diabetes mellitus (T2DM), the gut microbiome exhibits notable changes, termed dysbiosis, where there is an increase in bacteria that negatively impact metabolic health and a decrease in beneficial bacteria^[2,3]. This shift can lead to a cascade of health issues, including metabolic disorders, cardiovascular complications, neuronal diseases, and various inflammatory conditions, as depicted in Figure 1[4,5].

Despite understanding the broad impacts of gut microbiome dysbiosis, specific pathways, and interactions that lead to T2DM remain underexplored. The mechanisms through which microbial products and toxins contribute to increased intestinal permeability and subsequent inflammation are not fully elucidated. Furthermore, the exact nature of changes in the production of short-chain fatty acids (SCFAs), lipopolysaccharides, and bile acids in diabetics requires detailed investigation[6-8]. These gaps in knowledge hinder the full exploitation of the gut microbiome as a target for therapeutic interventions. Understanding these pathways in greater depth could unveil novel strategies to manage or potentially reverse the effects of dysbiosis in diabetic patients. The primary objective of this review is to synthesize current understandings of the microbiome's role in T2DM, with a particular focus on the pathophysiological changes associated with dysbiosis. We aimed to identify and discuss novel therapeutic strategies that target the microbiome to ameliorate the symptoms and complications associated with T2DM.

GUT MICROBIOTA AND ITS EFFECTS ON T2DM

Organisms with a positive correlation with the disease

The microbiome composition in individuals with diabetes, particularly T2DM, exhibits significant variations compared to non-diabetic individuals. Notably, the phyla Proteobacteria and Firmicutes[3] are predominantly observed in diabetics. Within Firmicutes, there is a noted increase in the genus Ruminococcus and a decrease in Clostridium species[3], as demonstrated in Figure 2, which outlines the microbiota profile variation in T2DM. However, observations in patients of different ethnicities may vary and needs further investigation.

Research highlights that the Firmicutes-to-Bacteroides ratio, which is typically below 0.8 in healthy individuals, is elevated in those with T2DM and obesity. This altered ratio is indicative of microbial dysbiosis associated with these conditions[9-11]. Using a Predomics approach, three species have been identified as significant biomarkers for T2DM: Enterococcus faecium, Eubacterium linosum, and Absiella spp. [12]. Eubacterium linosum, an anaerobic acetogenic bacterium, is



WJD | https://www.wjgnet.com



Figure 1 Communication channels of the gut with various systems of the body through metabolites produced by the gut microbiota. LPS: Lipopolysaccharides; SCFAs: Short-chain fatty acids.



Figure 2 Microbiota profile variation in type II diabetes mellitus. DM: Diabetes mellitus.

particularly noteworthy for its metabolic functions, including acetate production, sulfate reduction, and the degradation of urea and arginine. These metabolic activities contribute to chronic low-grade inflammation *via* pro-inflammatory cytokines and metabolites, a hallmark of T2DM[13,14]. Additionally, the production of acetate through the fermentation of galacto-oligosaccharides or inulin is linked to alterations in insulin sensitivity and weight gain[8,15]. Certain species within the Firmicutes phylum are recognized for their enhanced capacity to break down complex sugars and fatty acids, thereby potentially increasing the risk of obesity and T2DM[16]. On the other hand, some species within the *Proteobacteria phylum*, such as *Fusobacterium spp.*, are implicated in protein fermentation and degradation, leading to dysbiosis. *Fusobacterium spp.* also contributes to pathogenicity by inducing inflammatory cytokines [interleukin (IL)-1 β , tumor necrosis factor (TNF)- α , IL-17, *etc.*], which further exacerbate the inflammatory state[17,18].

Organisms with a negative correlation with the disease

Several organisms exhibit a negative correlation with T2DM, primarily through mechanisms that involve the production of anti-inflammatory and immunoregulatory metabolites such as butyrate and propionate[14]. Among these, the genus Bifidobacterium is notably protective against T2DM, attributed to its cross-feeding mechanisms that enhance metabolic

wJD https://www.wjgnet.com

health[19]. Animal studies further corroborate the benefits of various species within this genus, demonstrating an increase in glucose tolerance[20-24].

Another significant genus negatively correlated with T2DM is Bacteroides. This group, which includes species like *Bacteroides uniformis* and *Bacteroides acidifaciens*, is known to improve glucose tolerance and insulin sensitivity, and is instrumental in managing metabolic diseases exacerbated by poor diet[25,26]. Despite their reduced presence in individuals with T2DM, other genera such as *Faecalibacterium*, *Akkermansia*, and *Roseburia* also exhibit similar negative correlations with the disease, though they are not as abundantly reported[27]. *Lactobacillus* species display variable associations with T2DM; however, species like *Lactobacillus plantarum*[28], *Lactobacillus casei*[29], and *Lactobacillus reuteri*[30] have been shown to improve symptoms when administered as probiotics. In synergy with Bifidobacterium, they confer a collective protective effect against the disease[31-37]. SCFAs such as butyrate play crucial roles beyond their metabolic functions; they regulate pancreatic beta-cell activity, reduce hepatic gluconeogenesis, and modulate immune system functions. The reduction in these critical microbiota directly contributes to the pathogenesis of T2DM.

Gut metabolites

The fermentation of nutrients by gut microbiota leads to the production of metabolites like SCFAs (butyrate, propionate, acetate), branched-chain amino acids, indoles, imidazoles, and succinates. These are predominantly produced by genera including *Bacteroides, Akkermansia, Prevotella, Faecalibacterium, Lactobacillus, Clostridium,* and *Propionibacterium*. These metabolites have diverse and significant interactions within the gut environment, as illustrated in Figure 3.

Pathogenesis of dysbiosis contributing to worsening T2DM

Alteration in permeability and steatosis: Changes in the gut microbiome composition have been associated with T2DM progression, particularly a decrease in species such as *Bacteroides* and *Akkermansia*. These microbes are crucial for the regulation of tight junction proteins including occludin and claudin, which maintain intestinal barrier integrity[4,38]. The disruption of these tight junctions leads to increased intestinal permeability. This, in turn, facilitates enhanced nutrient absorption and altered glycemic control, propelling the progression toward steatosis. Concurrently, the reduction in microbial populations that regulate hepatic gluconeogenesis exacerbates liver steatosis.

Inflammation and altered lipopolysaccharides: Increased intestinal permeability allows greater absorption of dietary products and bacterial endotoxins, such as lipopolysaccharides, into the bloodstream. This elevates the production of proinflammatory cytokines, including TNF- α , IL-1 β , IL-6, IL-17, and other cytokines linked to T helper (Th)1, Th2, and Th17 responses. This state of endotoxemia drives systemic inflammation and reactive oxygen species production, leading to the destruction of pancreatic beta cells and the onset of insulin resistance[4,38]. The absence of protective microbial effects due to altered microbiota composition, such as those from *Roseburia intestinalis*, *Bacteroides fragilis*, *Akkermansia spp.*, and *Lactobacillus casei*, further compounds the problem. These species induce anti-inflammatory cytokines like IL-10, which may help mitigate low-grade inflammation and potentially improve insulin sensitivity. *Roseburia intestinalis*, for example, also promotes the production of IL-22, enhancing insulin sensitivity, and transforming growth factor-beta, inhibiting inflammatory processes.

Fatty acid metabolism: The altered composition of gut flora observed in T2DM patients may affect the production of SCFAs. These SCFAs, such as butyrate and propionate, are essential in promoting fatty acid oxidation by inhibiting the expression of peroxisome proliferator-activated receptor-gamma[39]. The absence of these regulatory mechanisms is evident in conditions like increased serum malonaldehyde, a marker for lipid oxidation, which is typically reduced by *Lactobacillus casei* and *Akkermansia muciniphila* in experimental models, but is elevated in diabetic subjects[40,41]. This disruption leads to enhanced fat accumulation in adipose tissue and the liver[42-44], triggering a modification in bile acid metabolism. Such alterations further propagate inflammation and microbial imbalance, perpetuating a cyclical exacerbation of T2DM pathogenesis[4]. Having discussed the potential associations between gut metabolites from microbiome dysbiosis and T2DM progression, specific targets to ameliorate their impact remains an area of future research.

Changes of the gut microbiome at different stages/complications of T2DM

The changes in the gut microbiome studied through various human and animal studies, a targeted replenishment of gut microbiota, can lead to delay or prevention in complications of diabetes through supplementation by probiotics, prebiotics, and fecal microbial transplantation, which in turn restores the stability of gut microbiota, as mentioned in Table 1.

Effects of diabetic medications on the gut microbiome

Metformin: Metformin is well-documented for its favorable modifications to the gut microbiome. According to Ismail and Evans-Molina[45], metformin treatment leads to changes in the composition of gut microbiota, enhancing the production of SCFAs and altering bile acids. These changes result in increased levels of glucagon-like peptide-1 (GLP-1), which promotes insulin secretion. Additionally, metformin helps normalize the altered microbial community by reversing the Firmicutes/Bacteroides ratio, which is often disrupted in diabetic conditions, thus restoring the microbiome toward a healthier state[4].

Sulfonylureas: The effects of sulfonylureas on the gut microbiome are less clear, with studies presenting conflicting data. One study reports that the use of sulfonylureas is associated with increased levels of phenylalanine and tryptophan, suggesting a potential impact on the microbiome[46]. However, another study found no significant changes in the composition of the gut microbiota with sulfonylurea treatment[47]. This indicates that the influence of sulfonylureas on



→ WJD https://www.wjgnet.com

Table 1 Dysbiosis observed in various stages of diabetes mellitus

Complication/stage	Dysbiosis observed				
observed	Decreased	Increased			
Diabetic nephropathy	Lactobacillus, Bifidobacterium, Bacteroides, Prevotella, Roseburia, Ruminococcaceae, and Faecalibacterium	Enterococcus, Enterobacteriaceae, Clostridaceae, Klebsiella, and Parabacterides			
Diabetic neuropathy	Bacteroides and Faecalibacterium	Escherichia, Blautia, Ruminococcus torques, and Lachno- clostridium			
Diabetic retinopathy	Bacteroidetes and Actinobacteria	Escherihia, Enterobacter, and Acidaminococcus			
Cerebrovascular disease	Lachnospiraceae, Ruminococcaceae, Bacteroidetes, Prevotella, and Faecalibacterium	Enterobacteriaceae, Veillonellaceae, Bifidobacterium, Lactoba- cillus, and Oscillobacter			
Cardiovascular disease	Roseburia, Eubacterium spp, Bacteroides and Faecalibacterium	<i>Collinsella, Escherichia-Shigella, Enterococcus,</i> and the ratio of <i>Firmicutes</i> to <i>Bacteroides</i>			
Peripheral vascular disease	-	Firmicutes, Actinobacteria, Verrucomicrobia, and Proteo- bacteria			



Figure 3 Metabolite breakdown products of gut microbiome. GLP-1: Glucagon-like peptide-1.

the gut microbiome may vary depending on other factors such as dosage, duration of treatment, and individual patient microbiome composition.

Alpha-glucosidase inhibitors: Alpha-glucosidase inhibitors prevent the breakdown of oligosaccharides in the small intestine, thereby increasing their availability as nutrients for gut bacteria. This alteration in nutrient availability promotes the growth of beneficial microbes such as Bacteroides, Lactobacillus, and Faecalibacterium, while reducing populations of potentially pathogenic bacteria like Ruminococcus and Butyricicoccus[48]. These microbial shifts are associated with changes in bile acid profiles and improvements in prognostic factors for T2DM.

GLP-1 agonists: GLP-1 agonists, which slow gastric emptying, induce significant shifts in the gut microbial community. Studies indicate a reduction in obesity-promoting organisms within the Firmicutes phylum and an increase in populations of Verrucomicrobia and microbes from the orders Clostridiales and Bacteroidales [49]. These drugs also promote the growth of SCFA-producing bacteria such as Bifidobacterium and Bacteroides, further supporting glycemic control and metabolic health[50].

Sodium-glucose co-transporter type 2 inhibitors: Sodium-glucose co-transporter type 2 inhibitors, such as sotagliflozin, impact the gut microbiome by decreasing the Firmicutes/Bacteroides ratio and enhancing fatty acid production[51]. This alteration not only affects the microbial landscape but also has broader implications for energy metabolism and insulin sensitivity in T2DM patients. A summary of the effects of diabetic medications on the gut microbiome is tabulated in



Zaishideng® WJD | https://www.wjgnet.com

Table 2.

Influence of diet and exercise on gut microbiome

The influence of diet and exercise on the gut microbiome has significant implications for the management of conditions such as T2DM. Various dietary patterns and physical activities have been shown to differentially affect the composition and functionality of the gut microbiota, with direct consequences on metabolic health (Table 3).

Dietary influences

A diet rich in plant-based fibers such as cellulose, inulin, pectin, and dextrin has been found to confer multiple health benefits, including reduced insulin resistance, lower serum cholesterol levels, and stable blood glucose levels. These effects are largely attributed to the enrichment of beneficial gut bacteria that negatively correlate with metabolic diseases. These bacteria produce SCFAs such as butyrates and propionates, which play a crucial role in reducing the production of pro-inflammatory cytokines, thereby mitigating inflammation associated with T2DM[4,52]. Conversely, diets high in fats and proteins, particularly from animal sources, tend to foster a pro-inflammatory gut environment. This is largely due to an increase in lipopolysaccharides, which are potent inflammatory agents. However, it is important to note that not all protein-rich diets have adverse effects. Diets rich in plant proteins can enhance the proliferation of beneficial microbes such as Lactobacillus spp. and Bifidobacterium spp. In contrast, diets high in animal proteins tend to increase levels of Bacteroides and Bilophila, which have mixed effects on health depending on the overall dietary context^[53]. The Mediterranean diet, which is rich in vegetables, fruits, nuts, seeds, and whole grains, has been specifically noted for its positive alterations to the gut microbiota. Adherence to this diet enhances the production of SCFAs, thereby improving insulin sensitivity. Observational studies in obese patients have shown increases in beneficial genera such as Roseburia and Oscillospira, alongside a reduction in Prevotella spp., which are associated with dysbiosis and metabolic disturbances. The beneficial shifts in the gut microbiome are contingent upon strict adherence to the dietary regimen[54].

Exercise influence

Physical exercise also plays a critical role in modulating the gut microbiome. Low-intensity physical activities have been associated with favorable changes in the gut microbiota composition, including an increase in the Bacteroides/Firmicutes ratio[55-57]. This shift helps mitigate the adverse effects of an unhealthy diet and contributes to overall metabolic health. Exercise-induced modifications to the microbiome can enhance the resilience of the gut ecosystem, promoting a balance that favors metabolic health and potentially reducing the risk or severity of T2DM.

THERAPIES FOR T2DM BASED ON THE GUT MICROBIOME

Nutraceutical agents

Lactobacillus and Bifidobacterium are among the most extensively studied probiotics, known for their beneficial effects on glycemic control. Meta-analyses have demonstrated that supplementation with these microbes can significantly decrease hemoglobin A1c, fasting blood glucose, and markers of oxidative stress[58]. Additionally, the introduction of Akkermansia muciniphila, a recent focus in microbiome research, has shown promising results. In murine models, this bacterium has been found to increase GLP-1 levels in colon cells, enhance glucose tolerance, and maintain gut barrier integrity, thereby reducing inflammation and ameliorating liver damage[59]. Research indicates that Akkermansia muciniphila also modulates the Firmicutes/Bacteroides ratio, leading to increased butyrate production. These changes collectively contribute to reduced oxidative stress and improved lipid profiles and glucose tolerance[39].

Fecal microbiota transplantation

Beyond probiotics, fecal microbiota transplantation (FMT) represents a more direct method for altering the gut microbiome. Initially popularized through its efficacy in treating Clostridium difficile infections, which often arise from chronic antibiotic use and the resultant suppression of native flora, FMT has since broadened its clinical applications[60]. Data from various human studies suggest that FMT can decrease inflammatory markers and increase the production of secondary bile acids, although its effects on insulin resistance have been relatively mild[61]. This indicates that, while FMT holds potential, its full spectrum of therapeutic benefits in T2DM remains to be fully elucidated through additional clinical trials. Despite its potential, FMT carries risks, particularly the possibility of transferring a dysbiotic microbiome and infectious pathogens. Therefore, while FMT is a promising tool in the arsenal against metabolic diseases, its application must be approached with caution, ensuring rigorous screening and monitoring protocols to mitigate these risks. A summary of evidence of the role of the gut microbiome in T2DM is tabulated in Table 4.

Challenges and limitations

One of the principal challenges in leveraging the gut microbiome for diabetes treatment is the inherent variability in microbial composition among individuals. Factors such as genetics, diet, age, and environment significantly influence the gut microbiota, creating a highly personalized microbial ecosystem. This variability can affect the efficacy of probiotic treatments, as the same probiotic strains may not produce identical effects in different individuals. The lack of standardization in probiotic formulations poses another significant challenge. Probiotics are available in various forms, from dietary supplements to fortified foods, with considerable differences in strain specificity, viability, and concentration. These discrepancies can lead to inconsistent study results and confusion about their clinical applicability. Moreover, the



Saishidena® WJD | https://www.wjgnet.com

Table 2 Effects of diabetic medications on gut microbiome			
Medication	Effects on microbiome	Observed outcomes	
Metformin[45]	Enhances SCFA production, normalizes Firmicutes/Bacteroides ratio	Increased GLP-1 levels, improved insulin secretion	
Sulfonylureas[46,47]	Conflicting data on impact	Variable influence on microbiome, potential increase in phenylalanine and tryptophan levels	
Alpha-glucosidase inhibitors[<mark>48</mark>]	Increases nutrient availability for beneficial bacteria	Growth of beneficial microbes like Bacteroides, improvement in T2DM prognostic factors	
GLP-1 agonists[49,50]	Changes in gastric emptying rates influence microbiota	Reduction in obesity-promoting organisms, increase in beneficial microbes like Bifidobacterium	
SGLT-2 inhibitors[51]	Alters microbial ratios favorably	Reduction in Firmicutes/Bacteroides ratio, enhanced fatty acid production	

DM: Diabetes mellitus; GLP-1: Glucagon-like peptide-1; SCFA: Short-chain fatty acid; SGLT-2: Sodium-glucose co-transporter type 2.

Table 3 Impact of diet and exercise on the gut microbiome in type 2 diabetes mellitus			
Factor	Description	Beneficial effects	
Diet	High fiber plant-based foods[4,52]	Decrease in insulin resistance, stabilization of blood glucose levels, reduction in serum cholesterol	
	High-fat and protein diets[53]	Increase in pro-inflammatory markers; variable effects based on protein source (plant vs animal)	
	Mediterranean diet[54]	Improvement in SCFA production, enhanced insulin sensitivity, increased beneficial genera like Roseburia	
Exercise	Low-intensity physical activity[55- 57]	Favorable shifts in microbiota composition, improvement in metabolic health markers	

SCFA: Short-chain fatty acid.

regulation of probiotics varies by region, affecting the quality and safety of available products.

While short-term studies have demonstrated the potential benefits of probiotics in managing T2DM, long-term safety and efficacy remain underexplored. Questions about the optimal duration of probiotic therapy, long-term side effects, and the sustainability of beneficial effects need to be addressed through longitudinal studies. A deeper mechanistic understanding of how probiotics interact with both the gut microbiota and the host is crucial. Current knowledge about the pathways through which probiotics influence metabolic health, including their effects on inflammation, insulin sensitivity, and lipid metabolism, is still rudimentary. Enhanced mechanistic insights would facilitate the development of more targeted and effective therapeutic strategies. Interactions between probiotics to affect drug metabolism and efficacy needs thorough investigation to avoid adverse effects and ensure complementary therapeutic outcomes.

Future directions

Advancing a personalized medicine approach in microbiome research could significantly enhance the efficacy of treatments. By understanding individual microbiome profiles, treatments can be tailored to optimize microbial composition and functionality. This approach would involve integrating detailed genomic, metabolic, and dietary data to predict individual responses to specific probiotic strains. Research into next-generation probiotics, which are specifically engineered or selected based on their beneficial characteristics, is a promising direction. These could include not only bacteria but also other components of the microbiota such as beneficial viruses or fungi that play a role in metabolic regulation.

Combining probiotics with other therapeutic modalities, such as dietary interventions, pharmacotherapy, and lifestyle changes, could enhance overall treatment outcomes. For example, synchronizing probiotic supplementation with a fiberrich diet might amplify the beneficial effects on the gut microbiome. Conducting more rigorous and comprehensive clinical trials that focus on various demographic groups and different stages of diabetes is crucial. These studies should aim to clarify optimal dosages, treatment durations, and combinations of probiotic strains. Moreover, trials should also assess the impact of probiotics on diabetes complications, offering a broader understanding of their potential benefits. Enhancing healthcare provider and patient education about the potential benefits and limitations of probiotics as part of diabetes management is essential. Increased awareness can lead to more informed decision-making and better clinical outcomes. The limitations and future directions in gut microbiome research for T2DM are summarized in Table 5.

Raisbideng® WJD | https://www.wjgnet.com

Table 4 Summary of global studies of the gut microbiome in type 2 diabetes mellitus			
Ref.	Place of study	Probiotics used	Observed effects
Kumari et al[62], 2021	India	Lactobacillus spp.	Decreased HbA1C, insulin resistance, TNF- α , IL-1 β
		Lactococcus spp.	
		Propionibacterium spp.	
		Bifidobacterium spp.	
Zhao et al[63], 2020	China	Selenium enhanced	Decreased FG, HbA1C, and insulin levels and improves glucose
		Bifidobacterium spp.	tolerance and lipid profile
Palacios <i>et al</i> [64], 2020	Australia	Lactobacillus plantarum	Decreased FG, HbA1C, and insulin resistance
		Lactobacillus bulgaricus	
		Lactobacillus gasseri	
		Bifidobacterium breve	
		Bifidobacterium animalis sbsp. lactis	
		Bifidobacterium bifidum	
		S. thermophiles	
		S. boulardii	
Razmpoosh et al[65], 2019	Iran	Lactobacillus acidophilus	Decreased FG, insulin resistance, and increased HDL cholesterol
		Lactobacillus casei	
		Lactobacillus rhamnosus	
		Lactobacillus bulgaricus	
		Bifidobacterium breve	
		Bifidobacterium longum	
		Streptococcus thermophilus	
Madempudi <i>et al</i> [66], 2019	India	Lactobacillus salivarius	Decreased HbA1C and effects on lipid profile are not significant
		Lactobacillus casei	
		Lactobacillus plantarum	
		Lactobacillus acidophilus	
		Bifidobacterium breve	
		Bifidobacterium coagulans	
Sabico <i>et al</i> [67], 2019	Saudi Arabia	Bifidobacterium bifidum W23	Decreased FG, insulin resistance, total cholesterol, and trigly-
		Bifidobacterium lactis W52	certues
		Lactobacillus acidophilus W37	
		Lactobacillus brevis W63	
		Lactobacillus casei W56	
		Lactobacillus salivarius W24	
		Lactobacillus lactis W19	
		Lactobacillus lacis W58	
Mazruei Arani <i>et al</i> [68], 2019	Iran	Bacillus coagulans T4	Decreased FG, insulin resistance, CRP, and improves lipid profile
Mohseni <i>et al</i> [37], 2018	Iran	Lactobacillus acidophilus	Decreased FG, insulin resistance, inflammatory markers, and
		Bifidobacterium bifidum	improves upid prome
		Lactobacillus casei	
		Lactobacillus fementum	



Jeyaraman M et al. Gut microbiome and diabetes mellitus

Kobyliak <i>et al</i> [61], 2018	Ukraine	14 probiotic strains of Lactobacillus	Decreased HbA1C and insulin resistance
		Lactococcus	
		Bifidobacterium spp.	
		Propionibacterium	
		Acetobacter	
Kassaian <i>et al</i> [69], 2018	Iran	Lactobacillus acidophilus	Decreased FG, HbA1C, and insulin resistance
		Bifidobacterium lactis	
		Bifidobacterium bifidum	
		Bifidobacterium longum	
Mohseni <i>et al</i> [37], 2018	Iran	Bifidobacterium bifidum	Decrease FG, insulin resistance, total cholesterol, and increased
		Lactobacillus casei	GSH level
		Lactobacillus acidophilus	
Mofidi <i>et al</i> [35], 2017	Iran	Lactobacillus casei	Decreased FG and triglycerides
		Lactobacillus rhamnosus	
		Streptococcus thermophilus	
		Bifidobacterium breve	
		Lactobacillus acidophilus	
		Bifidobacterium longum	
		Lactobacillus bulgaricus	
Firouzi <i>et al</i> [36], 2017	Malaysia	Lactobacillus acidophilus	Decreased HbA1C and does not affect lipid profile
		Lactobacillus casei	
		Lactobacillus lactis	
		Bifidobacterium bifidum	
		Bifidobacterium longum	
		Bifidobacterium infantis	
Tajabadi-Ebrahimi et al	Iran	Lactobacillus acidophilus	Decreased FG, increased insulin sensitivity, and does not affect
[33], 2017		Lactobacillus casei	lipid profile
		Bifidobacterium bifidum	
Ebrahimi <i>et al</i> [70], 2017	Iran	Lactobacillus spp.	Decreased FG, HbA1C, and no effect on lipid profile
		Bifidobacterium spp.	
		Streptococcus thermophilus and fructo- oligosaccharide	
Asemi et al[71], 2016	Iran	Probiotic: Lactobacillus sporogenes	Decreased in serum insulin, insulin resistance, triglycerides and
		Prebiotic: Inulin, beta-carotene	increased GSH levels
Madjd <i>et al</i> [72], 2016	Iran	Lactobacillus acidophilus LA5	Decreased HbA1C, 2-h postprandial glucose, insulin resistance,
) L J		Bifidobacterium lactis BB12	total cholesterol, and LDL levels
Karamali <i>et al</i> [73] , 2016	Iran	Lactobacillus acidophilus	Decreased fasting glucose, insulin resistance, triglycerides,
		Lactobacillus casei	VLDL, and increased insulin sensitivity
		Bifidobacterium bifidum	
Ostadrahimi et al[74], 2015	Iran	Lactobacillus acidophilus	Decreased HbA1C, and FG and does not affect lipid profile
		Lactobacillus casei	
		Bifidobacterium lactis	
Eslamparast <i>et al</i> [75], 2014	Iran	Lactobacillus casei	Decreased FG, insulin resistance and has no effect on lipid
· · · · · · · · · · · · · · · · · · ·		Lactobacillus rhamnosus	profile



		Streptococcus thermophilus	
		Bifidobacterium breve	
		Lactobacillus acidophilus	
		Bifidobacterium longum	
		Lactobacillus bulgaricus	
Rajkumar <i>et al</i> [76], 2014	India	Bifidobacterium longum	Decreased FG, insulin resistance, total cholesterol, triglycerides,
		Bifidobacterium infantis	
		Bifidobacterium breve	
		Lactobacillus acidophilus	
		Lactobacillus paracasei	
		Lactobacillus bulgaricus	
		Lactobacillus plantarum	
		Streptococcus thermophilus	
Ivey <i>et al</i> [77], 2014	Australia	Lactobacillus acidophilus La5	Increased FG and insulin resistance
		Bifidobacterium lactis Bb12	
Mohamadshahi et al[78],	Iran	Bifidobacterium lactis Bb12	Decreased HbA1C
2014		Lactobacillus acidophilus	
Asemi et al[79], 2014	Iran	Probiotic: Viable & heat-resistant Lactobacillus sporogenes	Decreased FG, HbA1C, insulin resistance, and inflammatory markers
		Prebiotic: Inulin	
Asemi et al[34], 2013	Iran	Lactobacillus spp.	Decreased FG and increased insulin levels, total GSH levels, and
		Bifidobacterium spp.	LDL levels
		<i>Streptococcus spp.</i> Fructo-oligosaccharide	
Mazloom <i>et al</i> [80], 2013	Iran	Lactobacillus acidophilus	Decreased FG, insulin resistance, and improves fasting insulin
		Lactobacillus bulgaricus	
		Lactobacillus bifidum	
		Lactobacillus casei	
Shavakhi <i>et al</i> [<mark>81</mark>], 2013	Iran	Lactobacillus acidophilus	Decreased FG, triglycerides, and total cholesterol
		Lactobacillus casei	
		Lactobacillus rhamnosus	
		Lactobacillus bulgaricus	
		Bifidobacterium breve	
		Bifidobacterium longum	
		Streptococcus thermophilus	
Asemi et al[82], 2013	Iran	Lactobacillus acidophilus LA5	Decreased insulin resistance
		Bifidobacterium animalis BB12	
Asemi et al[34], 2013	Iran	Lactobacillus acidophilus	Decreased FG, increased insulin resistance, and LDL levels
		Lactobacillus casei	
		Lactobacillus rhamnosus	
		Lactobacillus bulgaricus	
		Bifidobacterium breve	
		Bifidobacterium longum	
		Streptococcus thermophiles	
		r · · · · · · · · · · · · · · · · · · ·	



Jeyaraman M et al. Gut microbiome and diabetes mellitus

Moroti <i>et al</i> [31], 2012	Brazil	Lactobacillus acidophilus	Decreased FG and increased HDL levels
		Bifidobacterium bifidum	
Ejtahed <i>et al</i> [83], 2012	Iran	Lactobacillus acidophilus La5	Decreased FG and HbA1C and does not affect lipid profile
		Bifidobacterium lactis Bb12	
Laitinen <i>et al</i> [84], 2009	Finland	Lactobacillus rhamnosus GG	Decreased FG, insulin resistance, and increased insulin
		Bifidobacterium lactis Bb12	sensitivity

CRP: C-reactive protein; FG: Fasting glucose; GSH: Glutathione; HbA1C: Hemoglobin A1c; HDL: High-density lipoprotein; IL: Interleukin; LDL: Lowdensity lipoprotein; TNF: Tumor necrosis factor; VLDL: Very low-density lipoprotein.

Table 5 Limitations and future directions in gut microbiome research for type 2 diabetes mellitus

Limitations	Description	Future directions
Variability in microbial composition	Individual differences in microbiome composition complicate standard treatment outcomes	Personalized microbiome interventions: Develop treatments based on individual microbiome assessments to optimize efficacy
Lack of standardization	Inconsistencies in probiotic formulations affect study comparability and clinical applicability	Standardization of products: Establish regulations and standards for probiotic formulations to ensure quality and consistency
Short-term focus	Most studies have short duration and do not address long-term safety and effectiveness	Longitudinal studies: Conduct long-term studies to assess the sustained effects and safety of microbiome-based interventions
Incomplete mechanistic understanding	The pathways through which the microbiome influences diabetes are not fully elucidated	Mechanistic research: Deepen research into the biochemical interactions within the gut microbiome that affect diabetes pathogenesis and treatment
Drug-microbiome interactions	Potential interactions between probiotics and anti-diabetic medications are not well understood	Interaction studies: Explore how probiotics interact with common diabetic medications to refine treatment protocols
Regulatory hurdles	The global regulatory landscape for probiotics and microbiome therapies varies significantly	Harmonize regulations: Work toward an international consensus on the regulation of microbiome therapies to facilitate global research and application

CONCLUSION

T2DM significantly impacts the gut microbiota, leading to dysbiosis, which in turn promotes inflammation and contributes to the development of insulin resistance. The evidence gathered from various trials and studies underscores the pivotal role of nutraceuticals in restoring a balanced gut flora, enhancing insulin sensitivity, and reducing insulin resistance. These beneficial effects have been consistently observed in animal models and, to a lesser extent, in human studies. Although the initial outcomes from human trials have not met all expectations, they lay a solid foundation for future research. Continued exploration and more targeted studies are essential to refine these interventions and fully assess their therapeutic potential. With ongoing advancements and deeper insights into gut microbiome interactions, nutraceuticals hold promise as a novel and effective treatment strategy to manage T2DM. As research progresses, these natural compounds will become an integral part of comprehensive diabetes management, potentially shifting current therapeutic paradigms.

FOOTNOTES

Author contributions: Jeyaraman M, Mariappan T, and Jeyaraman N contributed to conceptualization; Mariappan T and Ramasubramanian S contributed to acquiring clinical data and performing the data analysis; Jeyaraman M, Mariappan T, and Ramasubramanian S contributed to manuscript writing; Jeyaraman M, Santos GS, da Fonseca LF, and Lana JF helped in manuscript revision; Muthu S contributed to image acquisition; Jeyaraman M contributed to proofreading; Jeyaraman M and Lana JF contributed to administration; All authors have agreed to the final version to be published and agree to be accountable for all aspects of the work.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country of origin: India



WJD https://www.wjgnet.com

ORCID number: Madhan Jeyaraman 0000-0002-9045-9493; Tejaswin Mariappan 0009-0007-0837-2334; Naveen Jeyaraman 0000-0002-4362-3326; Sathish Muthu 0000-0002-7143-4354; Swaminathan Ramasubramanian 0000-0001-8845-8427; Gabriel Silva Santos 0000-0002-0549-6821; Lucas Furtado da Fonseca 0000-0001-6497-833X; José Fábio Lana 0000-0002-2330-3982.

S-Editor: Wang JJ L-Editor: Filipodia P-Editor: Chen YX

REFERENCES

- Wang B, Yao M, Lv L, Ling Z, Li L. The Human Microbiota in Health and Disease. Engineering 2017; 3: 71-82 [DOI: 1 10.1016/J.ENG.2017.01.008]
- Barlow GM, Mathur R. Type 2 Diabetes and the Microbiome. J Endocr Soc 2022; 7: bvac184 [PMID: 36578879 DOI: 2 10.1210/jendso/bvac184]
- Dash NR, Al Bataineh MT, Alili R, Al Safar H, Alkhayyal N, Prifti E, Zucker JD, Belda E, Clément K. Functional alterations and predictive 3 capacity of gut microbiome in type 2 diabetes. Sci Rep 2023; 13: 22386 [PMID: 38104165 DOI: 10.1038/s41598-023-49679-w]
- Sadagopan A, Mahmoud A, Begg M, Tarhuni M, Fotso M, Gonzalez NA, Sanivarapu RR, Osman U, Latha Kumar A, Mohammed L. 4 Understanding the Role of the Gut Microbiome in Diabetes and Therapeutics Targeting Leaky Gut: A Systematic Review. Cureus 2023; 15: e41559 [PMID: 37554593 DOI: 10.7759/cureus.41559]
- Iatcu CO, Steen A, Covasa M. Gut Microbiota and Complications of Type-2 Diabetes. Nutrients 2021; 14 [PMID: 35011044 DOI: 5 10.3390/nu14010166
- Zhang D, Jian YP, Zhang YN, Li Y, Gu LT, Sun HH, Liu MD, Zhou HL, Wang YS, Xu ZX. Short-chain fatty acids in diseases. Cell Commun 6 Signal 2023; 21: 212 [PMID: 37596634 DOI: 10.1186/s12964-023-01219-9]
- Birkeland E, Gharagozlian S, Valeur J, Aas AM. Short-chain fatty acids as a link between diet and cardiometabolic risk: a narrative review. 7 Lipids Health Dis 2023; 22: 40 [PMID: 36915164 DOI: 10.1186/s12944-023-01803-5]
- Hernández MAG, Canfora EE, Jocken JWE, Blaak EE. The Short-Chain Fatty Acid Acetate in Body Weight Control and Insulin Sensitivity. 8 Nutrients 2019; 11 [PMID: 31426593 DOI: 10.3390/nu11081943]
- 9 Santos VM, Brito AKP, Amorim AT, Souza IR, Santos MB, Campos GB, Dos Santos DC, Júnior ACRB, Santana JM, Santos DB, Mancini MC, Timenetsky J, Marques LM. Evaluation of fecal microbiota and its correlation with inflammatory, hormonal, and nutritional profiles in women. Braz J Microbiol 2022; 53: 1001-1009 [PMID: 35277849 DOI: 10.1007/s42770-022-00729-x]
- Magne F, Gotteland M, Gauthier L, Zazueta A, Pesoa S, Navarrete P, Balamurugan R. The Firmicutes/Bacteroidetes Ratio: A Relevant Marker 10 of Gut Dysbiosis in Obese Patients? Nutrients 2020; 12 [PMID: 32438689 DOI: 10.3390/nu12051474]
- Stojanov S, Berlec A, Štrukelj B. The Influence of Probiotics on the Firmicutes/Bacteroidetes Ratio in the Treatment of Obesity and 11 Inflammatory Bowel disease. Microorganisms 2020; 8 [PMID: 33139627 DOI: 10.3390/microorganisms8111715]
- Kolmeder CA, de Vos WM. Roadmap to functional characterization of the human intestinal microbiota in its interaction with the host. J 12 Pharm Biomed Anal 2021; 194: 113751 [PMID: 33328144 DOI: 10.1016/j.jpba.2020.113751]
- 13 Hughes ER, Winter MG, Duerkop BA, Spiga L, Furtado de Carvalho T, Zhu W, Gillis CC, Büttner L, Smoot MP, Behrendt CL, Cherry S, Santos RL, Hooper LV, Winter SE. Microbial Respiration and Formate Oxidation as Metabolic Signatures of Inflammation-Associated Dysbiosis. Cell Host Microbe 2017; 21: 208-219 [PMID: 28182951 DOI: 10.1016/j.chom.2017.01.005]
- Crudele L, Gadaleta RM, Cariello M, Moschetta A. Gut microbiota in the pathogenesis and therapeutic approaches of diabetes. EBioMedicine 14 2023; 97: 104821 [PMID: 37804567 DOI: 10.1016/j.ebiom.2023.104821]
- Wong JM, de Souza R, Kendall CW, Emam A, Jenkins DJ. Colonic health: fermentation and short chain fatty acids. J Clin Gastroenterol 15 2006; 40: 235-243 [PMID: 16633129 DOI: 10.1097/00004836-200603000-00015]
- Binda C, Gibiino G, Coluccio C, Sbrancia M, Dajti E, Sinagra E, Capurso G, Sambri V, Cucchetti A, Ercolani G, Fabbri C. Biliary Diseases 16 from the Microbiome Perspective: How Microorganisms Could Change the Approach to Benign and Malignant Diseases. Microorganisms 2022; **10** [PMID: 35208765 DOI: 10.3390/microorganisms10020312]
- Yang Y, Weng W, Peng J, Hong L, Yang L, Toiyama Y, Gao R, Liu M, Yin M, Pan C, Li H, Guo B, Zhu Q, Wei Q, Moyer MP, Wang P, Cai 17 S, Goel A, Qin H, Ma Y. Fusobacterium nucleatum Increases Proliferation of Colorectal Cancer Cells and Tumor Development in Mice by Activating Toll-Like Receptor 4 Signaling to Nuclear Factor-KB, and Up-regulating Expression of MicroRNA-21. Gastroenterology 2017; 152: 851-866.e24 [PMID: 27876571 DOI: 10.1053/j.gastro.2016.11.018]
- Hall AB, Yassour M, Sauk J, Garner A, Jiang X, Arthur T, Lagoudas GK, Vatanen T, Fornelos N, Wilson R, Bertha M, Cohen M, Garber J, 18 Khalili H, Gevers D, Ananthakrishnan AN, Kugathasan S, Lander ES, Blainey P, Vlamakis H, Xavier RJ, Huttenhower C. A novel Ruminococcus gnavus clade enriched in inflammatory bowel disease patients. Genome Med 2017; 9: 103 [PMID: 29183332 DOI: 10.1186/s13073-017-0490-5]
- Falony G, Vlachou A, Verbrugghe K, De Vuyst L. Cross-feeding between Bifidobacterium longum BB536 and acetate-converting, butyrate-19 producing colon bacteria during growth on oligofructose. Appl Environ Microbiol 2006; 72: 7835-7841 [PMID: 17056678 DOI: 10.1128/AEM.01296-06
- 20 Rowland I, Gibson G, Heinken A, Scott K, Swann J, Thiele I, Tuohy K. Gut microbiota functions: metabolism of nutrients and other food components. Eur J Nutr 2018; 57: 1-24 [PMID: 28393285 DOI: 10.1007/s00394-017-1445-8]
- Le TK, Hosaka T, Nguyen TT, Kassu A, Dang TO, Tran HB, Pham TP, Tran QB, Le TH, Pham XD. Bifidobacterium species lower serum 21 glucose, increase expressions of insulin signaling proteins, and improve adipokine profile in diabetic mice. Biomed Res 2015; 36: 63-70 [PMID: 25749152 DOI: 10.2220/biomedres.36.63]
- Moya-Pérez A, Neef A, Sanz Y. Bifidobacterium pseudocatenulatum CECT 7765 Reduces Obesity-Associated Inflammation by Restoring the 22 Lymphocyte-Macrophage Balance and Gut Microbiota Structure in High-Fat Diet-Fed Mice. PLoS One 2015; 10: e0126976 [PMID: 26161548 DOI: 10.1371/journal.pone.0126976]
- 23 Kikuchi K, Ben Othman M, Sakamoto K. Sterilized bifidobacteria suppressed fat accumulation and blood glucose level. Biochem Biophys Res



WJD https://www.wjgnet.com

Commun 2018; 501: 1041-1047 [PMID: 29777696 DOI: 10.1016/j.bbrc.2018.05.105]

- Aoki R, Kamikado K, Suda W, Takii H, Mikami Y, Suganuma N, Hattori M, Koga Y. A proliferative probiotic Bifidobacterium strain in the 24 gut ameliorates progression of metabolic disorders via microbiota modulation and acetate elevation. Sci Rep 2017; 7: 43522 [PMID: 28252037 DOI: 10.1038/srep43522]
- Yang JY, Lee YS, Kim Y, Lee SH, Ryu S, Fukuda S, Hase K, Yang CS, Lim HS, Kim MS, Kim HM, Ahn SH, Kwon BE, Ko HJ, Kweon MN. 25 Gut commensal Bacteroides acidifaciens prevents obesity and improves insulin sensitivity in mice. Mucosal Immunol 2017; 10: 104-116 [PMID: 27118489 DOI: 10.1038/mi.2016.42]
- Wang J, Tang H, Zhang C, Zhao Y, Derrien M, Rocher E, van-Hylckama Vlieg JE, Strissel K, Zhao L, Obin M, Shen J. Modulation of gut 26 microbiota during probiotic-mediated attenuation of metabolic syndrome in high fat diet-fed mice. ISME J 2015; 9: 1-15 [PMID: 24936764 DOI: 10.1038/ismej.2014.99]
- Zhang X, Shen D, Fang Z, Jie Z, Qiu X, Zhang C, Chen Y, Ji L. Human gut microbiota changes reveal the progression of glucose intolerance. 27 PLoS One 2013; 8: e71108 [PMID: 24013136 DOI: 10.1371/journal.pone.0071108]
- 28 Okubo T, Takemura N, Yoshida A, Sonoyama K. KK/Ta Mice Administered Lactobacillus plantarum Strain No. 14 Have Lower Adiposity and Higher Insulin Sensitivity. Biosci Microbiota Food Health 2013; 32: 93-100 [PMID: 24936367 DOI: 10.12938/bmfh.32.93]
- 29 Naito E, Yoshida Y, Makino K, Kounoshi Y, Kunihiro S, Takahashi R, Matsuzaki T, Miyazaki K, Ishikawa F. Beneficial effect of oral administration of Lactobacillus casei strain Shirota on insulin resistance in diet-induced obesity mice. J Appl Microbiol 2011; 110: 650-657 [PMID: 21281408 DOI: 10.1111/j.1365-2672.2010.04922.x]
- Fåk F, Bäckhed F. Lactobacillus reuteri prevents diet-induced obesity, but not atherosclerosis, in a strain dependent fashion in Apoe-/- mice. 30 PLoS One 2012; 7: e46837 [PMID: 23056479 DOI: 10.1371/journal.pone.0046837]
- 31 Moroti C, Souza Magri LF, de Rezende Costa M, Cavallini DC, Sivieri K. Effect of the consumption of a new symbiotic shake on glycemia and cholesterol levels in elderly people with type 2 diabetes mellitus. Lipids Health Dis 2012; 11: 29 [PMID: 22356933 DOI: 10.1186/1476-511X-11-29
- Kijmanawat A, Panburana P, Reutrakul S, Tangshewinsirikul C. Effects of probiotic supplements on insulin resistance in gestational diabetes 32 mellitus: A double-blind randomized controlled trial. J Diabetes Investig 2019; 10: 163-170 [PMID: 29781243 DOI: 10.1111/jdi.12863]
- Tajabadi-Ebrahimi M, Sharifi N, Farrokhian A, Raygan F, Karamali F, Razzaghi R, Taheri S, Asemi Z. A Randomized Controlled Clinical 33 Trial Investigating the Effect of Synbiotic Administration on Markers of Insulin Metabolism and Lipid Profiles in Overweight Type 2 Diabetic Patients with Coronary Heart Disease. Exp Clin Endocrinol Diabetes 2017; 125: 21-27 [PMID: 27219886 DOI: 10.1055/s-0042-105441]
- Asemi Z, Zare Z, Shakeri H, Sabihi SS, Esmaillzadeh A. Effect of multispecies probiotic supplements on metabolic profiles, hs-CRP, and 34 oxidative stress in patients with type 2 diabetes. Ann Nutr Metab 2013; 63: 1-9 [PMID: 23899653 DOI: 10.1159/000349922]
- 35 Mofidi F, Poustchi H, Yari Z, Nourinayyer B, Merat S, Sharafkhah M, Malekzadeh R, Hekmatdoost A. Synbiotic supplementation in lean patients with non-alcoholic fatty liver disease: a pilot, randomised, double-blind, placebo-controlled, clinical trial. Br J Nutr 2017; 117: 662-668 [PMID: 28345499 DOI: 10.1017/S0007114517000204]
- Firouzi S, Majid HA, Ismail A, Kamaruddin NA, Barakatun-Nisak MY. Effect of multi-strain probiotics (multi-strain microbial cell 36 preparation) on glycemic control and other diabetes-related outcomes in people with type 2 diabetes: a randomized controlled trial. Eur J Nutr 2017; 56: 1535-1550 [PMID: 26988693 DOI: 10.1007/s00394-016-1199-8]
- 37 Mohseni S, Bayani M, Bahmani F, Tajabadi-Ebrahimi M, Bayani MA, Jafari P, Asemi Z. The beneficial effects of probiotic administration on wound healing and metabolic status in patients with diabetic foot ulcer: A randomized, double-blind, placebo-controlled trial. Diabetes Metab Res Rev 2018; 34 [PMID: 29193662 DOI: 10.1002/dmrr.2970]
- 38 Gurung M, Li Z, You H, Rodrigues R, Jump DB, Morgun A, Shulzhenko N. Role of gut microbiota in type 2 diabetes pathophysiology. EBioMedicine 2020; 51: 102590 [PMID: 31901868 DOI: 10.1016/j.ebiom.2019.11.051]
- 39 den Besten G, Bleeker A, Gerding A, van Eunen K, Havinga R, van Dijk TH, Oosterveer MH, Jonker JW, Groen AK, Reijngoud DJ, Bakker BM. Short-Chain Fatty Acids Protect Against High-Fat Diet-Induced Obesity via a PPARy-Dependent Switch From Lipogenesis to Fat Oxidation. Diabetes 2015; 64: 2398-2408 [PMID: 25695945 DOI: 10.2337/db14-1213]
- Zhang L, Qin Q, Liu M, Zhang X, He F, Wang G. Akkermansia muciniphila can reduce the damage of gluco/lipotoxicity, oxidative stress and 40 inflammation, and normalize intestine microbiota in streptozotocin-induced diabetic rats. Pathog Dis 2018; 76 [PMID: 29668928 DOI: 10.1093/femspd/fty028]
- Li X, Wang N, Yin B, Fang D, Jiang T, Fang S, Zhao J, Zhang H, Wang G, Chen W. Effects of Lactobacillus plantarum CCFM0236 on 41 hyperglycaemia and insulin resistance in high-fat and streptozotocin-induced type 2 diabetic mice. J Appl Microbiol 2016; 121: 1727-1736 [PMID: 27552342 DOI: 10.1111/jam.13276]
- Ojo O, Feng QQ, Ojo OO, Wang XH. The Role of Dietary Fibre in Modulating Gut Microbiota Dysbiosis in Patients with Type 2 Diabetes: A 42 Systematic Review and Meta-Analysis of Randomised Controlled Trials. Nutrients 2020; 12 [PMID: 33113929 DOI: 10.3390/nu12113239]
- Xi Y, Xu PF. Diabetes and gut microbiota. World J Diabetes 2021; 12: 1693-1703 [PMID: 34754371 DOI: 10.4239/wjd.v12.i10.1693] 43
- Gentile JKA, Oliveira KD, Pereira JG, Tanaka DY, Guidini GN, Cadona MZ, Siriani-Ribeiro DW, Perondini MT. THE INTESTINAL 44 MICROBIOME IN PATIENTS UNDERGOING BARIATRIC SURGERY: A SYSTEMATIC REVIEW. Arq Bras Cir Dig 2022; 35: e1707 [PMID: 36542005 DOI: 10.1590/0102-672020220002e1707]
- Ismail HM, Evans-Molina C. Does the Gut Microbiome Play a Role in Obesity in Type 1 Diabetes? Unanswered Questions and Review of the 45 Literature. Front Cell Infect Microbiol 2022; 12: 892291 [PMID: 35873174 DOI: 10.3389/fcimb.2022.892291]
- Đanić M, Pavlović N, Stanimirov B, Lazarević S, Vukmirović S, Al-Salami H, Mikov M. PAMPA model of gliclazide permeability: The 46 impact of probiotic bacteria and bile acids. Eur J Pharm Sci 2021; 158: 105668 [PMID: 33301903 DOI: 10.1016/j.ejps.2020.105668]
- van Bommel EJM, Herrema H, Davids M, Kramer MHH, Nieuwdorp M, van Raalte DH. Effects of 12-week treatment with dapagliflozin and 47 gliclazide on faecal microbiome: Results of a double-blind randomized trial in patients with type 2 diabetes. Diabetes Metab 2020; 46: 164-168 [PMID: 31816432 DOI: 10.1016/j.diabet.2019.11.005]
- Zhang X, Fang Z, Zhang C, Xia H, Jie Z, Han X, Chen Y, Ji L. Effects of Acarbose on the Gut Microbiota of Prediabetic Patients: A 48 Randomized, Double-blind, Controlled Crossover Trial. Diabetes Ther 2017; 8: 293-307 [PMID: 28130771 DOI: 10.1007/s13300-017-0226-y]
- 49 Madsen MSA, Holm JB, Pallejà A, Wismann P, Fabricius K, Rigbolt K, Mikkelsen M, Sommer M, Jelsing J, Nielsen HB, Vrang N, Hansen HH. Metabolic and gut microbiome changes following GLP-1 or dual GLP-1/GLP-2 receptor agonist treatment in diet-induced obese mice. Sci *Rep* 2019; **9**: 15582 [PMID: 31666597 DOI: 10.1038/s41598-019-52103-x]
- Zhang Q, Xiao X, Zheng J, Li M, Yu M, Ping F, Wang T, Wang X. Featured article: Structure moderation of gut microbiota in liraglutide-50



treated diabetic male rats. Exp Biol Med (Maywood) 2018; 243: 34-44 [PMID: 29171288 DOI: 10.1177/1535370217743765]

- Cefalo CMA, Cinti F, Moffa S, Impronta F, Sorice GP, Mezza T, Pontecorvi A, Giaccari A. Sotagliflozin, the first dual SGLT inhibitor: 51 current outlook and perspectives. Cardiovasc Diabetol 2019; 18: 20 [PMID: 30819210 DOI: 10.1186/s12933-019-0828-y]
- Singh RK, Chang HW, Yan D, Lee KM, Ucmak D, Wong K, Abrouk M, Farahnik B, Nakamura M, Zhu TH, Bhutani T, Liao W. Influence of 52 diet on the gut microbiome and implications for human health. J Transl Med 2017; 15: 73 [PMID: 28388917 DOI: 10.1186/s12967-017-1175-y
- Evans CC, LePard KJ, Kwak JW, Stancukas MC, Laskowski S, Dougherty J, Moulton L, Glawe A, Wang Y, Leone V, Antonopoulos DA, 53 Smith D, Chang EB, Ciancio MJ. Exercise prevents weight gain and alters the gut microbiota in a mouse model of high fat diet-induced obesity. PLoS One 2014; 9: e92193 [PMID: 24670791 DOI: 10.1371/journal.pone.0092193]
- Kerry RG, Patra JK, Gouda S, Park Y, Shin HS, Das G. Benefaction of probiotics for human health: A review. J Food Drug Anal 2018; 26: 54 927-939 [PMID: 29976412 DOI: 10.1016/j.jfda.2018.01.002]
- 55 Monda V, Villano I, Messina A, Valenzano A, Esposito T, Moscatelli F, Viggiano A, Cibelli G, Chieffi S, Monda M, Messina G. Exercise Modifies the Gut Microbiota with Positive Health Effects. Oxid Med Cell Longev 2017; 2017: 3831972 [PMID: 28357027 DOI: 10.1155/2017/3831972]
- Aragón-Vela J, Solis-Urra P, Ruiz-Ojeda FJ, Álvarez-Mercado AI, Olivares-Arancibia J, Plaza-Diaz J. Impact of Exercise on Gut Microbiota 56 in Obesity. Nutrients 2021; 13 [PMID: 34836254 DOI: 10.3390/nu13113999]
- Wegierska AE, Charitos IA, Topi S, Potenza MA, Montagnani M, Santacroce L. The Connection Between Physical Exercise and Gut 57 Microbiota: Implications for Competitive Sports Athletes. Sports Med 2022; 52: 2355-2369 [PMID: 35596883 DOI: 10.1007/s40279-022-01696-x]
- Singh VP, Sharma J, Babu S, Rizwanulla, Singla A. Role of probiotics in health and disease: a review. J Pak Med Assoc 2013; 63: 253-257 58 [PMID: 23894906]
- Tao YW, Gu YL, Mao XQ, Zhang L, Pei YF. Effects of probiotics on type II diabetes mellitus: a meta-analysis. J Transl Med 2020; 18: 30 59 [PMID: 31952517 DOI: 10.1186/s12967-020-02213-2]
- Svensson CK, Cold F, Ribberholt I, Zangenberg M, Mirsepasi-Lauridsen HC, Petersen AM, Helms M. The Efficacy of Faecal Microbiota 60 Transplant and Rectal Bacteriotherapy in Patients with Recurrent Clostridioides difficile Infection: A Retrospective Cohort Study. Cells 2022; 11 [PMID: 36291139 DOI: 10.3390/cells11203272]
- Kobyliak N, Falalyeyeva T, Mykhalchyshyn G, Kyriienko D, Komissarenko I. Effect of alive probiotic on insulin resistance in type 2 diabetes 61 patients: Randomized clinical trial. Diabetes Metab Syndr 2018; 12: 617-624 [PMID: 29661605 DOI: 10.1016/j.dsx.2018.04.015]
- Kumari M, Singh P, Nataraj BH, Kokkiligadda A, Naithani H, Azmal Ali S, Behare PV, Nagpal R. Fostering next-generation probiotics in 62 human gut by targeted dietary modulation: An emerging perspective. Food Res Int 2021; 150: 110716 [PMID: 34865747 DOI: 10.1016/j.foodres.2021.110716]
- Zhao D, Zhu H, Gao F, Qian Z, Mao W, Yin Y, Tan J, Chen D. Antidiabetic effects of selenium-enriched Bifidobacterium longum DD98 in 63 type 2 diabetes model of mice. Food Funct 2020; 11: 6528-6541 [PMID: 32638790 DOI: 10.1039/d0fo00180e]
- Palacios T, Vitetta L, Coulson S, Madigan CD, Lam YY, Manuel R, Briskey D, Hendy C, Kim JN, Ishoey T, Soto-Giron MJ, Schott EM, 64 Toledo G, Caterson ID. Targeting the Intestinal Microbiota to Prevent Type 2 Diabetes and Enhance the Effect of Metformin on Glycaemia: A Randomised Controlled Pilot Study. Nutrients 2020; 12 [PMID: 32660025 DOI: 10.3390/nu12072041]
- 65 Razmpoosh E, Javadi A, Ejtahed HS, Mirmiran P, Javadi M, Yousefinejad A. The effect of probiotic supplementation on glycemic control and lipid profile in patients with type 2 diabetes: A randomized placebo controlled trial. Diabetes Metab Syndr 2019; 13: 175-182 [PMID: 30641692 DOI: 10.1016/j.dsx.2018.08.008]
- Madempudi RS, Ahire JJ, Neelamraju J, Tripathi A, Nanal S. Efficacy of UB0316, a multi-strain probiotic formulation in patients with type 2 66 diabetes mellitus: A double blind, randomized, placebo controlled study. PLoS One 2019; 14: e0225168 [PMID: 31721790 DOI: 10.1371/journal.pone.0225168
- Sabico S, Al-Mashharawi A, Al-Daghri NM, Wani K, Amer OE, Hussain DS, Ahmed Ansari MG, Masoud MS, Alokail MS, McTernan PG. 67 Effects of a 6-month multi-strain probiotics supplementation in endotoxemic, inflammatory and cardiometabolic status of T2DM patients: A randomized, double-blind, placebo-controlled trial. Clin Nutr 2019; 38: 1561-1569 [PMID: 30170781 DOI: 10.1016/j.clnu.2018.08.009]
- Mazruei Arani N, Emam-Djomeh Z, Tavakolipour H, Sharafati-Chaleshtori R, Soleimani A, Asemi Z. The Effects of Probiotic Honey 68 Consumption on Metabolic Status in Patients with Diabetic Nephropathy: a Randomized, Double-Blind, Controlled Trial. Probiotics Antimicrob Proteins 2019; 11: 1195-1201 [PMID: 30218286 DOI: 10.1007/s12602-018-9468-x]
- Kassaian N, Feizi A, Aminorroaya A, Jafari P, Ebrahimi MT, Amini M. The effects of probiotics and synbiotic supplementation on glucose 69 and insulin metabolism in adults with prediabetes: a double-blind randomized clinical trial. Acta Diabetol 2018; 55: 1019-1028 [PMID: 29931423 DOI: 10.1007/s00592-018-1175-2]
- Ebrahimi ZS, Nasli-Esfahani E, Nadjarzade A, Mozaffari-Khosravi H. Effect of symbiotic supplementation on glycemic control, lipid profiles 70 and microalbuminuria in patients with non-obese type 2 diabetes: a randomized, double-blind, clinical trial. J Diabetes Metab Disord 2017; 16: 23 [PMID: 28589103 DOI: 10.1186/s40200-017-0304-8]
- Asemi Z, Alizadeh SA, Ahmad K, Goli M, Esmaillzadeh A. Effects of beta-carotene fortified synbiotic food on metabolic control of patients 71 with type 2 diabetes mellitus: A double-blind randomized cross-over controlled clinical trial. Clin Nutr 2016; 35: 819-825 [PMID: 26209256 DOI: 10.1016/j.clnu.2015.07.009]
- Madjd A, Taylor MA, Mousavi N, Delavari A, Malekzadeh R, Macdonald IA, Farshchi HR. Comparison of the effect of daily consumption of 72 probiotic compared with low-fat conventional yogurt on weight loss in healthy obese women following an energy-restricted diet: a randomized controlled trial. Am J Clin Nutr 2016; 103: 323-329 [PMID: 26702123 DOI: 10.3945/ajcn.115.120170]
- Karamali M, Dadkhah F, Sadrkhanlou M, Jamilian M, Ahmadi S, Tajabadi-Ebrahimi M, Jafari P, Asemi Z. Effects of probiotic 73 supplementation on glycaemic control and lipid profiles in gestational diabetes: A randomized, double-blind, placebo-controlled trial. Diabetes Metab 2016; 42: 234-241 [PMID: 27209439 DOI: 10.1016/j.diabet.2016.04.009]
- Ostadrahimi A, Taghizadeh A, Mobasseri M, Farrin N, Payahoo L, Beyramalipoor Gheshlaghi Z, Vahedjabbari M. Effect of probiotic 74 fermented milk (kefir) on glycemic control and lipid profile in type 2 diabetic patients: a randomized double-blind placebo-controlled clinical trial. Iran J Public Health 2015; 44: 228-237 [PMID: 25905057]
- Eslamparast T, Poustchi H, Zamani F, Sharafkhah M, Malekzadeh R, Hekmatdoost A. Synbiotic supplementation in nonalcoholic fatty liver 75 disease: a randomized, double-blind, placebo-controlled pilot study. Am J Clin Nutr 2014; 99: 535-542 [PMID: 24401715 DOI: 10.3945/ajcn.113.068890]



- Rajkumar H, Mahmood N, Kumar M, Varikuti SR, Challa HR, Myakala SP. Effect of probiotic (VSL#3) and omega-3 on lipid profile, insulin 76 sensitivity, inflammatory markers, and gut colonization in overweight adults: a randomized, controlled trial. Mediators Inflamm 2014; 2014: 348959 [PMID: 24795503 DOI: 10.1155/2014/348959]
- Ivey KL, Hodgson JM, Kerr DA, Lewis JR, Thompson PL, Prince RL. The effects of probiotic bacteria on glycaemic control in overweight 77 men and women: a randomised controlled trial. Eur J Clin Nutr 2014; 68: 447-452 [PMID: 24569536 DOI: 10.1038/ejcn.2013.294]
- Mohamadshahi M, Veissi M, Haidari F, Shahbazian H, Kaydani GA, Mohammadi F. Effects of probiotic yogurt consumption on 78 inflammatory biomarkers in patients with type 2 diabetes. Bioimpacts 2014; 4: 83-88 [PMID: 25035851 DOI: 10.5681/bi.2014.007]
- Asemi Z, Khorrami-Rad A, Alizadeh SA, Shakeri H, Esmaillzadeh A. Effects of synbiotic food consumption on metabolic status of diabetic 79 patients: a double-blind randomized cross-over controlled clinical trial. Clin Nutr 2014; 33: 198-203 [PMID: 23786900 DOI: 10.1016/j.clnu.2013.05.015]
- 80 Mazloom Z, Yousefinejad A, Dabbaghmanesh MH. Effect of probiotics on lipid profile, glycemic control, insulin action, oxidative stress, and inflammatory markers in patients with type 2 diabetes: a clinical trial. Iran J Med Sci 2013; 38: 38-43 [PMID: 23645956]
- Shavakhi A, Minakari M, Firouzian H, Assali R, Hekmatdoost A, Ferns G. Effect of a Probiotic and Metformin on Liver Aminotransferases in 81 Non-alcoholic Steatohepatitis: A Double Blind Randomized Clinical Trial. Int J Prev Med 2013; 4: 531-537 [PMID: 23930163]
- Asemi Z, Samimi M, Tabassi Z, Naghibi Rad M, Rahimi Foroushani A, Khorammian H, Esmaillzadeh A. Effect of daily consumption of 82 probiotic yoghurt on insulin resistance in pregnant women: a randomized controlled trial. Eur J Clin Nutr 2013; 67: 71-74 [PMID: 23187955 DOI: 10.1038/ejcn.2012.189]
- 83 Ejtahed HS, Mohtadi-Nia J, Homayouni-Rad A, Niafar M, Asghari-Jafarabadi M, Mofid V. Probiotic yogurt improves antioxidant status in type 2 diabetic patients. Nutrition 2012; 28: 539-543 [PMID: 22129852 DOI: 10.1016/j.nut.2011.08.013]
- Laitinen K, Poussa T, Isolauri E; Nutrition, Allergy, Mucosal Immunology and Intestinal Microbiota Group. Probiotics and dietary counselling 84 contribute to glucose regulation during and after pregnancy: a randomised controlled trial. Br J Nutr 2009; 101: 1679-1687 [PMID: 19017418 DOI: 10.1017/S0007114508111461]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

