



24766-Review

 Quotes Excluded
 Bibliography Excluded

 21%
 SIMILAR

Match Overview

Match Number	Source	Words	Similarity
1	CrossCheck	60 words	2%
	Xu, Xiao-Ling, Hui-Qin Yu, Wei Hu, Qian Song, and Wei-Min Mao. "A Novel Inflammation-Based Prognostic Score, the		
2	CrossCheck	45 words	2%
	De-shen Wang. "Comparison of the prognostic values of ... arious inflammation based factors in patients with pancreat		
3	CrossCheck	34 words	1%
	Salim, Derya, Hasan Mutlu, Melek Eryilmaz, Ozan Salim, Fatma Musri, Deniz Tural, Şeyda Gİ½ndİ½z, and Hasan		
4	CrossCheck	31 words	1%
	"Abstract", Breast Cancer Research and Treatment, 12/2005		
5	CrossCheck	27 words	1%
	Guthrie, Graeme J.K., Kellie A. Charles, Campbell S.D. Roxburgh, Paul G. Horgan, Donald C. McMillan, and Stephen		
6	CrossCheck	24 words	1%
	"Oral presentations", British Journal of Surgery, 05/2011		

21

Name of Journal: *World Journal of Gastrointestinal Oncology*

ESPS Manuscript NO: 24766

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

5

Prognostic value of inflammation-based markers in patients with pancreatic cancer administered gemcitabine and erlotinib

32

7

Jae Min Lee, Hong Sik Lee, Jong Jin Hyun, Hyuk Soon Choi, Eun Sun Kim, Bora Keum, Yeon Seok Seo, Yoon Tae Jeen, Hoon Jai Chun, Soon Ho Um, Chang Duck Kim

Prognostic value of inflammation-based markers in patients with pancreatic c



全部 图片 视频 新闻 购物 地图 图书

找到约 61,800 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

Google 学术: [Prognostic value of inflammation-based markers in patients with pancreatic cancer administered gemcitabine and erlotinib](#)

[... pancreatic cancer with nab-paclitaxel plus gemcitabine](#) - Von Hoff - 被引用次数: 825

[... inflammation based factors in patients with pancreatic ...](#) - Wang - 被引用次数: 55

[... versus gemcitabine for metastatic pancreatic cancer](#) - Conroy - 被引用次数: 1712

[Prognostic value of clinicopathological characteristics in patients ...](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC4626812/

A total of 103 pancreatic cancer patients were admitted to the Department of ... Six months of gemcitabine-based chemotherapy after surgical resection is the inhibitor of the EGFR receptor pathway erlotinib prolong the median survival time ... Comparison of the prognostic values of various inflammation based factors in ...

[Initial Metastatic Site as a Prognostic Factor in Patients With Stage IV ...](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC4504638/

26 Jun 2015 ... This study aimed to elucidate the prognostic value of the initial The median OS rates of patients who were administered FOLFIRINOX, gemcitabine-based, ... Prior to that, single gemcitabine, gemcitabine plus erlotinib, or gemcitabine plus inflammation-based markers in advanced pancreatic cancer.



[全部](#) [新闻](#) [图片](#) [购物](#) [视频](#) [更多 ▾](#) [搜索工具](#)

找到约 65,800 条结果 (用时 0.61 秒)

Google 学术: Prognostic value of inflammation-based markers in patients with pancreatic cancer administered gemcitabine and erlotinib

... [pancreatic cancer with nab-paclitaxel plus gemcitabine](#) - Von Hoff - 被引用次数: 892

... [inflammation based factors in patients with pancreatic ...](#) - Wang - 被引用次数: 59

... [versus gemcitabine for metastatic pancreatic cancer](#) - Conroy - 被引用次数: 1803

Prognostic value of clinicopathological characteristics in ...

www.ncbi.nlm.nih.gov > ... > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: M Geng - 2015 - [相关文章](#)

A total of 103 [pancreatic cancer patients](#) were admitted to the Department of ... Six months of [gemcitabine](#)-based chemotherapy after surgical resection is the ..., inhibitor of the EGFR receptor pathway [erlotinib](#) prolong the median survival time ... of the [prognostic values of various inflammation based factors in patients with ...](#)

Initial Metastatic Site as a Prognostic Factor in Patients With ...

www.ncbi.nlm.nih.gov > ... > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: HW Kim - 2015 - [相关文章](#)

2015年6月26日 - This study aimed to elucidate the [prognostic value of the initial ...](#) The median OS rates of [patients](#) who were [administered FOLFIRINOX, gemcitabine](#)-based,

[全部](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 66,900 条结果 (用时 0.73 秒)

[Google 学术: Prognostic value of inflammation-based markers in patients with pancreatic cancer administered gemcitabine and erlotinib](#)

... [pancreatic cancer with nab-paclitaxel plus gemcitabine](#) - Von Hoff - 被引用次数: 915

... [inflammation based factors in patients with pancreatic ...](#) - Wang - 被引用次数: 59

... [versus gemcitabine for metastatic pancreatic cancer](#) - Conroy - 被引用次数: 1835

[Prognostic value of clinicopathological characteristics in patie...](#)

[www.ncbi.nlm.nih.gov](#) > ... > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: M Geng - 2015 - [相关文章](#)

A total of 103 [pancreatic cancer patients](#) were admitted to the Department of ... Six months of [gemcitabine](#)-based chemotherapy after surgical resection is ... In the case of elevated preoperative serum CA19.9 levels the assessment of this [marker](#)

Comparison of the [prognostic values](#) of various [inflammation based](#) factors ...

[Initial Metastatic Site as a Prognostic Factor in Patients With ...](#)

[www.ncbi.nlm.nih.gov](#) > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: HW Kim - 2015 - [相关文章](#)

2015年6月26日 - This study aimed to elucidate the [prognostic value](#) of the initial The median OS rates of [patients](#) who were [administered](#) FOLFIRINOX, [gemcitabine](#)-based, ... Prior to that, single [gemcitabine](#), [gemcitabine plus erlotinib](#), or [gemcitabine plus](#)

[inflammation-based markers](#) in advanced [pancreatic cancer](#).

[Prognostic and predictive markers in pancreatic adenocarcino...](#)

[www.sciencedirect.com/science/article/.../S1590865815006908](#) ▾ [翻译此页](#)