



15916-Review
BY TÜNDE HALÁSZ

Quotes Included
Bibliography Included

0%
SIMILAR

Name of journal: World Journal of Gastroenterology
ESPS Manuscript NO: 15916
Columns: ORIGINAL ARTICLE

Retrospective Study
miR-122 negatively correlates with liver fibrosis as detected by histology and transient elastography

Tünde Halász, Gábor Horváth, Gabriella Pár, Klára Werling, András Kiss, Zsuzsa Schaff, Gábor Lendvai

Abstract

Match Overview

There are no matching words to be displayed.

网页 图片 新闻 视频 更多 搜索工具

找到约 9,720 条结果 (用时 0.21 秒)

Google 学术: miR-122 negatively correlated with liver fibrosis as detected by histology and transient elastography

Hepatic and serum levels of miR-122 after chronic HCV ... - Trebicka - 被引用次数: 44

... of hepatocellular carcinoma in non-alcoholic fatty liver ... - Tian - 被引用次数: 11

AbstractBook HEPDART2013 final - Informed Horizons, LLC

<https://www.informedhorizons.com/.../pdf/.../page82.html> - 翻译此页

detectable levels. Moreover, we ... The Expression of miR-122 Negatively Correlates with Severity of Liver Fibrosis Detected by Transient Elastography and Histology. T Halász. 1,2 ... by transient elastography (TE) and METAVIR guideline.

[PDF] Hepatic and serum levels of miR-122 after chronic HCV ...

www.lcgdbzz.org/UpFiles/Article/2012102914946.pdf - 翻译此页

2012年10月29日 - In addition, circulating miR-122 levels correlated negatively with ... using transient elastography can detect severe fibrosis [11, 12], but this technique is ... who underwent liver biopsy and who were characterized by histology,

Unbound MEDLINE | Liver international journal articles from ...

www.unboundmedicine.com/medline/.../Liver_international?... - 翻译此页

A significant correlation was detected between serum and hepatic miR-122 expression ... HBsAg IHC displayed a negative correlation with gamma glutamyl ... Liver histology, profibrogenic genes expression, bacterial DNA translocation, serum ... using transient elastography allows for accurate evaluation of liver fibrosis in ...

Hepatic and serum levels of miR-122 after chronic HCV ...

vadda.icm.edu.pl/.../bwm/meta1/element/elsevier-3e94d0ad-4df7-... - 翻译此页

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

英语网页

 包括专利 包含引用 创建快讯您是不是要找: miR-122 negatively **correlated** with liver fibrosis as detected by histology and transient elastography**Hepatic and serum levels of miR-122 after chronic HCV-induced fibrosis**

journal-of-hepatology.eu 中的 [HTML]

J Trebicka, E Anadol, N Elfimova, I Strack... - Journal of ..., 2013 - Elsevier

... In addition, circulating miR-122 levels **correlated negatively** with increasing stages of **fibrosis**, although ... **122** in the **liver** after chronic **liver** disease, which inversely **correlates** with progression ... Thus, with **fibrosis** progression, **hepatic** levels of **miR-122** decrease and the release in ...

被引用次数: 45 相关文章 所有 7 个版本 引用 保存

miR-33a levels in hepatic and serum after chronic HBV-induced fibrosis

CF Huang, CC Sun, F Zhao, YD Zhang... - Journal of gastroenterology, 2014 - Springer

... significantly **correlates** with progression of **liver fibrosis**. ... with CHB patients in comparison with controls and higher circulating miR-33a levels were **correlated** with the ... multiple target genes by miRNA [36, 61, 62], though Smad7 was previously shown to **negatively** modulate **liver** ...

被引用次数: 1 相关文章 所有 3 个版本 引用 保存

DSS-induced Chronic Colitis Aggravates Inflammation and Fibrogenesis in Mice with CCl4-induced hepatic Fibrosis

EP Plompen, J Schouten, DW Loth, BE Hansen... - Wiley Online Library

... Conclusions: Our digital histopathologic approach **correlates** extremely well with both a pathologist's ... Finally, NK cell cytolytic activity was also **correlated negatively** with **liver fibrosis** scores in ... **Mir-122** expression was assessed in a total of 127 percutaneous **liver** biopsies and 83 ...

引用 保存

REGRESSION OF HEPATIC INFLAMMATION AND FIBROSIS FOLLOWING TRANSPLANTATION OF HUMAN AMNION EPITHELIAL CELLS INTO IMMUNO-

ADSLF IN - Wiley Online Library

... Conclusions: High CCL5 expression **correlates** significantly with a Th1 cytokine signature in PB and ... A higher **correlation** emerged when VirtualTouch **liver** values were multiplied by VirtualTouch spleen ... than 200 individuals with stage III or greater **fibrosis** and **correlated** with the ...

引用 保存

Biomarkers of liver fibrosis.

KT Suk, DY Kim, KM Sohn, DJ Kim - ADVANCES IN CLINICAL ..., 2013 - books.google.com

... Haptoglobin is **negatively** associated with **fibrosis** because its syn- thesis is decreased by hepatocyte ... PIIINP is increased in acute hepatitis and **correlates** with aminotransferases [94, 95]. ... **sis** but does **correlate** with necrosis ...