

World Journal of *Orthopedics*

World J Orthop 2022 November 18; 13(11): 949-1037



EDITORIAL

- 949 Genetics of congenital anomalies of the hand
Kyriazis Z, Kollia P, Grivea I, Sotiriou S, Dailiana ZH

REVIEW

- 955 Changes in trends of orthopedic services due to the COVID-19 pandemic: A review
Obamiro E, Trivedi R, Ahmed N

ORIGINAL ARTICLE**Basic Study**

- 969 Role of the posterior deep deltoid ligament in ankle fracture stability: A biomechanical cadaver study
McCormack DJ, Solan M, Aziz S, Faroug R, Kirmani S, Wright G, Mangwani J

- 978 Anatomic relationship of extensor indicis propius and extensor digitorum communis: Implications for tendon transfer
Zhou J, Frey C, Segovia N, Yao J

Retrospective Cohort Study

- 986 Can immediate postoperative radiographs predict outcomes in pediatric clubfoot?
Leeprakobboon D

Retrospective Study

- 993 Septic arthritis of the hand: From etiopathogenesis to surgical treatment
Lipatov KV, Asatryan A, Melkonyan G, Kazantsev AD, Solov'eva EI, Gorbacheva IV, Vorotyntsev AS, Emelyanov AY

Observational Study

- 1006 Patient preference for trigger finger treatment
Blough C, Najdawi J, Kuschner S

SYSTEMATIC REVIEWS

- 1015 Viral infections in orthopedics: A systematic review and classification proposal
Sidiropoulos K, Christofilos SI, Tsikopoulos K, Kitridis D, Drago L, Meroni G, Romanò CL, Kavarthapu V
- 1029 Patient-reported dissatisfaction following second side in staged bilateral total knee arthroplasty: A systematic review
Gruenberger E, Bae AS, Kelly T, Ponce BA, McGrory J

ABOUT COVER

Editorial Board Member of *World Journal of Orthopedics*, Konstantinos Tsikopoulos, MD, MSc, Senior Researcher, Department of Orthopaedic, Harrogate and District Foundation Trust, Harrogate HG2 7SX, United Kingdom. kostastsikop@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Orthopedics (WJO, World J Orthop)* is to provide scholars and readers from various fields of orthopedics with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJO mainly publishes articles reporting research results and findings obtained in the field of orthopedics and covering a wide range of topics including arthroscopy, bone trauma, bone tumors, hand and foot surgery, joint surgery, orthopedic trauma, osteoarthropathy, osteoporosis, pediatric orthopedics, spinal diseases, spine surgery, and sports medicine.

INDEXING/ABSTRACTING

WJO is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 Journal Citation Indicator (JCI) for *WJO* as 0.62. The *WJO*'s CiteScore for 2021 is 2.4 and Scopus CiteScore rank 2021: Orthopedics and Sports Medicine is 139/284.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*, Production Department Director: *Xiang Li*, Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Orthopedics

ISSN

ISSN 2218-5836 (online)

LAUNCH DATE

November 18, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Massimiliano Leigheb

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/2218-5836/editorialboard.htm>

PUBLICATION DATE

November 18, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Genetics of congenital anomalies of the hand

Zisis Kyriazis, Panagoula Kollia, Ioanna Grivea, Sotirios Sotiriou, Zoe H Dailiana

Specialty type: Orthopedics

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Chisthi MM, India; Kotlyarov S, Russia

Received: August 29, 2022

Peer-review started: August 29, 2022

First decision: September 26, 2022

Revised: October 9, 2022

Accepted: October 14, 2022

Article in press: October 14, 2022

Published online: November 18, 2022



Zisis Kyriazis, Zoe H Dailiana, Department of Orthopaedic Surgery, Faculty of Medicine, School of Health Sciences, University of Thessaly, Larissa 41500, Greece

Panagoula Kollia, Department of Genetics and Biotechnology, Faculty of Biology, National and Kapodistrian University of Athens, Athens 10679, Greece

Ioanna Grivea, Department of Paediatrics, Faculty of Medicine, School of Health Sciences, University of Thessaly, Larissa 41500, Greece

Sotirios Sotiriou, Laboratory of Histology and Embryology, Faculty of Medicine, School of Health Sciences, University of Thessaly, Larissa 41500, Greece

Corresponding author: Zoe H Dailiana, MD, PhD, Professor, Surgeon, Department of Orthopaedic Surgery, Faculty of Medicine, School of Health Sciences, University of Thessaly, No. 3 Panepistimiou Street, Biopolis, Larissa 41500, Greece. dailiana@med.uth.gr

Abstract

Congenital anomalies of the hand are malformations occurring during the development of the human limb, and present as isolated disorders or as a part of a syndrome. During the last years, molecular analysis techniques have offered increasing knowledge about the molecular basis of hand malformations. Disturbances in the signaling pathways during the development of the upper limb result in malformations of the upper extremity. At present, several genes have been identified as responsible for hand anomalies and other have been recognized as suspect genes related to them. Different and new high throughput methods have been introduced for the identification of the gene mutations. In the current editorial, we summarize concisely the current molecular status of isolated hand genetic disorders and the recent progress in molecular genetics, including the genes related to the disorder. This progress improves the knowledge of these disorders and has implications on genetic counselling and prenatal diagnosis.

Key Words: Hand; Gene; Mutation; Molecular; Diagnosis; Disorders

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The genetic basis of hand disorders is elucidated by the expansion of knowledge and introduction of molecular analysis techniques which contribute to the identification of new genes responsible for them. New genes and mutations are being isolated and correlated with the disorder based on the advances in sequencing technology, such as next generation sequencing and genetic consultation, and future therapeutic developments are enhanced. There appears to be a gap in the literature concerning the knowledge about the genetic basis of all hand disorders. The current molecular status of them is discussed and a summary of different genes, already identified or suspected to be related with them, is presented.

Citation: Kyriazis Z, Kollia P, Grivea I, Sotiriou S, Dailiana ZH. Genetics of congenital anomalies of the hand. *World J Orthop* 2022; 13(11): 949-954

URL: <https://www.wjgnet.com/2218-5836/full/v13/i11/949.htm>

DOI: <https://dx.doi.org/10.5312/wjo.v13.i11.949>

INTRODUCTION

Congenital anomalies of the hand very often have an autosomal dominant pattern of inheritance and most of them have a monogenic genetic basis with variable penetrance[1]. They are related with a disturbance of the normal procedure of the development of the limb, with diverse aetiology and variable clinical features, and their proposed classification system has changed throughout the years, incorporating a rather molecular than anatomic scope of the phenotypes[2]. Initially, the Swanson Classification of congenital anomalies of the hand, including nine categories of malformations, was used by the International Federation of Societies for Surgery of the Hand Committee on Congenital Conditions, as it was considered effective according to the knowledge at the time. Later on, as the awareness of pathogenetic routes and molecular basis of limb formation expanded, the object modeling technique classification was presented, dividing hand anomalies into four groups: Malformations (which include the majority of the disorders), deformations, dysplasias, and syndromes[3].

To date, several loci and disease-causing genes, including all four categories of hand disorders, have been identified in humans, and correlated to specific phenotypes.

Since these phenotype manifestations are indicators that the fetus or the newborn may suffer from a syndrome, the ability to identify the potential syndromes associated with these anomalies, is important for the clinician. Additionally, it is important to distinguish between syndromic and non-syndromic cases for reasons of genetic counselling. Therefore, we present a concise summary of the main genes that are responsible for the disorders, whose etiology is mainly based on known genetic and not external factors, and lead to hand disorder phenotypes when mutated.

POLYDACTYLY

Preaxial/Radial polydactyly

Preaxial polydactyly is a malformation described by an extra digit on the radial side of the hand with an incidence as high as 1 in 3000 births. It follows an autosomal dominant inheritance model with reduced penetrance[4]. Thumb polydactyly has been further subdivided into six subtypes by Wassel according to the level of (metacarpal, proximal, or distal phalange) and the extent of duplication (partial and complete)[5,6]. Next generation sequencing (NGS) analysis in a patient with Wassel III polydactyly identified three gene mutations as follows: (1) Substitution c.1639 G>T in *RPGRIP1* gene; (2) Insertion of adenine in *TMEM216* gene; and (3) A>G nucleotide substitution (c.490) in *FBN1* gene. In a patient with Wassel IV duplication, the following mutations were identified: (1) Adenine duplicated in exon 45 of *CEP290* gene; (2) Two substitutions in *RPGRIP1* gene, c.1639 G>T and c.685 G>A; (3) Adenine insertion in *TMEM216* gene, c.432-11 432-10 insA; (4) Substitution G>C c.8249 in *MEGF8* gene; and (5) Substitution T>A c.548 in *CEP164* gene. These mutations are suspected to be involved in the formation of thumb duplication phenotype[7]. Another suspected way of development of preaxial polydactyly is the overexpression of *HES1* gene. The produced protein is considered to intervene in SHH/GLI signaling axis and results in the manifestation of preaxial polydactyly[8]. The disease gene locus with triphalangeal thumb was identified in chromosome 7q36[9]. Point mutations (105C>G, 305A>T, 323T>C, 404G>A, 295T>C, 4909C>T, 297G>A, 334T>G, 402C>T, and 545G>A) have been identified, and a 739A>G transition near the 5-end of the zone of polarizing activity regulatory sequence (*ZRS*) and a 621C>G mutation in the *ZRS* of the *LMBR1* gene have also been mapped[10,11]. Two more novel mutations (*ZRS131A>T* and *ZRS474C>G*) correlated with preaxial polydactyly were identified in a recent study of a Chinese family[12]. No mutations have been identified for index finger polydactyly, which is inherited with an autosomal dominant trait[13].

Postaxial polydactyly

Postaxial polydactyly presents with extra digits on the ulnar side of the hands. Mutations in genes *ZNF141*, *GLI3*, *IQCE*, *GLI1*, *FAM92A1*, *KIAA0825*, and *DACH1* have been isolated and their involvement in this manifestation is identified[14].

Responsible gene loci have been mapped to 7p15-q11.23, 13q21-q32, 19p13.2-p13.1, 7q21-q34, and 13q13.3-13q21.2 regions using genome-wide scan[15]. Subsequently, two heterozygous mutations, p.A765PfsX14 and p.R539TfsX12 in *GLI3* gene, and P.T474I mutation in the *ZNF141* gene have been identified using exome sequencing[16-18]. Recently, a new suspected mutation in *GLI1* gene (c.1133 C>T) was isolated in an Indian family with the disorder and a mutation in *KIAA0825* gene has been isolated and suspected, although the role of the protein encoded by this gene in limb formation is still unclear[19,20].

Central polydactyly and complex types of polydactyly

Central polydactyly phenotype is characterized by duplication of the 2nd, 3rd, or 4th digit[21]. No disease causing locus or gene responsible for central polydactyly has been identified. Mirror image polydactyly is characterized by mirror-image duplication of fingers and toes[22]. A mutation of the *MIPOL1* gene at 14q13 and two heterozygous deletions including the *PITX1* gene were identified[23].

SPLIT-HAND MALFORMATION

Split-hand malformation may occur as an isolated trait or accompanied with other defects. It manifests as a clinically heterogeneous disorder characterized by absent central digital rays, which result in median clefts of the hand. Responsible mutations map to chromosome 7q21.3-q22.1, chromosome Xq26, and chromosome 10q25[24]. *LBX1*, *BTRC*, *POLL*, *FBXW4*, and *BTRC* gene mutations are reported as responsible for the disease[25]. Recent molecular studies have expanded the list of suspected gene mutations. A *TP63* gene translocation and *FGFR1*, *BHLHA9*, *LRP6*, *UBA2*, and *WNT10B* gene mutations have been recently identified[26-30].

RADIAL RAY DEFECTS

Radial ray defects occur as an isolated malformation or syndromic. They are characterized by partial or complete absence of radial ray structures. Radial defects comprise a large group of diseases. They are associated with *TBX3* gene, coding for a T-box transcriptional factor. *TBX3* is widely expressed in a variety of tissues including forelimbs and hindlimbs, epithelium of the mammary gland, the genital tubercle, and the uterus[31].

DEFECTS IN DORSO-VENTRAL PATTERNING

This disorder category involves nail-patella syndrome, which is autosomal dominant and is expressed with defects affecting the nails, skeleton, kidneys, and eyes. Loss of function mutations in the *LMX1B* gene lead to the syndrome[32,33]. *LMX1B* is involved in determination of dorso-ventral patterning of the limb. A mutation of *WIF1* gene has been isolated as a potential novel cause of the phenotype[34].

BRACHYDACTYLY

Brachydactyly phenotype may present as an isolated defect or in association with other malformations and refers to disproportionately short fingers and toes. Isolated brachydactyly usually occur as autosomal dominant traits and show a high degree of phenotypic variability. A locus on chromosome 5p13.3-p13.2 and the Indian hedgehog gene on chromosome 2q35-36 are involved in Type A1 brachydactyly[35]. A mutation in the human bone morphogenetic protein receptor 1B gene (*BMPRI1B*) on chromosome 4q can cause Type A2 brachydactyly. Mutations in growth/differentiation factor-5 gene (*GDF5*) alter the receptor binding affinities and can also cause symphalangism. No gene or locus for Type A3 brachydactyly has been identified. Type B Brachydactyly phenotype involves isolated mutations in the receptor kinase-like orphan receptor 2 gene (*ROR2*) on 9q22[36]. Type C Brachydactyly phenotype is considered to be caused by mutations in *GDF5*[37].

SYNDACTYLY

Syndactyly is characterized by the fusion of soft and/or bony tissue of the fingers of the hand and is the most common congenital malformation of the hand in North America and Europe. Syndactyly can occur as an isolated malformation or as part of a syndrome. *HOXD13*, *FBLN1*, *GJA1*, *LMBR1*, *LRP4*, *GREM*, *FGF16*, and *BHLHA9* genes are incriminated for the disorder, when mutated[38]. Syndactyly type I presents as fusion between the middle and ring fingers. It is an autosomal dominant malformation and the most common type of syndactyly. Mutations in human chromosome 2q34-q36 have been isolated. Syndactyly type II is a dominantly inherited malformation. Its phenotype contains soft tissue syndactyly between the middle and ring fingers and sometimes clinodactyly or camptodactyly of the little finger. *HOXD13* gene mutations are involved in this disorder[39]. Syndactyly type III affects the ring and little fingers with the middle phalanx of the little finger being absent or rudimentary. Mutations in Connexin 43 are involved in this type of syndactyly. Syndactyly type IV manifests as syndactyly of all fingers, and syndactyly type V as metacarpal synostosis. The genetic background of the last two types is not yet investigated[40].

CONCLUSION

Genetic hand disorders and their genetic heterogeneity and allelic heterogeneity between families indicate more complex mechanisms besides simple Mendelian inheritance. These mechanisms include underlying genetic, epigenetic, and environmental factors[41]. With the advent of NGS technology, including exome sequencing and whole-genome sequencing, new mutations causing hand malformations are isolated and the molecular pathogenesis is exposed. Systematic bioinformatics analysis of the responsible genes, using high-throughput sequencing, is a valuable tool in establishing the precise genotype-phenotype correlations of hand genetic disorders. Diagnosis is still largely postbirth, although prenatal diagnosis *via* molecular and genetic methodologies exists. The expansion of our knowledge related to the mutations leading to different phenotypes, with the use of next generation sequencing analysis, will contribute to prenatal diagnosis, prediction of operative treatment strategy, and potential future applications in gene therapy.

FOOTNOTES

Author contributions: Kyriazis Z wrote the paper and participated in the analysis and drafting of the literature data; Kollia P participated in the conception and design of the study, analysis and interpretation of the literature data, and critical revision of the article; Grivea I and Sotiriou S participated in analysis and interpretation of the literature data and critical revision of the article; Dailiana ZH participated in the conception and design of the study, analysis and interpretation of the literature data, and drafting and critical revision of the article; all authors actively reviewed and revised the manuscript and approved the finally submitted manuscript.

Conflict-of-interest statement: All the authors have no conflict of interest related to the manuscript.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Greece

ORCID number: Zisis Kyriazis 0000-0003-2465-8774; Panagoula Kollia 0000-0002-0635-3996; Ioanna Grivea 0000-0002-9221-5628; Sotirios Sotiriou 0000-0002-9466-6086; Zoe H Dailiana 0000-0003-3890-0832.

S-Editor: Wang LL

L-Editor: Wang TQ

P-Editor: Wang LL

REFERENCES

- 1 Talamillo A, Bastida MF, Fernandez-Teran M, Ros MA. The developing limb and the control of the number of digits. *Clin Genet* 2005; **67**: 143-153 [PMID: 15679824 DOI: 10.1111/j.1399-0004.2005.00404.x]
- 2 Kornak U, Mundlos S. Genetic disorders of the skeleton: a developmental approach. *Am J Hum Genet* 2003; **73**: 447-474 [PMID: 12900795 DOI: 10.1086/377110]

- 3 **Tonkin MA**, Oberg KC. The OMT Classification of Congenital Anomalies of the Hand and Upper Limb. *Hand Surg* 2015; **20**: 336-342 [PMID: 26387992 DOI: 10.1142/S0218810415400055]
- 4 **Wessel LE**, Daluiski A, Trehan SK. Polydactyly a review and update of a common congenital hand difference. *Curr Opin Pediatr* 2020; **32**: 120-124 [PMID: 31851054 DOI: 10.1097/MOP.0000000000000871]
- 5 **Manske MC**, Kennedy CD, Huang JI. Classifications in Brief: The Wassel Classification for Radial Polydactyly. *Clin Orthop Relat Res* 2017; **475**: 1740-1746 [PMID: 27613532 DOI: 10.1007/s11999-016-5068-9]
- 6 **Nguyen JL**, Ho CA. Congenital Disorders of the Pediatric Thumb. *JBJS Rev* 2022; **10** [PMID: 35230999 DOI: 10.2106/JBJS.RVW.21.00147]
- 7 **Kyriazis Z**, Kollia P, Grivea I, Varitimidis SE, Constantoulakis P, Dailiana ZH. Thumb duplication: molecular analysis of different clinical types. *Eur J Orthop Surg Traumatol* 2019; **29**: 421-426 [PMID: 30498907 DOI: 10.1007/s00590-018-2343-3]
- 8 **Sharma D**, Mirando AJ, Leinroth A, Long JT, Karner CM, Hilton MJ. HES1 is a novel downstream modifier of the SHH-GLI3 Axis in the development of preaxial polydactyly. *PLoS Genet* 2021; **17**: e1009982 [PMID: 34928956 DOI: 10.1371/journal.pgen.1009982]
- 9 **Heutink P**, Zguricas J, van Oosterhout L, Breedveld GJ, Testers L, Sandkuijl LA, Snijders PJ, Weissenbach J, Lindhout D, Hovius SE. The gene for triphalangeal thumb maps to the subtelomeric region of chromosome 7q. *Nat Genet* 1994; **6**: 287-292 [PMID: 8012392 DOI: 10.1038/ng0394-287]
- 10 **Balci S**, Demirtas M, Civelek B, Piskin M, Sensoz O, Akarsu AN. Phenotypic variability of triphalangeal thumb-polysyndactyly syndrome linked to chromosome 7q36. *Am J Med Genet* 1999; **87**: 399-406 [PMID: 10594878]
- 11 **Kloppocki E**, Ott CE, Benatar N, Ullmann R, Mundlos S, Lehmann K. A microduplication of the long range SHH limb regulator (ZRS) is associated with triphalangeal thumb-polysyndactyly syndrome. *J Med Genet* 2008; **45**: 370-375 [PMID: 18178630 DOI: 10.1136/jmg.2007.055699]
- 12 **Zeng L**, Jin JY, Luo FM, Sheng Y, Wu PF, Xiang R. ZPA Regulatory Sequence Variants in Chinese Patients With Preaxial Polydactyly: Genetic and Clinical Characteristics. *Front Pediatr* 2022; **10**: 797978 [PMID: 35652055 DOI: 10.3389/fped.2022.797978]
- 13 **Atasu M**. Hereditary index finger polydactyly: phenotypic, radiological, dermatoglyphic, and genetic findings in a large family. *J Med Genet* 1976; **13**: 469-476 [PMID: 1018306 DOI: 10.1136/jmg.13.6.469]
- 14 **Ahmad Z**, Liaqat R, Palander O, Bilal M, Zeb S, Ahmad F, Jawad Khan M, Umair M. Genetic overview of postaxial polydactyly: Updated classification. *Clin Genet* 2022 [PMID: 36071556 DOI: 10.1111/cge.14224]
- 15 **Tsai LP**, Liao HM, Chen YJ, Fang JS, Chen CH. A novel microdeletion at chromosome 2q31.1-31.2 in a three-generation family presenting duplication of great toes with clinodactyly. *Clin Genet* 2009; **75**: 449-456 [PMID: 19459884 DOI: 10.1111/j.1399-0004.2008.01147.x]
- 16 **Radhakrishna U**, Blouin JL, Mehenni H, Patel UC, Patel MN, Solanki JV, Antonarakis SE. Mapping one form of autosomal dominant postaxial polydactyly type A to chromosome 7p15-q11.23 by linkage analysis. *Am J Hum Genet* 1997; **60**: 597-604 [PMID: 9042919]
- 17 **Schrauwen I**, Giese AP, Aziz A, Lafont DT, Chakchouk I, Santos-Cortez RLP, Lee K, Acharya A, Khan FS, Ullah A, Nickerson DA, Bamshad MJ, Ali G, Riazuddin S, Ansar M, Ahmad W, Ahmed ZM, Leal SM. FAM92A Underlies Nonsyndromic Postaxial Polydactyly in Humans and an Abnormal Limb and Digit Skeletal Phenotype in Mice. *J Bone Miner Res* 2019; **34**: 375-386 [PMID: 30395363 DOI: 10.1002/jbmr.3594]
- 18 **Umm-e-Kalsoom**, Basit S, Kamran-ul-Hassan Naqvi S, Ansar M, Ahmad W. Genetic mapping of an autosomal recessive postaxial polydactyly type A to chromosome 13q13.3-q21.2 and screening of the candidate genes. *Hum Genet* 2012; **131**: 415-422 [PMID: 21877132 DOI: 10.1007/s00439-011-1085-7]
- 19 **Bakar A**, Ullah A, Bibi N, Khan H, Rahman AU, Ahmad W, Khan B. A novel homozygous variant in the GLI1 underlies postaxial polydactyly in a large consanguineous family with intra familial variable phenotypes. *Eur J Med Genet* 2022; **65**: 104599 [PMID: 36067927 DOI: 10.1016/j.ejmg.2022.104599]
- 20 **Ullah I**, Kakar N, Schrauwen I, Hussain S, Chakchouk I, Liaqat K, Acharya A, Wasif N, Santos-Cortez RLP, Khan S, Aziz A, Lee K, Couthouis J, Horn D, Kragestein BK, Spielmann M, Thiele H, Nickerson DA, Bamshad MJ, Gitler AD, Ahmad J, Ansar M, Borck G, Ahmad W, Leal SM. Variants in KIAA0825 underlie autosomal recessive postaxial polydactyly. *Hum Genet* 2019; **138**: 593-600 [PMID: 30982135 DOI: 10.1007/s00439-019-02000-0]
- 21 **Haber LL**, Adams HB, Thompson GH, Duncan LS, Didomenico LA, McCluskey WP. Unique case of polydactyly and a new classification system. *J Pediatr Orthop* 2007; **27**: 326-328 [PMID: 17414019 DOI: 10.1097/bpo.0b013e3180342ff5]
- 22 **Kloppocki E**, Kähler C, Foulds N, Shah H, Joseph B, Vogel H, Lüttgen S, Bald R, Besoke R, Held K, Mundlos S, Kurth I. Deletions in PITX1 cause a spectrum of lower-limb malformations including mirror-image polydactyly. *Eur J Hum Genet* 2012; **20**: 705-708 [PMID: 22258522 DOI: 10.1038/ejhg.2011.264]
- 23 **Malik S**. Syndactyly: phenotypes, genetics and current classification. *Eur J Hum Genet* 2012; **20**: 817-824 [PMID: 22333904 DOI: 10.1038/ejhg.2012.14]
- 24 **Li QY**, Newbury-Ecob RA, Terrett JA, Wilson DI, Curtis AR, Yi CH, Gebuhr T, Bullen PJ, Robson SC, Strachan T, Bonnet D, Lyonnet S, Young ID, Raeburn JA, Buckler AJ, Law DJ, Brook JD. Holt-Oram syndrome is caused by mutations in TBX5, a member of the Brachyury (T) gene family. *Nat Genet* 1997; **15**: 21-29 [PMID: 8988164 DOI: 10.1038/ng0197-21]
- 25 **Qiu L**, Li C, Zheng G, Yang T, Yang F. Microduplication of BTRC detected in a Chinese family with split hand/foot malformation type 3. *Clin Genet* 2022; **102**: 451-456 [PMID: 35908152 DOI: 10.1111/cge.14204]
- 26 **Peng Y**, Yang S, Xi H, Hu J, Jia Z, Pang J, Liu J, Yu W, Tang C, Wang H. Whole genome sequencing reveals translocation breakpoints disrupting TP63 gene underlying split hand/foot malformation in a Chinese family. *Mol Genet Genomic Med* 2021; **9**: e1604 [PMID: 33471964 DOI: 10.1002/mgg3.1604]
- 27 **Papasozomenou P**, Papoulidis I, Mikos T, Zafrakas M. Split Hand Foot Malformation Syndrome: A Novel Heterozygous FGFR1 Mutation Detected by Next Generation Sequencing. *Curr Genomics* 2019; **20**: 226-230 [PMID: 31929729 DOI: 10.2174/1389202920666190530092856]
- 28 **Caylor R**, Fee T, Lay A, Skinner C, Everman D, Blue E, Bamshad M, Schwartz C, Friez M, Stevenson R. eP326: Genome

- sequencing reveals BHLHA9 gene duplication as cause of multi-generational split-hand/foot malformation with long bone deficiency. *Genet Med* 2022; **24**: S204 [DOI: [10.1016/j.gim.2022.01.361](https://doi.org/10.1016/j.gim.2022.01.361)]
- 29 **Yamoto K**, Saito H, Nishimura G, Kosaki R, Takayama S, Haga N, Tonoki H, Okumura A, Horii E, Okamoto N, Suzumura H, Ikegawa S, Kato F, Fujisawa Y, Nagata E, Takada S, Fukami M, Ogata T. Comprehensive clinical and molecular studies in split-hand/foot malformation: identification of two plausible candidate genes (LRP6 and UBA2). *Eur J Hum Genet* 2019; **27**: 1845-1857 [PMID: [31332306](https://pubmed.ncbi.nlm.nih.gov/31332306/) DOI: [10.1038/s41431-019-0473-7](https://doi.org/10.1038/s41431-019-0473-7)]
- 30 **Kantaputra PN**, Kapoor S, Verma P, Intachai W, Ketudat Cairns JR. Split hand-foot malformation and a novel WNT10B mutation. *Eur J Med Genet* 2018; **61**: 372-375 [PMID: [29427788](https://pubmed.ncbi.nlm.nih.gov/29427788/) DOI: [10.1016/j.ejmg.2018.02.001](https://doi.org/10.1016/j.ejmg.2018.02.001)]
- 31 **Davenport TG**, Jerome-Majewska LA, Papaioannou VE. Mammary gland, limb and yolk sac defects in mice lacking Tbx3, the gene mutated in human ulnar mammary syndrome. *Development* 2003; **130**: 2263-2273 [PMID: [12668638](https://pubmed.ncbi.nlm.nih.gov/12668638/) DOI: [10.1242/dev.00431](https://doi.org/10.1242/dev.00431)]
- 32 **Dreyer S**, Zhou G, Baldini A, Winterpacht A, Zabel B, Cole W, Johnson R, Lee B. Mutations in LMX1 B cause abnormal skeletal patterning and renal dysplasia in nail patella syndrome. *Nat Genet* 1998; **19**: 47-50 [PMID: [9590287](https://pubmed.ncbi.nlm.nih.gov/9590287/) DOI: [10.1038/ng0598-47](https://doi.org/10.1038/ng0598-47)]
- 33 **Lindelöf H**, Horemuzova E, Voss U, Nordgren A, Grigelioniene G, Hammarsjö A. Case Report: Inversion of *LMX1B* - A Novel Cause of Nail-Patella Syndrome in a Swedish Family and a Longtime Follow-Up. *Front Endocrinol (Lausanne)* 2022; **13**: 862908 [PMID: [35769074](https://pubmed.ncbi.nlm.nih.gov/35769074/) DOI: [10.3389/fendo.2022.862908](https://doi.org/10.3389/fendo.2022.862908)]
- 34 **Jones MC**, Topol SE, Rueda M, Oliveira G, Phillips T, Spencer EG, Torkamani A. Mutation of WIF1: a potential novel cause of a Nail-Patella-like disorder. *Genet Med* 2017; **19**: 1179-1183 [PMID: [28383544](https://pubmed.ncbi.nlm.nih.gov/28383544/) DOI: [10.1038/gim.2017.20](https://doi.org/10.1038/gim.2017.20)]
- 35 **Zeng F**, Liu H, Xia X, Shu Y, Cheng W, Xu H, Yin G, Xie Q. Case Report: Brachydactyly Type A1 Induced by a Novel Variant of in-Frame Insertion in the *IHH* Gene. *Front Genet* 2022; **13**: 814786 [PMID: [35669189](https://pubmed.ncbi.nlm.nih.gov/35669189/) DOI: [10.3389/fgene.2022.814786](https://doi.org/10.3389/fgene.2022.814786)]
- 36 **Shao J**, Liu Y, Zhao S, Sun W, Zhan J, Cao L. A novel variant in the ROR2 gene underlying brachydactyly type B: a case report. *BMC Pediatr* 2022; **22**: 528 [PMID: [36064339](https://pubmed.ncbi.nlm.nih.gov/36064339/) DOI: [10.1186/s12887-022-03564-z](https://doi.org/10.1186/s12887-022-03564-z)]
- 37 **Dudley AT**, Ros MA, Tabin CJ. A re-examination of proximodistal patterning during vertebrate limb development. *Nature* 2002; **418**: 539-544 [PMID: [12152081](https://pubmed.ncbi.nlm.nih.gov/12152081/) DOI: [10.1038/nature00945](https://doi.org/10.1038/nature00945)]
- 38 **Cassim A**, Hettiarachchi D, Dissanayake VHW. Genetic determinants of syndactyly: perspectives on pathogenesis and diagnosis. *Orphanet J Rare Dis* 2022; **17**: 198 [PMID: [35549993](https://pubmed.ncbi.nlm.nih.gov/35549993/) DOI: [10.1186/s13023-022-02339-0](https://doi.org/10.1186/s13023-022-02339-0)]
- 39 **Patel R**, Singh SK, Bhattacharya V, Ali A. Novel HOXD13 variants in syndactyly type 1b and type 1c, and a new spectrum of TP63-related disorders. *J Hum Genet* 2022; **67**: 43-49 [PMID: [34321610](https://pubmed.ncbi.nlm.nih.gov/34321610/) DOI: [10.1038/s10038-021-00963-5](https://doi.org/10.1038/s10038-021-00963-5)]
- 40 **Bell J**. On brachydactyly and symphalangism. In: Penrose LS, editor. *The Treasury of Human Inheritance*. Cambridge: Cambridge University Press, 1951: 1-31 [DOI: [10.1007/s12142-000-1047-2](https://doi.org/10.1007/s12142-000-1047-2)]
- 41 **Bosse K**, Betz RC, Lee YA, Wienker TF, Reis A, Kleen H, Propping P, Cichon S, Nöthen MM. Localization of a gene for syndactyly type 1 to chromosome 2q34-q36. *Am J Hum Genet* 2000; **67**: 492-497 [PMID: [10877983](https://pubmed.ncbi.nlm.nih.gov/10877983/) DOI: [10.1086/303028](https://doi.org/10.1086/303028)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

