

Certificate of Biostatistics

29 May 2024

To whom it may concern

Title of Study: GATIS score for predicting the prognosis of rectal neuroendocrine neoplasms: a Chinese multicenter study of twelve-year experience.

After a rigorous evaluation of the manuscript, I think the statistical method of the manuscript is appropriate. I think this manuscript meets the following requirements: (1) Statistical methods are adequately and appropriately described when they are used to verify the results; (2) Statistical techniques are suitable or correct, and compliant with the following Baishideng Publishing Group (Baishideng) directives; (3) Only homogeneous data can be averaged. Standard deviation (SD) is preferred to standard error (SE). The number of observations and subjects (n) is given. Losses in observations, such as drop-outs from the study, are reported; (4) Values, such as ED50, LD50, and IC50, have the 95% confidence limits calculated and have been compared by weighted probit modeling (using the functions described by Bliss and Finney); and (5) The word "significantly" is replaced by its synonyms (if it indicates extent) or the P value (if it indicates statistical significance). Statistical data should be expressed as mean \pm SD or mean \pm SE. Common statistical expressions are identified as: t-test as t ; F-test as F ; chi-square test as χ^2 ; relative coefficient as r ; degree of freedom as df ; number of samples as n ; and probability as P .

Please contact me if you have any queries.

Yong Gan, PhD

Huazhong University of Science and Technology, Wuhan, China

Email: [REDACTED]