

找到约 17,600 条结果 (用时 0.42 秒)

### "No go" donor hepatectomy in living-donor liver transplantation. - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/29170994> - 翻译此页

作者: V Pamecha - 2017 - 相关文章

2017年11月24日 - "No go" donor **hepatectomy in living-donor** liver transplantation. Pamecha ... One patient had unexpected biliary pathology with fibrosis **found** ...

缺少字词: **neuroendocrine tumor incidentally**

### Liver Transplantation For Malignant Tumors - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3229298/> - 翻译此页

作者: B Koneru - 1988 - 被引用次数: 82 - 相关文章

\*One patient was transplanted for metastatic **carcinoid tumors**. A bile duct cancer was **detected in the hepatectomy** specimen. ... the transplant **can** be discontinued and the **donor** liver is transplanted into the "back-up" candidate rather .... The Group I patients who had an **incidental** carcinoma **did** extremely well with a single ....

### Organ transplantation from donors (cadaveric or living) with a history ...

<https://www.sciencedirect.com/science/article/pii/S0955470X1400055X> - 翻译此页

作者: S Zhang - 2014 - 被引用次数: 12 - 相关文章

**Donors** with low-grade skin tumors, carcinomas **in situ** of the uterine cervix, and .... History of melanoma, leukemia or lymphoma, small cell lung/**neuroendocrine carcinoma** .... kidneys from patients with small, localized, **incidentally detected** RCC. .... to the organ, partial **hepatectomy** or retransplantation **can** be **life-saving**.

### Nesimi Mecit's scientific contributions | Sisli Etfal Training and ...

[https://www.researchgate.net/scientific.../2042644623\\_Nesimi\\_Mecit](https://www.researchgate.net/scientific.../2042644623_Nesimi_Mecit) - 翻译此页

Nesimi Mecit's scientific contributions including: Liver Transplant **in** Children ... This report presents

**Name of Journal: World Journal of Hepatology**

**Manuscript NO: 39863**

**Manuscript Type: CASE REPORT**

**Neuroendocrine Tumor Incidentally Detected During Living Donor Hepatectomy:  
Two cases treated with complete response**

Sami Akbulut, Burak Isik, Egemen Cicek, Emine Samdanci, Sezai Yilmaz

#### Abstract

To our best knowledge, no case of a tumor that was incidentally detected during living donor hepatectomy (LDH) has been reported in the English language medical literature. We aimed to present two cases who were incidentally detected to have a grade I neuroendocrine tumor (NET) during our twelve-year LDH experience. First Case: A 29-year-old male underwent right lobe LDH for his father with HBV-related chronic liver disease (CLD). An abdominal exploration immediately prior to the closure of the incision revealed that the appendix vermiformis was edematous and had a firmness with a size of 8-10 mm at its tip. An appendectomy was performed. Second Case: A 26-year-old male underwent LDH for his brother suffering from HBV-related CLD. After right lobe LDH, intestinal length was

#### Match Overview

1	<b>Internet</b> 15 words crawled on 19-Nov-2017 <a href="http://www.uptodate.com">www.uptodate.com</a>	1%
2	<b>Crossref</b> 15 words Sorbye, Halfdan, Jonathan Strosberg, Eric Baudin, David S. Klimstra, and James C. Yao. "Gastroenteropancreatic high ...	1%
3	<b>Crossref</b> 12 words Gokhan Yagci. "Perforation of Meckel's Diverticulum by a C... icken Bone, a Rare Complication: Report of a Case", Surger	1%



[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 5 条结果 (用时 0.40 秒)

### Update on pancreatic neuroendocrine tumors - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4244504/> ▼ [翻译此页](#)作者：LR McKenna - 2014 - 被引用次数：51 - [相关文章](#)

2014年6月3日 - Pancreatic **neuroendocrine tumors** (pNETs) are relatively rare tumors ... Commonly found in the gastrointestinal (GI) tract and lung, they also ..... Partial **hepatectomy** ... orthotopic or **living donor** liver transplantation has been shown to be ... labeled PRRT reported **complete** and partial tumor response in 2% ...

缺少字词：incidentally

### Hepatic surgery for metastases from neuroendocrine tumors. - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/12735141> ▼ [翻译此页](#)作者：JM Sarmiento - 2003 - 被引用次数：224 - [相关文章](#)

In functioning tumors, clinical endocrinopathies are relieved in most patients and this response usually lasts for several months. ... success of the operative approach in such complex cases (54% of patients received a ... in every patient with **carcinoid tumors** and repair the valves prior to **hepatic resection** when indicated [27].

缺少字词：incidentally donor

### Treatment of Neuroendocrine Tumor Liver Metastases - Hindawi

<https://www.hindawi.com/journals/ijh/2012/973946/> ▼ [翻译此页](#)作者：MA Lewis - 2012 - 被引用次数：17 - [相关文章](#)

2012年10月27日 - In the care of patients with hepatic **neuroendocrine** metastases, medical ... 45% were treated with major **hepatectomy**, and 14% required .... 12 (92%) of 13 patients had **complete** ablation of all visible **tumors**, with 2 recurrences at the .... response rates, with reductions in tumor size seen in <10% of cases [65 ...

缺少字词：incidentally

### A Revolution in the Treatment of Neuroendocrine Tumors - Cure Today

<https://www.curetoday.com/...cancer.../a-revolution-in-the-treatment-of-neur...> ▼ [翻译此页](#)



Neuroendocrine tumor incidentally detected during living donor hepatectomy: T



全部 图片 新闻 视频 更多

设置 工具

找到约 5 条结果 (用时 0.54 秒)

### Treatment of Neuroendocrine Tumor Liver Metastases - Hindawi

<https://www.hindawi.com/journals/ijh/2012/973946/> ▼ 翻译此页

作者: MA Lewis - 2012 - 被引用次数: 17 - 相关文章

2012年10月27日 - In the care of patients with hepatic neuroendocrine metastases, medical ... 45% were treated with major hepatectomy, and 14% required .... 12 (92%) of 13 patients had complete ablation of all visible tumors, with 2 recurrences at the .... response rates, with reductions in tumor size seen in <10% of cases [65 ...

缺少字词: incidentally

### Hepatic surgery for metastases from neuroendocrine tumors. - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/12735141> ▼ 翻译此页

作者: JM Sarmiento - 2003 - 被引用次数: 224 - 相关文章

In functioning tumors, clinical endocrinopathies are relieved in most patients and this response usually lasts for several months. ... success of the operative approach in such complex cases (54% of patients received a ... in every patient with carcinoid tumors and repair the valves prior to hepatic resection when indicated [27].

缺少字词: incidentally donor

### Hepatic resection for metastatic gastrointestinal and pancreatic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2225511/> ▼ 翻译此页

作者: D Gomez - 2007 - 被引用次数: 41 - 相关文章

Treatment modalities for hepatic metastases from neuroendocrine tumours (NETs) .... of the primary tumour and detection of hepatic metastases in these six cases was ... Two patients with recurrent disease underwent further hepatic resection .... partial or complete response of hormonally induced symptoms in all cases that ...

缺少字词: incidentally donor

### Multimodal management of neuroendocrine liver metastases

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3028577/> ▼ 翻译此页

作者: A Frilling - 2010 - 被引用次数: 108 - 相关文章

Hepatic resection is the only potentially curative treatment. ... In 12.9% of the patients, metastases are already detectable at the time of initial tumour ... Impact of liver metastases on prognosis of neuroendocrine tumours ..... of living donor liver transplantation for neuroendocrine tumour patients to spare the deceased donor ...

缺少字词: incidentally