



Match Overview

1	Crossref 15 words Wai-Kay Seto, Rex W.H. Hui, Lung-Yi Mak, James Fung et al. "Association Between Hepatic Steatosis, Measured by C	1%
2	Crossref 14 words "Abstracts of the 26th Annual Conference of APASL, February 15-19, 2017, Shanghai, China", Hepatology International	1%
3	Crossref 13 words "Abstracts of the 25th Annual Conference of APASL, February 20-24, 2016, Tokyo, Japan", Hepatology International,	1%

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 41819

Manuscript Type: EDITORIAL

Chronic hepatitis B and metabolic risk factors: A call for rigorous longitudinal studies

Wai-Kay Seto

Abstract

Long-term nucleos(t)ide analogue therapy in chronic hepatitis B virus (HBV) infection is effective in suppressing viral replication and reducing liver-related complications. However, HBV-related liver events can still occur in different patient sub-groups. There

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 690,000 条结果 (用时 0.62 秒)

Google 学术 : Interaction between chronic hepatitis B and metabolic risk factors: emerging evidence

Diabetes in Asia: epidemiology, risk factors, and ... - Chan - 被引用次数 : 1534

Further evidence for an association between non- ... - Caronia - 被引用次数 : 486

Nonalcoholic fatty liver disease and hepatocellular ... - Starley - 被引用次数 : 1032

(PDF) Emerging metabolic risk factors in... - ResearchGate

https://www.researchgate.net/.../321384344_Emerging_metabolic_risk_facto... - 翻译此页

2018年8月1日 - miological evidence showing the significant relationship of these con- ... Interleukin 18; IKK β , inhibitor of nuclear factor kappa-B kinase subunit beta; ... chronic hepatitis patients, even more if both metabolic risk factors were.

Serum from Chronic Hepatitis B Patients Promotes Growth and ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/29763915> - 翻译此页

作者 : Y Ji - 2018 - 相关文章

2018年5月9日 - Emerging evidence implicates insulin-like growth factor (IGF)-II as a major risk factor for the growth and development of HCC. However, the relationship between HBV infection and IGF-II functions remains to be elucidated. ... Hepatitis B, Chronic/complications; Hepatitis B, Chronic/metabolism*; Humans ...

Influence of Metabolic Risk Factors on Risk of Hepatocellular ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/28711626> - 翻译此页

作者 : MW Yu - 2017 - 被引用次数 : 14 - 相关文章

2017年7月12日 - Influence of Metabolic Risk Factors on Risk of Hepatocellular Carcinoma and Liver-Related Death in Men With Chronic Hepatitis B: A Large ...



全部 新闻 图片 视频 更多 设置 工具

找到约 314,000 条结果 (用时 0.46 秒)

Google 学术: Chronic hepatitis B and metabolic risk factors: A call for rigorous longitudinal studies

Hepatitis C infection and risk of diabetes: a systematic ... - White - 被引用次数: 337

Chronic hepatitis B virus infection in Asian countries - Merican - 被引用次数: 586

Chronic hyperlactatemia in HIV-infected patients taking ... - John - 被引用次数: 316

Metabolic risk factors are associated with non - hepatitis B non ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5983169/> - 翻译此页

作者: SF Huang - 2018 - 被引用次数: 1 - 相关文章

2018年4月18日 - **Metabolic risk factors**, such as obesity, fatty liver, high lipidemia, and diabetes mellitus are ... Of these patients, 411 (10.69%) who were negative for **hepatitis B virus (HBV)**, surface ... These viral marker **data** were used to sort the 3,843 patients with HCC into four Policies and Guidelines | **Contact**.

缺少字词: rigorous longitudinal

Influence of Metabolic Risk Factors on Risk of Hepatocellular ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/28711626> - 翻译此页

作者: MW Yu - 2017 - 被引用次数: 17 - 相关文章

2017年7月12日 - (1)Graduate Institute of Epidemiology and Preventive Medicine, College of ... We used patients' baseline **data** on obesity, diabetes, hypertriglyceridemia, and ... **HBV** carriers with different **metabolic risk factors** had significant ...

缺少字词: call rigorous longitudinal

找到约 393,000 条结果 (用时 0.54 秒)

Google 学术: Chronic hepatitis B and metabolic risk factors: A call for rigorous longitudinal studies

Hepatitis C infection and risk of diabetes: a systematic ... - White - 被引用次数: 337

Chronic hepatitis B virus infection in Asian countries - Merican - 被引用次数: 586

Chronic hyperlactatemia in HIV-infected patients taking ... - John - 被引用次数: 316

Metabolic risk factors are associated with non - hepatitis B non ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5983169/> - 翻译此页

作者: SF Huang - 2018 - 被引用次数: 1 - 相关文章

2018年4月18日 - Metabolic risk factors, such as obesity, fatty liver, high lipidemia, and diabetes mellitus are ... Of these patients, 411 (10.69%) who were negative for hepatitis B virus (HBV), surface ... These viral marker data were used to sort the 3,843 patients with HCC into four Policies and Guidelines | Contact.

缺少字词: rigorous longitudinal

Influence of Metabolic Risk Factors on Risk of Hepatocellular ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/28711626> - 翻译此页

作者: MW Yu - 2017 - 被引用次数: 17 - 相关文章

2017年7月12日 - (1)Graduate Institute of Epidemiology and Preventive Medicine, College of ... We used patients' baseline data on obesity, diabetes, hypertriglyceridemia, and ... HBV carriers with different metabolic risk factors had significant ...

缺少字词: call rigorous longitudinal

Hepatocellular carcinoma among US and non-US-born patients with ...

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone...> - 翻译此页

作者: K Kennedy - 2018

2018年9月25日 - Background Risk factors for hepatocellular carcinoma (HCC) have not been well ... We identified a total of 278 patients with HCC and chronic HBV seen at two and other longitudinal cohort studies of patients with chronic HBV that have ... Diabetes and metabolic syndrome have both been shown to be ...