

Supplementary Table 1 Effect on endoscopic resection for gastric cancer

	Univariate analysis		Multivariate analysis		
	Odds ratio	<i>P</i> value	Regression coefficient	Odds ratio (95% confidence interval)	<i>P</i> value
Age	1.04	<.001	-0.013	0.99 (0.95-1.02)	.48
Male sex	1.23	.43			
Body mass index	1.22	<.001	0.212	1.24 (1.08-1.41)	.002
Surveillance endoscopy within 2 years	10.26	<.001	2.713	15.08 (3.91-58.11)	<.001
Current <i>H. pylori</i> infection	0.22	<.001	0.254	1.29 (0.30-5.56)	.73
Endoscopic findings					
Atrophy score	0.77	.30			
Intestinal metaplasia score	1.39	.03	0.004	1.00 (0.59-1.70)	.99
Enlarged folds score	0.18	<.001	-1.303	0.27 (0.11-0.67)	.004
Nodularity score	0.24	.003	-0.355	0.70 (0.15-3.36)	.66
Diffuse redness score	0.51	<.001	-0.223	0.80 (0.38-1.67)	.55
Kyoto score	0.77	.001			
Lauren's intestinal-type	28.06	<.001	3.911	49.96 (12.81-194.77)	<.001

P values were calculated using binomial logistic regression analysis.