

Supplemental Table 1. Multivariable Cox regression model for 1-year mortality

Variable	HR	95% CI	p-value
Male sex	1.56	1.15–2.10	0.004
Peripheral vascular disease	0.97	0.66–1.40	0.89
MELD-XI	1.03	1.00–1.06	0.036
LVAD	1.27	0.97–1.60	0.087
ECMO at transplant	2.66	1.75–4.00	<0.001
Infection	1.27	0.91–1.70	0.16
KDPI	1.49	0.68–3.20	0.32

HR = hazard ratio; CI = confidence interval.

Supplemental Table 2 Multivariable Cox regression model for 90-day kidney graft failure

Variable	HR	95% CI	p-value
Male sex	1.7	1.25–2.30	<0.001
Peripheral vascular disease	0.88	0.59–1.30	0.53
Kidney pump use	—	—	>0.99
Cold ischemia time	1.02	1.00–1.04	0.028
MELD-XI	1.06	1.03–1.09	<0.001
LVAD	1.87	1.43–2.40	<0.001
ECMO at transplant	2.25	1.43–3.50	<0.001
Infection	1.15	0.82–1.60	0.41
KDPI	2.17	1.00–4.70	0.051

HR = hazard ratio; CI = confidence interval.

Supplemental Table 3 Multivariable Cox regression model for 1-year kidney graft failure

Characteristic	HR	95% CI	p-value
Male sex	0.64	0.49–0.84	0.001
Peripheral vascular disease	0.85	0.60–1.21	0.36

Donor age	1.02	1.01–1.03	0.002
Donor BMI	1.02	1.01–1.04	0.01
Cold ischemia time	1.02	1.01–1.03	0.005
MELD-XI	1.05	1.03–1.07	<0.001
LVAD	1.53	1.21–1.93	<0.001
ECMO at transplant	2.19	1.48–3.23	<0.001
Infection	1.35	1.02–1.79	0.037

HR = hazard ratio; CI = confidence interval.

Supplemental Table 4 Multivariable Cox regression model for 1-year kidney graft failure using MELD-XI tertiles

Characteristic	HR	95% CI	p-value
Male sex	0.63	0.48–0.83	<0.001
Peripheral vascular disease	0.84	0.59–1.20	0.34
Donor age	1.02	1.01–1.03	0.002
Donor BMI	1.02	1.01–1.04	0.011
Cold ischemia time	1.02	1.01–1.03	0.004
MELD-XI tertile (reference = low)			
Medium	1.53	1.18–1.99	0.001
High	1.81	1.25–2.62	0.002
LVAD	1.53	1.21–1.93	<0.001
ECMO at transplant	2.27	1.53–3.36	<0.001
Infection	1.36	1.02–1.80	0.034

HR = hazard ratio; CI = confidence interval.