

# World Journal of *Hepatology*

*World J Hepatol* 2024 June 27; 16(6): 863-972



**EDITORIAL**

- 863 From non-alcoholic fatty liver disease to metabolic-associated steatotic liver disease: Rationale and implications for the new terminology  
*Malnick SDH, Zamir D*
- 867 Immunological crossroads: The intriguing dance between hepatitis C and autoimmune hepatitis  
*Soldera J*
- 871 Sarcopenia and metabolic dysfunction associated steatotic liver disease: Time to address both  
*Wong R, Yuan LY*
- 878 Importance of the gut microbiota in the gut-liver axis in normal and liver disease  
*Kotlyarov S*
- 883 Cold ischemia time in liver transplantation: An overview  
*Cesaretti M, Izzo A, Pellegrino RA, Galli A, Mavrothalassitis O*
- 891 Milestones to optimize of transjugular intrahepatic portosystemic shunt technique as a method for the treatment of portal hypertension complications  
*Garbuzenko DV*

**MINIREVIEWS**

- 900 Hepatitis B cure: Current situation and prospects  
*Li YP, Liu CR, He L, Dang SS*

**ORIGINAL ARTICLE****Observational Study**

- 912 In-hospital outcomes in COVID-19 patients with non-alcoholic fatty liver disease by severity of obesity: Insights from national inpatient sample 2020  
*Srikanth S, Garg V, Subramanian L, Verma J, Sharma H, Klair HS, Kavathia SA, Teja JK, Vasireddy NS, Anmol K, Kolli D, Bodhankar SS, Hashmi S, Chauhan S, Desai R*

- 920 Liver histological changes in untreated chronic hepatitis B patients in indeterminate phase  
*Huang DL, Cai QX, Zhou GD, Yu H, Zhu ZB, Peng JH, Chen J*

**Basic Study**

- 932 Diagnostic and prognostic role of LINC01767 in hepatocellular carcinoma  
*Zhang L, Cui TX, Li XZ, Liu C, Wang WQ*

**SCIENTOMETRICS**

- 951** Mapping the global research landscape on nonalcoholic fatty liver disease and insulin resistance: A visualization and bibliometric study

*Zyoud SH, Hegazi OE, Alalalmeh SO, Shakhshir M, Abushamma F, Khilfeh S, Al-Jabi SW*

**CASE REPORT**

- 966** Successful treatment of severe hepatic impairment in erythropoietic protoporphyria: A case report and review of literature

*Zeng T, Chen SR, Liu HQ, Chong YT, Li XH*

**ABOUT COVER**

Editorial Board Member of *World Journal of Hepatology*, Chien-Hao Huang, MD, PhD, Associate Professor, Department of Gastroenterology and Hepatology, Chang-Gung Memorial Hospital, Linkou Medical Center, Taoyuan, Taiwan, TaoYuan 330, Taiwan. huangchianhou@gmail.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Hepatology (WJH, World J Hepatol)* is to provide scholars and readers from various fields of hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

*WJH* mainly publishes articles reporting research results and findings obtained in the field of hepatology and covering a wide range of topics including chronic cholestatic liver diseases, cirrhosis and its complications, clinical alcoholic liver disease, drug induced liver disease autoimmune, fatty liver disease, genetic and pediatric liver diseases, hepatocellular carcinoma, hepatic stellate cells and fibrosis, liver immunology, liver regeneration, hepatic surgery, liver transplantation, biliary tract pathophysiology, non-invasive markers of liver fibrosis, viral hepatitis.

**INDEXING/ABSTRACTING**

The *WJH* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (ESCI), Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJH* as 2.4. The *WJH*'s CiteScore for 2023 is 4.1 and Scopus CiteScore rank 2023: Hepatology is 41/82.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Yi-Xuan Cai, Production Department Director: Xiang Li, Cover Editor: Xiang Li.

**NAME OF JOURNAL**

*World Journal of Hepatology*

**ISSN**

ISSN 1948-5182 (online)

**LAUNCH DATE**

October 31, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Nikolaos Pyrsopoulos, Ke-Qin Hu, Koo Jeong Kang

**EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF**

Shuang-Suo Dang

**EDITORIAL BOARD MEMBERS**

<https://www.wjnet.com/1948-5182/editorialboard.htm>

**PUBLICATION DATE**

June 27, 2024

**COPYRIGHT**

© 2024 Baishideng Publishing Group Inc

**PUBLISHING PARTNER**

Department of Infectious Diseases, the Second Affiliated Hospital of Xi'an Jiaotong University

**INSTRUCTIONS TO AUTHORS**

<https://www.wjnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjnet.com/bpg/gerinfo/208>

**POLICY OF CO-AUTHORS**

<https://www.wjnet.com/bpg/GerInfo/310>

**ARTICLE PROCESSING CHARGE**

<https://www.wjnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

**PUBLISHING PARTNER'S OFFICIAL WEBSITE**

[http://2yuan.xjtu.edu.cn/Html/Departments/Main/Index\\_21148.html](http://2yuan.xjtu.edu.cn/Html/Departments/Main/Index_21148.html)

## From non-alcoholic fatty liver disease to metabolic-associated steatotic liver disease: Rationale and implications for the new terminology

Stephen David Howard Malnick, Doron Zamir

**Specialty type:** Gastroenterology and hepatology

**Provenance and peer review:** Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's classification**

**Scientific Quality:** Grade B, Grade D

**Novelty:** Grade B, Grade C

**Creativity or Innovation:** Grade B, Grade C

**Scientific Significance:** Grade B, Grade C

**P-Reviewer:** Peltec A, Moldova; Yao SK, China

**Received:** December 7, 2023

**Revised:** February 12, 2024

**Accepted:** April 25, 2024

**Published online:** June 27, 2024



**Stephen David Howard Malnick**, Department of Internal Medicine C, Kaplan Medical Center, Rehovot 76100, Israel

**Doron Zamir**, Department of Internal Medicine D, Barzilai Medical Center, Ashkelon 7830604, Israel

**Corresponding author:** Stephen David Howard Malnick, AGAF, MA, MBBS, MSc, Adjunct Associate Professor, Professor, Department of Internal Medicine C, Kaplan Medical Center, Rehov Pasternak, Rehovot 76100, Israel. [stephen@malnick.net](mailto:stephen@malnick.net)

### Abstract

Non-alcoholic fatty liver disease (NAFLD) was the term first used to describe hepatic steatosis in patients with the metabolic syndrome who did not consume excess amounts of alcohol. Alcoholic liver disease (ALD) has many similarities to NAFLD in both pathogenesis and histology. This entity is now the most prevalent chronic liver disease worldwide as a consequence of the epidemic of obesity. Attempts to incorporate the importance of the metabolic syndrome in the development of steatosis resulted in the renaming of NAFLD as metabolic-associated fatty liver disease. This new term, however, has the disadvantage of the use of terms that may be perceived as derogatory. The terms fatty and non-alcoholic have negative connotations in many cultures. In addition, non-alcoholic is not usually a term applicable to pediatric cases of hepatic steatosis. Recently, an international collaborative effort, with participants from 56 countries, after a global consultation process, recommended to change the nomenclature to steatotic liver disease -including metabolic dysfunction- associated steatotic liver disease, metabolic-associated steatohepatitis and metabolic dysfunction-associated ALD. The new terminology is consistent with most of the previously published epidemiological studies and will have a major impact on research into diagnosis, prognosis and treatment.

**Key Words:** Non-alcoholic fatty liver disease; Steatosis metabolic-associated steatotic liver disease; Nomenclature

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Due to the epidemic of obesity, there has been an increase in the prevalence of the metabolic syndrome and hepatic steatosis. The term non-alcoholic fatty liver disease (NAFLD) was given to describe the hepatic manifestation of the metabolic syndrome. Recently the nomenclature has been changed to metabolic dysfunction-associated steatotic liver disease. This removes stigmata associated with the term fatty and include a new entity reflecting both the metabolic syndrome and alcohol as causes of steatosis. These new terms do not alter the inclusion criteria for most of the published studies on NAFLD and will facilitate future studies.

**Citation:** Malnick SDH, Zamir D. From non-alcoholic fatty liver disease to metabolic-associated steatotic liver disease: Rationale and implications for the new terminology. *World J Hepatol* 2024; 16(6): 863-866

**URL:** <https://www.wjgnet.com/1948-5182/full/v16/i6/863.htm>

**DOI:** <https://dx.doi.org/10.4254/wjh.v16.i6.863>

## INTRODUCTION

The excess consumption of alcohol and food was first described in biblical times. Noah, after leaving the ark at the end of the flood, planted a vineyard, consumed too much of the wine he produced and lay naked in his tent (Genesis 9: 21-27). The two daughters of Lot, fearing that there were no men left in the world to provide them with children, made their father inebriated and then became pregnant by him (Genesis 19: 30-38). The Bible also warns against the excess consumption of food and alcohol, "Do not join those who drink too much wine or gorge themselves on meat (Proverbs 23: 19-21). Our failure to comply with these biblical words of advice, has resulted in an epidemic of obesity and alcohol consumption worldwide. As a consequence of this there is now a large increase in obesity and the associated metabolic syndrome resulting in an increase in hepatic steatosis.

## STEATOTIC LIVER DISEASE

Steatosis of the liver, or fatty liver, with macrovesicular steatosis and Mallory bodies was described by Addison in 1836 [1]. Connor [2] and Chaikoff *et al* [3] found a link between steatosis and either alcohol consumption or diabetes in 1938. In the 1980s Ludwig *et al* [4] described patients who had a similar feature but denied alcohol consumption. The initial nomenclature was non-alcoholic fatty liver disease (NAFLD). The metabolic syndrome is based on the presence of hypertension, impaired glucose tolerance, elevated triglyceride levels, low high-density lipoprotein and increased abdominal circumference, with some differences between different ethnic groups [5].

Subsequently, NAFLD was noted to be the hepatic manifestation of the metabolic syndrome [6]. NAFLD is now the major cause of chronic liver disease worldwide and one of the main indications for liver transplantation [7]. The definition of non-alcoholic is 30 G per day for a male and 20 g per day for a female. However, many patients who are considered to have NAFLD do in fact have significant alcohol consumption [8].

The pathogenesis of steatotic liver disease involves the development of a dysbiosis, resulting in endogenous alcohol production, increased intestinal permeability, the production of endotoxins, which reach the liver by way of the hepatic portal vein. The insulin resistance that is associated with the metabolic syndrome results in an increase in intrahepatic fat, increased gluconeogenesis and increased free fatty acid levels [9]. In both alcoholic and non-alcoholic steatosis the liver injury is in the parenchyma. Progressive liver injury results in an evolution from simple steatosis, to steatohepatitis, fibrosis and cirrhosis. Lifestyle modifications can decrease and even reverse the disease progression [9].

As a result of this a new nomenclature was proposed: metabolic dysfunction-associated fatty liver disease (MAFLD) [10]. This term was felt to reflect the importance of metabolic factors in the etiology of MAFLD and to assist in providing a better understanding of this entity by patients.

There are, however, problems with this nomenclature as well. For example, not all cases of MAFLD are seen in patients with obesity. This is reflected in the term lean NAFLD [11]. In many of these patients a history of drinking sweet sugary beverages or soda with artificial sweeteners may be present. Some patients with cirrhotic liver disease suspected to be caused by MAFLD may not have fat in the liver when assessed by biopsy. The change to MAFLD has been suggested to have been premature [12].

Treatment for MAFLD is mainly based on lifestyle changes. Clinical trials with anti-obesity treatments, lipid-lowering agents and insulin sensitizers in the main were not successful. There is concern that concentrating just on metabolic factors could impede the development of other therapeutic mechanisms [13]. In addition, in many trials the end-point selected was a decrease in the NASH score and not in the more important degree of fibrosis or clinical end-points including cardiovascular events or death [14]. Other factors affecting the development of steatosis include the intestinal microbiome, genetic factors including PNPLA3 mutations [15], and sarcopenia.

Recently, the name has been revised once more. A global consultation process using the well-established Delphi technique was performed from 2020 to 2023 [16-18]. This involved 236 participants from 56 countries. The terms non-alcoholic and fatty were rejected for use since they were associated with a stigma in terms of patient understanding. Steatosis was chosen to replace "fatty". The new nomenclature to be recommended was metabolic dysfunction-associated steatotic liver disease (MASLD).

The presence of at least one of the five cardiometabolic risk factors is essential to make the diagnosis. The previous diagnosis of non-alcoholic steatohepatitis (NASH) was recommended to be replaced by metabolic dysfunction associated steatohepatitis (MASH). Cryptogenic steatotic liver disease was coined to cover those cases with no clear cause.

Metabolic dysfunction-associated ALD (MetALD) is a new term to describe those cases with excessive alcohol consumption. This entity has not been previously defined and will now have to be investigated further in appropriate clinical trials.

---

## CONSEQUENCES OF THE CHANGE IN NOMENCLATURE

---

The change in nomenclature preserves the existing data on natural history of the disease, biomarkers and clinical trials, which is very important. An analysis of the European NAFLD registry cohort has found that 98% of the participants would fulfill the new criteria for MASLD[19]. In addition, a study of 1333 patients from Hong Kong with NAFLD found that only 4 (0.3%) of patients did not meet the new MASLD criteria[20].

The diagnosis of MASLD has an advantage of removing the stigma associated with the term fatty, and patient advocacy groups were consulted as part of the process. The importance of this effect differs between different cultures [21].

Steatotic liver disease reflects the range of causes of hepatic steatosis and permits characterization of fibrotic severity – for example MASH with stage 3 fibrosis. It is to be noted that disease stage and severity are not changed in this new definition and will still be valid with the increasing use of non-invasive testing for determining these parameters.

There is a strong similarity between the metabolic criteria for diagnosing MASLD and those previously suggested for MAFLD. The new definition of MASLD is based on simple and accessible clinical criteria and biological measurements without the need for specialized testing for insulin resistance such as homeostasis model assessment of insulin resistance. There may be a place for subsequent testing in the minority of patients with hepatic steatosis in the absence of cardiometabolic risk factors.

The new classification also enables subgroups to be included within the umbrella term steatotic liver disease. This includes medications and rare genetic diseases in children. The newly defined term MetALD reflects the importance of alcohol consumption in the development of steatosis, even in the absence of metabolic factors.

---

## CONCLUSION

---

This new nomenclature for the most common liver disease worldwide is important for developing a cohesive definition between groups worldwide which will enable extensive data collection with implications for the development of personalized medical therapeutic and enabling incorporation of existing data sets. The public health implications of this are likely to be very significant.

---

## FOOTNOTES

---

**Author contributions:** Malnick SDH and Zamir D contributed equally to the conception, writing and editing of this manuscript.

**Conflict-of-interest statement:** There are no conflicts of interest.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** Israel

**ORCID number:** Stephen David Howard Malnick [0000-0003-1865-8313](https://orcid.org/0000-0003-1865-8313).

**Corresponding Author's Membership in Professional Societies:** American Gastroenterological Association, C--175497.

**S-Editor:** Zhang H

**L-Editor:** A

**P-Editor:** Cai YX

---

## REFERENCES

---

- 1 Addison T. Observations on fatty degeneration of the liver. *Guys Hosp Rep* 1836; 1: 476
- 2 Connor CL. Fatty infiltration of the liver and the development of cirrhosis in diabetes and chronic alcoholism. *Am J Pathol* 1938; 14: 347-364



- 3 **Chaikoff IL**, Connor CL, Biskind GR. Fatty infiltration and cirrhosis of the liver in depancreatized dogs maintained with insulin. *Am J Pathol* 1938; **14**: 101-110
- 4 **Ludwig J**, Viggiano TR, McGill DB, Oh BJ. Nonalcoholic steatohepatitis: Mayo Clinic experiences with a hitherto unnamed disease. *Mayo Clin Proc* 1980; **55**: 434-438
- 5 **Lorenzo C**, Williams K, Hunt KJ, Haffner SM. The National Cholesterol Education Program - Adult Treatment Panel III, International Diabetes Federation, and World Health Organization definitions of the metabolic syndrome as predictors of incident cardiovascular disease and diabetes. *Diabetes Care* 2007; **30**: 8-13 [PMID: [17192325](#) DOI: [10.2337/dc06-1414](#)]
- 6 **Kim CH**, Younossi ZM. Nonalcoholic fatty liver disease: a manifestation of the metabolic syndrome. *Cleve Clin J Med* 2008; **75**: 721-728 [PMID: [18939388](#) DOI: [10.3949/ccjm.75.10.721](#)]
- 7 **Riazi K**, Azhari H, Charette JH, Underwood FE, King JA, Afshar EE, Swain MG, Congly SE, Kaplan GG, Shaheen AA. The prevalence and incidence of NAFLD worldwide: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol* 2022; **7**: 851-861 [PMID: [35798021](#) DOI: [10.1016/S2468-1253\(22\)00165-0](#)]
- 8 **Staufer K**, Huber-Schönauer U, Strebinger G, Pimingsstorfer P, Suesse S, Scherzer TM, Paulweber B, Ferenci P, Stimpfl T, Yegles M, Datz C, Trauner M. Ethyl glucuronide in hair detects a high rate of harmful alcohol consumption in presumed non-alcoholic fatty liver disease. *J Hepatol* 2022; **77**: 918-930 [PMID: [35605744](#) DOI: [10.1016/j.jhep.2022.04.040](#)]
- 9 **Malnick SDH**, Alin P, Somin M, Neuman MG. Fatty Liver Disease-Alcoholic and Non-Alcoholic: Similar but Different. *Int J Mol Sci* 2022; **23** [PMID: [36555867](#) DOI: [10.3390/ijms232416226](#)]
- 10 **Eslam M**, George J. Reply to: correspondence regarding "A new definition for metabolic dysfunction-associated fatty liver disease: An international expert consensus statement": Bringing evidence to the NAFLD-MAFLD debate. *J Hepatol* 2020; **73**: 1575 [PMID: [32933781](#) DOI: [10.1016/j.jhep.2020.07.045](#)]
- 11 **Xu R**, Pan J, Zhou W, Ji G, Dang Y. Recent advances in lean NAFLD. *Biomed Pharmacother* 2022; **153**: 113331 [PMID: [35779422](#) DOI: [10.1016/j.biopha.2022.113331](#)]
- 12 **Younossi ZM**, Rinella ME, Sanyal AJ, Harrison SA, Brunt EM, Goodman Z, Cohen DE, Loomba R. From NAFLD to MAFLD: Implications of a Premature Change in Terminology. *Hepatology* 2021; **73**: 1194-1198 [PMID: [32544255](#) DOI: [10.1002/hep.31420](#)]
- 13 **Zhang C**, Yang M. Current Options and Future Directions for NAFLD and NASH Treatment. *Int J Mol Sci* 2021; **22** [PMID: [34299189](#) DOI: [10.3390/ijms22147571](#)]
- 14 **Malnick S**, Somin M. Clinical endpoints are necessary in the interim analysis of REGENERATE. *Lancet* 2020; **396**: 663 [PMID: [32891201](#) DOI: [10.1016/S0140-6736\(20\)30810-2](#)]
- 15 **Llovet JM**, Willoughby CE, Singal AG, Greten TF, Heikenwälder M, El-Serag HB, Finn RS, Friedman SL. Nonalcoholic steatohepatitis-related hepatocellular carcinoma: pathogenesis and treatment. *Nat Rev Gastroenterol Hepatol* 2023; **20**: 487-503 [PMID: [36932227](#) DOI: [10.1038/s41575-023-00754-7](#)]
- 16 **Rinella ME**, Lazarus JV, Ratziu V, Francque SM, Sanyal AJ, Kanwal F, Romero D, Abdelmalek MF, Anstee QM, Arab JP, Arrese M, Bataller R, Beuers U, Boursier J, Bugianesi E, Byrne CD, Castro Narro GE, Chowdhury A, Cortez-Pinto H, Cryer DR, Cusi K, El-Kassas M, Klein S, Eskridge W, Fan J, Gawrieh S, Guy CD, Harrison SA, Kim SU, Koot BG, Korenjak M, Kowdley KV, Lacaillle F, Loomba R, Mitchell-Thain R, Morgan TR, Powell EE, Roden M, Romero-Gómez M, Silva M, Singh SP, Sookoian SC, Spearman CW, Tiniakos D, Valenti L, Vos MB, Wong VW, Xanthakos S, Yilmaz Y, Younossi Z, Hobbs A, Villota-Rivas M, Newsome PN; NAFLD Nomenclature consensus group. A multisociety Delphi consensus statement on new fatty liver disease nomenclature. *J Hepatol* 2023; **79**: 1542-1556 [PMID: [37364790](#) DOI: [10.1016/j.jhep.2023.06.003](#)]
- 17 **Rinella ME**, Lazarus JV, Ratziu V, Francque SM, Sanyal AJ, Kanwal F, Romero D, Abdelmalek MF, Anstee QM, Arab JP, Arrese M, Bataller R, Beuers U, Boursier J, Bugianesi E, Byrne CD, Castro Narro GE, Chowdhury A, Cortez-Pinto H, Cryer DR, Cusi K, El-Kassas M, Klein S, Eskridge W, Fan J, Gawrieh S, Guy CD, Harrison SA, Kim SU, Koot BG, Korenjak M, Kowdley KV, Lacaillle F, Loomba R, Mitchell-Thain R, Morgan TR, Powell EE, Roden M, Romero-Gómez M, Silva M, Singh SP, Sookoian SC, Spearman CW, Tiniakos D, Valenti L, Vos MB, Wong VW, Xanthakos S, Yilmaz Y, Younossi Z, Hobbs A, Villota-Rivas M, Newsome PN; NAFLD Nomenclature consensus group. A multisociety Delphi consensus statement on new fatty liver disease nomenclature. *Hepatology* 2023; **78**: 1966-1986 [PMID: [37363821](#) DOI: [10.1097/HEP.0000000000000520](#)]
- 18 **Rinella ME**, Lazarus JV, Ratziu V, Francque SM, Sanyal AJ, Kanwal F, Romero D, Abdelmalek MF, Anstee QM, Arab JP, Arrese M, Bataller R, Beuers U, Boursier J, Bugianesi E, Byrne CD, Narro GEC, Chowdhury A, Cortez-Pinto H, Cryer DR, Cusi K, El-Kassas M, Klein S, Eskridge W, Fan J, Gawrieh S, Guy CD, Harrison SA, Kim SU, Koot BG, Korenjak M, Kowdley KV, Lacaillle F, Loomba R, Mitchell-Thain R, Morgan TR, Powell EE, Roden M, Romero-Gómez M, Silva M, Singh SP, Sookoian SC, Spearman CW, Tiniakos D, Valenti L, Vos MB, Wong VW, Xanthakos S, Yilmaz Y, Younossi Z, Hobbs A, Villota-Rivas M, Newsome PN; NAFLD Nomenclature consensus group. A multisociety Delphi consensus statement on new fatty liver disease nomenclature. *Ann Hepatol* 2024; **29**: 101133 [PMID: [37364816](#) DOI: [10.1016/j.aohep.2023.101133](#)]
- 19 **Hardy T**, Wonders K, Younes R, Aithal GP, Aller R, Allison M, Bedossa P, Betsou F, Boursier J, Brosnan MJ, Burt A, Cobbold J, Cortez-Pinto H, Day CP, Dufour JF, Ekstedt M, Francque S, Harrison S, Miele L, Nasr P, Papatheodoridis G, Petta S, Tiniakos D, Torstenson R, Valenti L, Holleboom AG, Yki-Jarvinen H, Geier A, Romero-Gomez M, Ratziu V, Bugianesi E, Schattenberg JM, Anstee QM; LITMUS Consortium. The European NAFLD Registry: A real-world longitudinal cohort study of nonalcoholic fatty liver disease. *Contemp Clin Trials* 2020; **98**: 106175 [PMID: [33045403](#) DOI: [10.1016/j.cct.2020.106175](#)]
- 20 **Song SJ**, Lai JC, Wong GL, Wong VW, Yip TC. Can we use old NAFLD data under the new MASLD definition? *J Hepatol* 2024; **80**: e54-e56 [PMID: [37541393](#) DOI: [10.1016/j.jhep.2023.07.021](#)]
- 21 **Younossi ZM**, Alqahtani SA, Alswat K, Yilmaz Y, Keklikkiran C, Funuyet-Salas J, Romero-Gómez M, Fan JG, Zheng MH, El-Kassas M, Castera L, Liu CJ, Wai-Sun Wong V, Zelber-Sagi S, Allen AM, Lam B, Treeprasertsuk S, Hameed S, Takahashi H, Kawaguchi T, Schattenberg JM, Duseja A, Newsome PN, Francque S, Spearman CW, Castellanos Fernández MI, Burra P, Roberts SK, Chan WK, Arrese M, Silva M, Rinella M, Singal AK, Gordon S, Fuchs M, Alkhoury N, Cusi K, Loomba R, Ranagan J, Eskridge W, Kautz A, Ong JP, Kugelmas M, Eguchi Y, Diago M, Yu ML, Gerber L, Fornaresio L, Nader F, Henry L, Racila A, Golabi P, Stepanova M, Carrieri P, Lazarus JV; Global NASH Council. Global survey of stigma among physicians and patients with nonalcoholic fatty liver disease. *J Hepatol* 2024; **80**: 419-430 [PMID: [37984709](#) DOI: [10.1016/j.jhep.2023.11.004](#)]





Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [office@baishideng.com](mailto:office@baishideng.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

