

[全部](#) [图片](#) [视频](#) [新闻](#) [购物](#) [地图](#) [图书](#)

找到约 3,090 条结果

时间不限[过去 1 小时内](#)[过去 24 小时内](#)[过去 1 周内](#)[过去 1 个月内](#)[过去 1 年内](#)**所有结果**[精确匹配](#)

[Google 学术: Is Endoscopic Papillary Balloon Dilatation \(EPBD\) Really a Risk Factor for Post-ERCP Pancreatitis?](#)

[Endoscopic balloon dilation of the biliary sphincter ...](#) - Baron - 被引用次数: 292

[... managed by endoscopic papillary balloon dilation](#) - Tsujino - 被引用次数: 74

[Endoscopic sphincterotomy plus large-balloon dilation ...](#) - Heo - 被引用次数: 208

[Reappraisal of endoscopic papillary balloon dilation for the ...](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC4325312/ ▼

16 Feb 2015 ... The risk factors of pancreatitis included dysfunction of sphincter of Oddi, young ... tract soon after endoscopic papillary balloon dilation (EPBD) and four patients The balloon is innocent and the pancreatitis may actually result from the dilation for removal of large common bile duct stones during ERCP.

[Is the Balloon Dilation Duration in Endoscopic Papillary Balloon ...](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC4323435/

31 Jan 2015 ... Endoscopic papillary balloon dilation (EPBD) was proposed as an alternative to ... pancreatic stent in patients at a high risk of post-ERCP pancreatitis. As a result, all factors that might affect the occurrence of post-EPBD pancreatitis were ... Kato T, Moriwaki H. Can endoscopic papillary balloon dilation really ...

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 24905

Manuscript Type: TOPIC HIGHLIGHT

2016 Gastrointestinal Endoscopy: Global view

**Is endoscopic papillary balloon dilatation really a risk factor for Post-ERCP
pancreatitis?**

Toshio Fujisawa, Koichi Kagawa, Kantaro Hisatomi, Kensuke Kubota, Atsushi

Nakajima, Nobuyuki Matsuhashi

Match Overview

There are no matching sources for this report

Is endoscopic papillary balloon dilatation really a risk factor for Post-ERCP p

全部 图片 视频 新闻 购物 地图 图书

找到约 44,400 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Endoscopic Papillary Balloon Dilation for Removal of ... - NCBI](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC3065083/

16 Mar 2011 ... Endoscopic papillary balloon dilation (EPBD) is an alternative method of ...
Actually, meta-analysis demonstrated that post-ERCP pancreatitis occurred ... risk factor for
post-ERCP pancreatitis (OR, 1.59; 95% CI, 1.06-2.39).

[Balloon dilation itself may not be a major determinant of post ... - NCBI](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC4258560/

7 Dec 2014 ... Acute pancreatitis after endoscopic papillary balloon dilation ... Thus,
pancreatitis -associated EPBD is a very controversial and serious topic[53,54]. ... Younger
age is a risk factor for post-ERCP pancreatitis[3,4,6,12,15], but not ...

[Endoscopic Papillary Balloon Dilation and Extraction of Bile Duct ...](#)

www.sciencedirect.com/science/article/pii/S2212097113702012

28 Sep 2013 ... Endoscopic Papillary Balloon Dilation and Extraction of Bile Duct Stone. J
Pohl ... To lower the risk of post-ERCP pancreatitis after balloon dilation of the papilla,
prophylactic stent ... Complications and Risk Factors ... Can Endoscopic Papillary Balloon
Dilation Really Preserve Sphincter of Oddi Function?

[Prevention of Post-ERCP Pancreatitis](#)

全部 图片 新闻 视频 购物 更多 ▾ 搜索工具

找到约 45,200 条结果 (用时 0.73 秒)

The Ballooning Time in Endoscopic Papillary Balloon ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: BW Bang - 2010 - 被引用次数: 9 - 相关文章

2010年8月31日 - Endoscopic papillary balloon dilation (EPBD) is a safe and effective method for This study did not assess risk factors for post-ERCP pancreatitis, but our ... Kato T, Moriwaki H. Can endoscopic papillary balloon dilation really

Endoscopic Papillary Balloon Dilation for Removal of ... - NCBI

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: JW Chung - 2011 - 被引用次数: 26 - 相关文章

2011年3月16日 - Endoscopic papillary balloon dilation (EPBD) is an alternative method of Actually, meta-analysis demonstrated that post-ERCP pancreatitis occurred ... risk factor for post-ERCP pancreatitis (OR, 1.59; 95% CI, 1.06-2.39).

Endoscopic Papillary Balloon Dilation and Extraction of Bile ...

www.sciencedirect.com/science/article/.../S2212097113702012 ▾ 翻译此页

作者: J Pohl - 2013 - 相关文章

2013年9月28日 - Endoscopic Papillary Balloon Dilation and Extraction of Bile Duct Stone. J Pohl To lower the risk of post-ERCP pancreatitis after balloon dilation of the papilla, prophylactic stent ... Complications and Risk Factors ... Can Endoscopic Papillary Balloon Dilation Really Preserve Sphincter of Oddi Function?

^[PDF] Complications of ERCP

www.spg.pt/wp-content/.../11/2012-ERCP-complications.pdf ▾ 翻译此页

monly performed endoscopic procedure.2 The diagnostic ... risk of PEP with endoscopic papillary balloon dilation ... Risk factors for post-ERCP pancreatitis in.

^[PDF] Risk Factors for Post-ERCP Pancreatitis - University of ...

umanitoba.ca/faculties/medicine/.../shermanchengPEP.pdf ▾ 翻译此页

作者: CL Cheng - 2006 - 被引用次数: 514 - 相关文章