



学术搜索

获得 3 条结果 (用时0.11秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

包括专利

包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[引用] Sa1074 Is Split-Dose Regimen of 2L Polyethylene Glycol Plus Ascorbic Acid tolerable for Early Morning Visit in Health Check-up Center?

C Lee, JY Seo, EH Jin, JH Lim, HY Kang... - Gastrointestinal ..., 2016 - giejournal.org

Skip to Main Content. Gastrointestinal Endoscopy Home. Mobile. ASGE; YouTube;

Twitter; Facebook; RSS Feeds. Login Email/Username: Password: Remember me

Forgot password? Register Register as a Guest. Register and ...

引用 保存

SATURDAY ABSTRACTS

[XC Zhang, PH Zhou - giejournal.org](#)

... 8.9%) Grade I 126 (7.5%) Grade II 24 (1.4%) Mucosa injury, n(%) 291 (17.3%) Type I 257 (15.3%)

Type II 34 (2.0 ... Methods: All ERCP and EUS scopes are disinfected using Custom Ultrasonic 83

plus AERs after hand cleaning by strict OEM ... 2 million other ambulatory procedures ...

相关文章 引用 保存

Colonic and Anorectal Disorders

BC YOON - Journal of Gastroenterology and Hepatology, 2014 - Wiley Online Library

... Tissue acid-fast bacilli staining and tuberculin skin test was negative, but interferon-gamma assay

was ... Key Word(s): 1. ulcerative colitis; 2. dysplasia; 3. endoscopic submucosal dissection. ... can

be easily examined, and the possibility to be connected for the early detection of the ...

所有 3 个版本 引用 保存

创建快讯

**Name of Journal:** *World Journal of Gastroenterology*

**ESPS Manuscript NO:** 31681

**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Cohort Study*

**Is a split-dose regimen of 2 L polyethylene glycol plus ascorbic acid tolerable for an early morning visit to a comprehensive medical check-up?**

Seo JY *et al.* Split-dose regimen of low volume polyethylene glycol

<sup>2</sup> Ji Yeon Seo, Changhyun Lee, Eun Hyo Jin, Mi Hyun Yun, Joo Hyun Lim, Hae Yeon Kang, Jong In Yang, Su Jin Chung, Sun Young Yang, Joo Sung Kim

<sup>4</sup> Ji Yeon Seo, Changhyun Lee, Eun Hyo Jin, Joo Hyun Lim, Hae Yeon Kang, Jong In Yang, Su Jin Chung, Sun Young Yang, Joo Sung Kim, Department of Internal Medicine and Healthcare Research Institute, Healthcare System Gangnam Center, Seoul National University Hospital, Seoul 135-984, South Korea

Mi Hyun Yun, Department of Nursing, Healthcare System Gangnam Center, Seoul

### Match Overview

1	<b>Internet</b> 159 words crawled on 03-Jul-2016 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	4%
2	<b>Crossref</b> 122 words Lee, Changhyun, Ji Yeon Seo, Eun Hyo Jin, Joo Hyun Lim, Hae Yeon Kang, Jong In Yang, Su Jin Chung, Sun Young	3%
3	<b>Crossref</b> 25 words Siddiqui, A.A.. "Duration of the interval between the completion of bowel preparation and the start of colonoscopy pre..."	1%
4	<b>Crossref</b> 25 words "ASGE Program", <i>Gastrointestinal Endoscopy</i> , 2016.	1%
5	<b>Crossref</b> 21 words Yang, Hyo-Joon, Seon Hee Lim, Changhyun Lee, Ji Min Choi, Jong In Yang, Su Jin Chung, Seung Ho Choi, Jong Pil I	1%
6	<b>Internet</b> 20 words crawled on 29-Jun-2015 <a href="http://www.e-sciencecentral.org">www.e-sciencecentral.org</a>	1%

找到约 1,550 条结果 (用时 0.86 秒)

### Bisacodyl plus split 2-L polyethylene glycol-citrate-simethicone ... - NCBI

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: F Valiante - 2013 - 被引用次数: 16 - 相关文章

2013年9月7日 - US National Library of Medicine ... CONCLUSION: BIS plus split 2-L PEG-CS was as effective as but ... Screening for early detection and removal of premalignant ... A split dose regimen, in which the first half dose is taken on the day ... sulphate-free bowel preparation consisting of PEG 4000, citric acid, ...

### Efficacy of morning-only 4 liter sulfa free polyethylene glycol vs 2 liter ...

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: JM Rivas - 2014 - 被引用次数: 1 - 相关文章

2014年8月14日 - Colonoscopy is considered the gold standard test for early detection and prevention of ... The addition of ascorbic acid to a 2-L PEG solution has been found to work ... by adding up the three segments (0-12) plus the fluid score (0-2) ... the purgative in its entirety in the morning or using a split dose regimen.

### Ascorbic acid PEG-2L is superior for early morning colonoscopies in ...

<https://dialnet.unirioja.es/servlet/articulo?codigo=5283866> - 翻译此页

作者: CR Miguel - 2015 - 相关文章

The quality of colon cleansing and the tolerability of antegrade preparation are ... Ascorbic acid PEG-2L is superior for early morning colonoscopies in .... High-volume polyethylene glycol plus electrolytes solution (PEG-ELS) is ... that for morning colonoscopies, split-dose regimens are superior to single-dose regimens.

### [PDF] expert - ASGE

[www.asge.org/uploadedfiles/asge\\_news/aejan11.pdf](http://www.asge.org/uploadedfiles/asge_news/aejan11.pdf) ▾ 翻译此页

the morning of colonoscopy for some afternoon procedures. ... ascorbic acid (MoviPrep, Salix Pharmaceuticals, Morrisville, N.C.) ... might be as effective as MoviPrep if it were given in split doses, but it is not, ... select a 2-liter PEG-ELS when patients lack risk factors for failed preparation, i.e., ... Medical Systems, Braintree.

### Efficacy and acceptability of sodium picosulphate/magnesium citrate ...

[全部](#)[购物](#)[图片](#)[新闻](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 1,540 条结果 (用时 0.75 秒)

小提示：仅限搜索简体中文结果。您可以在设置中指定搜索语言

### [Bisacodyl plus split 2-L polyethylene glycol-citrate-simethicone ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - [翻译此页](#)

作者：F Valiante - 2013 - 被引用次数：16 - [相关文章](#)

2013年9月7日 - US National Library of **Medicine** ... CONCLUSION: BIS **plus** split 2-L PEG-CS was as effective as but ... Screening for **early** detection and removal of premalignant ... A **split dose regimen**, in which the first half dose is taken on the day ... sulphate-free bowel preparation consisting of PEG 4000, citric **acid**, ...

### [Efficacy of morning-only 4 liter sulfa free polyethylene glycol vs 2 liter ...](#)

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - [翻译此页](#)

作者：JM Rivas - 2014 - 被引用次数：1 - [相关文章](#)

2014年8月14日 - Colonoscopy is considered the gold standard **test** for **early** detection and prevention of ... The addition of **ascorbic acid** to a 2-L PEG solution has been found to work ... by adding **up** the three segments (0-12) **plus** the fluid score (0-2). .... the purgative in its entirety in the **morning** or using a **split dose regimen**.

### [Ascorbic acid PEG-2L is superior for early morning colonoscopies in ...](#)

<https://dialnet.unirioja.es/servlet/articulo?codigo=5283866> - [翻译此页](#)

作者：CR Miguel - 2015 - [相关文章](#)

The quality of colon cleansing and the **tolerability** of antegrade preparation are ... **Ascorbic acid PEG-2L** is superior for **early morning** colonoscopies in .... High-volume **polyethylene glycol plus** electrolytes solution (PEG-ELS) is ... that for **morning** colonoscopies, **split-dose regimens** are superior to single-dose **regimens**.

### [\[PDF\] expert - ASGE](#)

[www.asge.org/uploadedfiles/asge\\_news/aejan11.pdf](http://www.asge.org/uploadedfiles/asge_news/aejan11.pdf) ▼ [翻译此页](#)

the **morning** of colonoscopy for some afternoon procedures. ... **ascorbic acid** (MoviPrep, Salix