

World Journal of *Radiology*

World J Radiol 2024 September 28; 16(9): 375-496



EDITORIAL

- 375 Innovative approaches beyond periprocedural hydration for preventing contrast-induced acute kidney injury
Cheng CH, Hao WR, Cheng TH

ORIGINAL ARTICLE**Retrospective Study**

- 380 Intentionally unilateral prostatic artery embolization: Patient selection, technique and potential benefits
Moschouris H, Stamatou K
- 389 Cryoablation of osteoid osteomas: Is it a valid treatment option?
Michailidis A, Panos A, Samoladas E, Dimou G, Mingou G, Kosmoliaptis P, Arvaniti M, Giankoulof C, Petsatodis E
- 398 Radiological findings of February 2023 twin earthquakes-related spine injuries
Bolukçu A, Erdemir AG, İdilman İS, Yıldız AE, Çoban Çifçi G, Onur MR, Akpınar E

Observational Study

- 407 Retinal microcirculation changes in prediabetic patients with short-term increased blood glucose using optical coherence tomography angiography
Hu K, Lv BJ, Zuo HJ, Li QF, Huang FF, Zhang T, Huang RX, Zheng SJ, Wan WJ
- 418 Nomogram for predicting short-term response to anti-vascular endothelial growth factor treatment in neovascular age-related macular degeneration: An observational study
Huang ZH, Tu XZ, Lin Q, Tu M, Lin GC, Zhang KP
- 429 Cerebral perfusion in patients with unilateral internal carotid artery occlusion by dual post-labeling delays arterial spin labeling imaging
Zhang GR, Zhang YY, Liang WB, Ding D

CASE REPORT

- 439 Acquired factor XIII deficiency presenting with multiple intracranial hemorrhages and right hip hematoma: A case report
Wang L, Zhang N, Liang DC, Zhang HL, Lin LQ
- 446 Myelin oligodendrocyte glycoprotein-associated transverse myelitis after SARS-CoV-2 infection: A case report
Zheng JR, Chang JL, Hu J, Lin ZJ, Lin KH, Lu BH, Chen XH, Liu ZG
- 453 Extralobar pulmonary sequestration in children with abdominal pain: Four case reports
Jiang MY, Wang YX, Lu ZW, Zheng YJ

- 460 Behcet's disease-related panuveitis following COVID-19 vaccination: A case report
Lin RT, Liu PK, Chang CW, Cheng KC, Chen KJ, Chang YC
- 466 Hyperparathyroidism presented as multiple pulmonary nodules in hemodialysis patient status post parathyroidectomy: A case report
Chiang PH, Ko KH, Peng YJ, Huang TW, Tang SE
- 473 Secondary rectal linitis plastica caused by prostatic adenocarcinoma - magnetic resonance imaging findings and dissemination pathways: A case report
Labra AA, Schiappacasse G, Cocio RA, Torres JT, González FO, Cristi JA, Schultz M
- 482 *Pneumocystis* pneumonia in stage IIIA lung adenocarcinoma with immune-related acute kidney injury and thoracic radiotherapy: A case report
Zheng YW, Pan JC, Wang JF, Zhang J
- 489 Prolonged course of Paxlovid administration in a centenarian with COVID-19: A case report
Zhang YX, Tang J, Zhu D, Wu CY, Liang ML, Huang YT

ABOUT COVER

Editorial Board Member of *World Journal of Radiology*, Roberto Grassi, MD, Professor, Chief, Department of Radiology, University of Campania Luigi Vanvitelli, Napoli, 80138, Italy. roberto.grassi@unicampania.it

AIMS AND SCOPE

The primary aim of *World Journal of Radiology (WJR, World J Radiol)* is to provide scholars and readers from various fields of radiology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJR mainly publishes articles reporting research results and findings obtained in the field of radiology and covering a wide range of topics including state of the art information on cardiopulmonary imaging, gastrointestinal imaging, genitourinary imaging, musculoskeletal imaging, neuroradiology/head and neck imaging, nuclear medicine and molecular imaging, pediatric imaging, vascular and interventional radiology, and women's imaging.

INDEXING/ABSTRACTING

The *WJR* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for *WJR* as 1.4; JIF without journal self cites: 1.4; 5-year JIF: 1.8; JIF Rank: 132/204 in radiology, nuclear medicine and medical imaging; JIF Quartile: Q3; and 5-year JIF Quartile: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Wen-Bo Wang*; Production Department Director: *Xu Guo*; Cover Editor: *Jia-Ping Yan*.

NAME OF JOURNAL

World Journal of Radiology

ISSN

ISSN 1949-8470 (online)

LAUNCH DATE

January 31, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Thomas J Vogl

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1949-8470/editorialboard.htm>

PUBLICATION DATE

September 28, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Retrospective Study

Radiological findings of February 2023 twin earthquakes-related spine injuries

Ayşenur Bolukçu, Ahmet Gürkan Erdemir, İlkey Sedakat İdilman, Adalet Elçin Yıldız, Gökçen Çoban Çifçi, Mehmet Ruhi Onur, Erhan Akpınar

Specialty type: Radiology, nuclear medicine and medical imaging

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade B

Novelty: Grade A

Creativity or Innovation: Grade A

Scientific Significance: Grade B

P-Reviewer: Elzeiny A

Received: May 27, 2024

Revised: August 20, 2024

Accepted: August 28, 2024

Published online: September 28, 2024

Processing time: 122 Days and 6.4 Hours



Ayşenur Bolukçu, Ahmet Gürkan Erdemir, İlkey Sedakat İdilman, Adalet Elçin Yıldız, Gökçen Çoban Çifçi, Mehmet Ruhi Onur, Erhan Akpınar, Department of Radiology, Hacettepe University, Faculty of Medicine, Ankara 06100, Türkiye

Erhan Akpınar, Department of Radiology, University of Chicago, Chicago, IL 60637, United States

Corresponding author: Ahmet Gürkan Erdemir, MD, Doctor, Instructor, Department of Radiology, Hacettepe University, Faculty of Medicine, No. 6/1 Adnan Saygun Street, Ankara 06100, Türkiye. a.gurkan.erdemir@gmail.com

Abstract

BACKGROUND

The February 6, 2023, twin earthquakes in Türkiye caused significant structural damage and a high number of injuries, particularly affecting the spine, which underscores the importance of understanding the distribution and nature of vertebral injuries in disaster victims.

AIM

To investigate the distribution of radiological findings of vertebral injuries in patients referred to a major tertiary center during the February 6, 2023 twin earthquakes in Türkiye.

METHODS

With the approval of the institutional ethics committee, 1216 examinations of 238 patients transferred from the region to a tertiary major hospital after the twin earthquakes of February 6, 2023, were retrospectively analyzed for spine injuries.

RESULTS

Spine computed tomography (CT) scans were performed in 192 of 238 patients with a suspected spinal injury, 42 of whom also had an magnetic resonance imaging (MRI). In 86 of 192 patients (44.79%; M:F = 33:53) a spinal fracture was detected on CT and in 33 of 42 patients (78.57%; M:F = 20:13) a spinal injury was found on MRI. Of the 86 patients in whom vertebral injury was detected, fractures were detected in the Denis-B group in 33, Denis-C in 4, Denis-D in 20 and Denis-E in 11 patients. Among the vertebral bodies: 40 "compression fractures", 17 "burst fractures", 5 "translational dislocation fractures", 5 "flexion-distraction fractures" and

58 "prolonged forced fetal posture fractures" were detected. In addition, isolated transverse or spinous process fractures were found in eighteen vertebrae.

CONCLUSION

Our study highlights the prevalence and diverse spectrum of spinal injuries following the February 6, 2023 twin earthquakes in Turkey underscoring the urgent need for effective management strategies in similar disaster scenarios, and emphasizing the "prolonged forced fetal posture" damage we encountered in earthquake victims who remained under the collapse for a long time.

Key Words: Accidental injuries; Compression fractures; Crush injuries; Earthquakes; Spine

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: In the classification of damages from the earthquakes in Turkey on February 6, 2023, it was found that the injuries were predominantly crush-related rather than escape-related. Observations revealed that patients frequently sustained damage to the thoracolumbar regions due to prolonged fetal positioning under debris. It was determined that a significant portion of patients with spinal canal injuries had damage consistent with the thoracolumbar junction.

Citation: Bolukçu A, Erdemir AG, İdilman İS, Yıldız AE, Çoban Çifçi G, Onur MR, Akpınar E. Radiological findings of February 2023 twin earthquakes-related spine injuries. *World J Radiol* 2024; 16(9): 398-406

URL: <https://www.wjgnet.com/1949-8470/full/v16/i9/398.htm>

DOI: <https://dx.doi.org/10.4329/wjr.v16.i9.398>

INTRODUCTION

The catastrophic twin earthquakes measuring 7.8 and 7.7 on the Richter scale struck Kahramanmaraş, Turkey, on February 6, 2023, marked a shattering chapter in the region's history, claiming the lives of over 50000 people and leaving thousands more injured[1,2]. The aftermath of these seismic events, saw a saturation of health services, as the scale of the devastation exceeded the capacity of local medical facilities. Unlike other natural disasters, earthquakes impose a disproportionately high toll on human lives, underscoring the urgent need for effective response mechanisms in the face of such crises[3]. Among the myriad of injuries, spinal trauma is of particular concern as it affects a significant proportion of earthquake survivors. In particular, the unique circumstances surrounding these injuries, including the prevalence of crush injuries and falls, pose a particular challenge in diagnosis and treatment[4].

Following the earthquakes, the increase in crush injuries and falls precipitated a wave of spinal trauma that strained already overstretched healthcare resources[5]. Notably, earthquake survivors were found to be at heightened risk of developing crush syndrome, a potentially life-threatening condition characterized by the myonecrosis, rapid hypovolemia and excessive third-space edema[6,7]. In addition, the prevalence of vertebral fractures, often due to the high-force impacts associated with earthquakes, underscored the complex nature of spinal injuries in this context[8].

Compression fractures typically occur due to axial loading, where the vertebral body is crushed under excessive pressure, leading to a wedge-shaped deformity. Of particular concern was the emergence of a specific injury pattern known as "forced prolonged fetal posture", observed predominantly among earthquake survivors and requires specialized diagnostic approaches for accurate assessment[9]. The likely mechanism for a consecutive spine flexion injury due to forced prolonged fetal posture involves excessive and sustained forward bending of the spine, which specifically exerts abnormal compressive forces on the vertebral bodies. This stress can lead to microfractures, compression fractures, or other traumatic injuries to the vertebrae, compromising their structural integrity and resulting in significant spinal damage. Burst fractures, caused by a more severe axial load, result in the vertebral body shattering and dispersing fragments, which can impinge on the spinal canal. Translational dislocation fractures, on the other hand, involve the vertebrae being displaced horizontally, often due to high-energy trauma, leading to severe instability. Flexion-distraction fractures arise from hyperflexion, where the anterior vertebral structures are compressed while the posterior elements are distracted, causing a failure in the vertebral integrity.

The main objective of this paper is to describe the distribution and characteristics of spinal injuries in the survivors of the February 6 twin earthquakes. By analyzing multidetector row computed tomography (CT) and magnetic resonance imaging (MRI) findings from a cohort of individuals who had spinal injuries after the earthquake, we attempt to clarify the incidence and patterns of spinal trauma in this population.

MATERIALS AND METHODS

Study design

The study was conducted following the guidelines of the Declaration of Helsinki and approval for this retrospective study was obtained from the Institutional Ethics Board. Our study included patients who were referred to our hospital from the disaster area during the period from February 6 to February 28, 2023. We included all patients from the disaster area in our study, without considering any age restrictions.

CT and MRI protocol

All CT examinations were performed in the emergency radiology station, *via* Somatom Perspective 64-Slice Siemens® CT device (Erlangen/Germany). CT scan parameters were as follows: Tube voltage: 120 kV, tube current set with optimized automatic exposure control (the ref mAs value of CT was 140 mAs), collimation thickness: 0.6-2 mm, tube rotation time: 0.6-1 seconds and collimated section thickness: 2-5 mm. The iterative reconstruction (ADMIRE, strength 2) with 30 seconds soft tissue and 60 seconds bone kernel was used as a reconstruction algorithm to reduce the radiation dose.

MRI imaging was performed with a Siemens® 1.5 Tesla Aera device (Erlangen/Germany). Imaging parameters for the cervical, thoracic, lumbar and sacral regions included a slice thickness of approximately 3-5 mm, a repetition time of approximately 2000-2500 ms, an Time to Echo of approximately 20-40 ms, a matrix size of approximately 256 × 256 and a field of view of 200-350 mm, depending on the region scanned. The MRI protocol for the spine examination included axial T1-weighted, axial T2-weighted, axial fat-suppressed T2-weighted, sagittal T1-weighted, sagittal T2-weighted, and sagittal fat-suppressed T2-weighted sequences.

Data collection and statistics

All CT and MRI examinations of the spine of these patients were included in the study. The distribution of spinal fractures across the anatomical regions, including the cervical, lumbar, thoracic and sacral regions, was delineated. Fractures were categorized using the Denis, Association for Osteosynthesis (AO), and TLICS classifications[10-13]. Patients were classified according to the AO classification and grouped according to their instabilities, considering the Denis spinal column distributions. To investigate the traumatic effects of prolonged fetal position due to confinement under the debris, we also investigated the incidence of typical consecutive spinal fractures of the thoracolumbar junction, which are more vulnerable and correspond to levels T10-L2.

Statistical analysis included descriptive statistics and incidence calculations to describe the distribution of CT and MRI findings of spinal fractures and to investigate consecutive spinal fractures due to prolonged fetal position.

RESULTS

1216 examinations were carried out on 238 patients who were transferred from the disaster area to our hospital. Of these, 192 patients underwent a CT scan and 42 an MRI scan. Of the 192 CT scans, 163 examinations of the cervical, thoracic and lumbar vertebra were performed, while the remaining scans examined specific anatomical regions based on clinical symptoms.

Of the patients who underwent CT scans, 86 patients had vertebral fractures, of which 63 (73.25%) had multiple fractures and 23 (27.75%) had single fractures. Total fracture number was 287. In addition, 47 patients out of 63 (74.60%) with multiple fractures had consecutive fractures, while 16 patients (25.40%) had scattered fractures. Grouping patients according to TLICS groups of involvement revealed various types of vertebral fractures, including 40 compression fractures, 17 burst fractures, 5 flexion-distraction fractures, 5 translational dislocation fractures, and 58 fractures due to prolonged fetal posture. For examples of vertebral fractures in these patients, please see Figures 1, 2, 3, 4, and 5.

Most injuries occurred in the lumbar region. Sixty-three patients out of 86 patients (73.25%) sustained lumbar vertebra fractures. Furthermore, 140 vertebral fractures were assessed within lumbar region. Of these 63 patients, 33 patients (52.38%) had type B fractures according to the Denis classification, and 26 patients (41.26%) had type A4 fractures according to the AO classification.

Injuries to the thoracolumbar junction (T10-L2) were most common in earthquake victims. Fifty-eight patients had fractures at this level with total 93 fractures. Looking at the distribution of thoracolumbar region injuries based on their vertebral levels, ten (10.75%) fractures were at the T10 Level, sixteen (17.20%) at the T11 level, nineteen (20.43%) at the T12 level, twenty-three (24.73%) at the L1 level and twenty-five (26.88%) at the L2 level. In addition, according to the Denis classification, 27 of them had unstable 3-column fractures in the lumbar region.

Fractures affecting the thoracic vertebrae were identified in 48 out of 86 earthquake victims. Among detected 287 vertebral fractures, 131 were localized to the thoracic vertebrae. Patients with thoracic fractures had different fracture types, with A0 fractures being the most common. Of the 48 patients with thoracic vertebral fractures, 13 patients (27.08%) had thoracic A0 fractures with 44 A0 fractures (33.58%) identified among the 131 thoracic vertebral fractures. According to the Denis classification, nine patients had unstable 3-column fractures in the thoracic region.

No unstable 3-column fractures were observed in patients with cervical vertebral fractures. Ten patients with fractures in the sacral region were grouped according to the Denis zonal distribution classification, and an equal number of zone 2 and zone 3 fractures were observed.

Concomitant fractures in patients with vertebral fractures included ribs (43), upper extremities (10), lower extremities (19), scapulas (11), sternum (4), viscerocranium (6), clavicles (6), and neurocranium (2).




Figure 1 Prolonged fetal posture injury. A 23-year old male who jammed under debris about 15 minutes. A: Computed tomography image reveals T8 to L1 consecutive mild compression fractures (arrows), which corresponds to thoracolumbar area injury; B and C: T1W image and T2-STIR sequence show T8 to L1 vertebrae fracture and edema around them; D: The image of the volume rendering technique shows especially the fractures of the T12 and L1 vertebrae.




Figure 2 Compression fracture. A 27-year old male who was injured while escaping from the earthquake. A: Computed tomography image reveals T1-T5 vertebrae and T7 vertebra compression fractures (arrows); B and C: T2-STIR and T2W images show C6-T5 vertebrae and T7 vertebra compression fractures (arrows) and edema around them.




Figure 3 Flexion-distraction fracture. A 35-year old male who jammed under debris about 3 hours. A: Computed tomography image reveals the flexion-distraction type of fracture involving T12 to L4 vertebrae, especially with anteriorly losing height (orange arrows) and distraction of posterior processes (yellow arrows); B: STIR sequence shows fractures of T12 to L4 vertebrae and edema around fractures.

MRI showed retropulsion and narrowing of the spinal canal in 12 patients, and CT showed this in 19 patients. When examining the distribution of fractures causing spinal canal narrowing with retro-pulsed fragments, the most common fractures occurred at the L1 vertebra in 7 patients, followed by the T12 and L2 vertebrae in every three patients, the L4 vertebra in 2 patients, and one patient each in the L3, L5, T11, and T9 vertebrae.

Detailed information on the distribution of fracture types and the positive CT and MRI findings can be found in Tables 1, 2 and 3.

DISCUSSION

The twin earthquakes that struck southeastern Turkey on February 6, 2023, represent a major disaster. Considering the persistent presence of humanity, various types of disasters, especially earthquakes, remain inevitable. A critical aspect requiring attention among earthquake victims is the incidence of vertebral fractures, as documented in the existing literature from various earthquakes worldwide and corroborated by our observations following the twin earthquakes[5, 14]. Analysis of earthquake victims reveals that a substantial portion sustain injuries from being trapped under debris, while others experience vertebral fractures during attempts to escape the earthquake's effects. Li *et al*[15] demonstrated in their research (focused on an earthquake with a magnitude of 8.0) that crush-related injuries constitute the majority of cases; similarly, our study found that most patients were trapped under debris. Consequently, contrast material examination was limited in these patients due to the potential risk of renal damage from crush injuries. Additionally, literature indicates that spinal injuries related to non-earthquake crush injuries are less common[5]. Our demographic analysis of earthquake victims shows a predominance of female patients, mirroring findings from another study and suggesting a possible link to how individuals cope with earthquake hazards. Since crush injuries are the primary cause of trauma rather than escape-related injuries, it is likely that young male victims are better able to avoid injury[5]. This observation aids in understanding the characteristics of the earthquake victim patient group.

In our earthquake-related cohort, the risk of lumbar spinal injury due to retro-pulsed bony fragments was most prevalent, with the highest incidence of fractures occurring in the lumbar region. This finding is consistent with both our observations and results from a study on spinal cord injuries sustained during an earthquake in China (Sichuan, 2008) [16]. Furthermore, another study on earthquakes in China suggests that the extensive coverage area of the lumbar region may contribute to its increased susceptibility to crush injuries, further validating our findings. Our study makes a unique contribution to the existing literature by observing that a substantial proportion of patients exhibited fractures in multiple vertebrae, a finding that aligns with previous studies. However, unlike the study by Dong *et al*[5], which identified minor fractures across multiple widely dispersed areas, our patient cohort predominantly involved consecutive fractures.

A study examining spinal fractures during the Wenchuan earthquake (with a magnitude of 8.0) identified burst and compression fractures as the most prevalent types[15]. Similarly, in our patient group, compression fractures emerged as the most frequently observed fracture type. Additionally, as indicated in the study by Dong *et al*[5], compression fractures

Table 1 Positivity rates and age-gender distribution

Total number of patients	Positive findings (%)	Gender distribution (%)	Mean age	Standard deviation	Median
CT (192)	86 (44.79)	Female: 53 (61.62); Male: 33 (38.37)	45.25	19.56	46.00
MRI (42)	33 (78.57)	Female: 20 (60.60); Male: 13 (39.39)	52.00	19.91	55.00

MRI: Magnetic resonance imaging; CT: Computed tomography.

Table 2 The distribution of injury types

Injury type	Number of patients (86 pts w 287 vertebra fractures) (%)	Mean age	Standard deviation	Median
Compression	40 (46.51)	40.82	20.97	38.50
Burst	17 (19.76)	53.47	18.53	61.00
Translation dislocation	5 (5.81)	49.40	14.97	52.00
Flexion distraction	5 (5.81)	30.20	17.66	26.00
Isolated transverse or spinous process fracture	18 (20.93)	49.88	15.72	52.00
Prolonged fetal posture injury	58 (67.44)	45.22	19.52	48.00

Table 3 The distribution of injury types according Denis-grading system

Denis	Number of patients (86 pts w 287 vertebra fractures) (%)	Mean age	Standard deviation	Median
B	33 (38.37)	40.06	22.04	36.00
C	4 (4.65)	54.00	11.31	54.00
D	20 (23.25)	50.75	18.38	60.00
E	11 (12.79)	43.09	20.95	52.00

may result from direct vertical force or axial loading due to falling objects, indicative of crush injury mechanisms. The high number of compression fractures in our study is correlated with the predominance of patients presenting with crush injuries. Although literature on patients with triple column fractures is limited, our study observed a significant occurrence of these unstable fractures among patients. The prevalence of such fractures in earthquake-related trauma exhibits high-energy trauma characteristics. Upon assessing the likelihood of spinal cord injury in our patient group through clinical evaluation and MRI findings, positive indications were observed in a minority of cases, consistent with findings reported in the literature[16,17]. Furthermore, the absence of myelopathy despite evident spinal canal compression on MRI was correlated with the lack of clinical symptoms in these patients.

The thoracolumbar junction, corresponding to the T10-L2 vertebral levels, is one of the most frequently traumatized areas due to its mechanical vulnerability. This region serves as a transition zone between the thoracic levels, which are stabilized by the ribs and sternum, and the more mobile lumbar levels[18]. Various studies have demonstrated how different body positions affect spinal injuries, and the thoracolumbar region is particularly susceptible to injury in earthquake scenarios[5,9,19-22]. Our research supports this, revealing that the majority of earthquake victims with vertebral fractures had injuries at the thoracolumbar junction (T10-L2). Specifically, when examining patients likely to experience neurological damage due to bony fragment retropulsion into the spinal canal, we found that the majority of these cases, totaling fourteen patients, involved the thoracolumbar junction.

The literature suggests that fractures in the thoracolumbar region occur more frequently due to the kyphotic posture patients often adopt, particularly within the "life triangle," as a protective mechanism against crush injuries[9,23]. This kyphosis-related damage is more likely because earthquake victims tend to minimize their surface area and adopt a hyper-flexed fetal position instinctively to shield themselves from harm. Additionally, the human spine's center of gravity is located anterior to the thoracolumbar junction, further contributing to the tendency of patients to assume this protective fetal position.

The study has several limitations. It is retrospective in nature and confined to a single-center setting. Additionally, while all patients underwent CT imaging, the availability of MRI imaging was limited. A significant limitation of our study is the lack of comprehensive data, as we only have information on the duration patients were trapped under rubble and the time elapsed between their rescue and hospital admission, which limits our ability to investigate the impact of time on vertebral injury patterns and distribution, highlighting the need for further research in this area. Despite these




Figure 4 Burst fracture. A 63-year old female who jammed under debris about 48 hours. A: Computed tomography image reveals the burst fracture of the L4 vertebra (arrows) with the loss of height; B and C: T2W and STIR sequence show the retropulsed bone fragment and narrowing of the spinal canal (arrows).




Figure 5 Translation-dislocation fracture. A 52-year old male who jammed under debris about 12 hours. A: Sagittal reformat computed tomography (CT) image reveals the translation type of L4 vertebra fracture (arrow), extending into the right facet joint. B: Axial CT image of the same patient reveals the right transverse process fracture of L4 extending into the right facet joint.

limitations, the study's strengths lie in its comprehensive analysis of vertebral fractures among earthquake victims, particularly its focus on the thoracolumbar junction and the correlation between injury patterns and trauma mechanisms.

CONCLUSION

In the classification of damages from the earthquakes in Turkey on February 6, 2023, it was found that the injuries were predominantly crush-related rather than escape-related. Observations revealed that patients frequently sustained damage to the thoracolumbar regions due to prolonged fetal positioning under debris. It was determined that a significant portion of patients with spinal canal injuries had damage consistent with the thoracolumbar junction.

FOOTNOTES

Author contributions: Bolukçu A designed research and wrote the paper; Erdemir AG performed research and wrote the paper; İdilman İS, Yildiz AE, Çifçi GC contributed figures, analytic tools and analyzed data; Onur MR conceptualized the research; Akpınar E supervised the research.

Institutional review board statement: The study was conducted following the guidelines of the Declaration of Helsinki and approval for this retrospective study was obtained from the Institutional Ethics Board (SBA 24/077).

Informed consent statement: Informed consent was not obtained due to the retrospective nature of the study.

Conflict-of-interest statement: There is no conflict of interest.

Data sharing statement: There is no data to be shared.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: Türkiye

ORCID number: Ahmet Gürkan Erdemir 0000-0001-7427-1039; İlky Sedakat İdilman 0000-0002-1913-2404; Adalet Elçin Yildiz 0000-0001-7208-8097; Gökçen Çoban Çifçi 0000-0002-0277-2042; Mehmet Ruhi Onur 0000-0003-1732-7862; Erhan Akpınar 0000-0001-8245-4804.

S-Editor: Qu XL

L-Editor: A

P-Editor: Zhao YQ

REFERENCES

- 1 Provost F, Van der Woerd J, Malet J, Maggi A, Klinger Y, Michéa D, Pointal E, Pacini F. Mapping the ruptures of the Mw7.8 and Mw7.7 Turkey-Syria Earthquakes using optical offset tracking with Sentinel-2 images. *EGU General Assembly 2023*; 2023 Apr 23–28; Vienna, Austria. EGU23-17612 [DOI: 10.5194/egusphere-egu23-17612]
- 2 Naddaf M. One MRI for 4.7 million people: the battle to treat Syria's earthquake survivors. *Nature* 2023; **615**: 193-194 [PMID: 36849818 DOI: 10.1038/d41586-023-00547-7]
- 3 Erdemir AG, Yurttutan N, Onur MR, İdilman İS, Öztürk MH, Ertürk ŞM, Çevikol C, Akpınar E. Radiological management and challenges of the twin earthquakes of February 6th. *Emerg Radiol* 2023; **30**: 659-666 [PMID: 37535144 DOI: 10.1007/s10140-023-02162-5]
- 4 Wolfson N, Lerner A, Roshal L. Orthopedics in Disasters. Orthopedic Injuries in Natural Disasters and Mass Casualty Events. Heidelberg: Springer Berlin, 2016 [DOI: 10.1007/978-3-662-48950-5]
- 5 Dong ZH, Yang ZG, Chen TW, Chu ZG, Wang QL, Deng W, Denor JC. Earthquake-related versus non-earthquake-related injuries in spinal injury patients: differentiation with multidetector computed tomography. *Crit Care* 2010; **14**: R236 [PMID: 21190568 DOI: 10.1186/cc9391]
- 6 Bywaters EG, Beall D. Crush Injuries with Impairment of Renal Function. *Br Med J* 1941; **1**: 427-432 [PMID: 20783577 DOI: 10.1136/bmj.1.4185.427]
- 7 Bywaters EG. 50 years on: the crush syndrome. *BMJ* 1990; **301**: 1412-1415 [PMID: 2279155 DOI: 10.1136/bmj.301.6766.1412]
- 8 Dong ZH, Yang ZG, Chu ZG, Chen TW, Bai HL, Shao H, Tang SS, Denor JC. Earthquake-related injuries: evaluation with multidetector computed tomography and digital radiography of 1491 patients. *J Crit Care* 2012; **27**: 103.e1-103.e6 [PMID: 21514091 DOI: 10.1016/j.jcrc.2011.03.007]
- 9 Alpergin BC, Mete EB, Zaimoglu M, Caglar YS, Orhan O, Hasimoglu S, Eroglu U. Common Vertebral Fracture Level After the 2023 Turkey Earthquake: Thoracolumbar Junction - Due to Hyper-Flexed and Fixed Posture - at Triangle of Life Areas. *Turk Neurosurg* 2024; **34**: 485-489 [PMID: 37528724 DOI: 10.5137/1019-5149.JTN.44241-23.1]
- 10 Reinhold M, Audigé L, Schnake KJ, Bellabarba C, Dai LY, Oner FC. AO spine injury classification system: a revision proposal for the thoracic and lumbar spine. *Eur Spine J* 2013; **22**: 2184-2201 [PMID: 23508335 DOI: 10.1007/s00586-013-2738-0]
- 11 Vaccaro AR, Koerner JD, Radcliff KE, Oner FC, Reinhold M, Schnake KJ, Kandziora F, Fehlings MG, Dvorak MF, Aarabi B, Rajasekaran S, Schroeder GD, Kepler CK, Vialle LR. AOSpine subaxial cervical spine injury classification system. *Eur Spine J* 2016; **25**: 2173-2184 [PMID: 25716661 DOI: 10.1007/s00586-015-3831-3]
- 12 Pneumatics SG, Karampinas PK, Triantafilopoulos G, Koufos S, Polyzois V, Vlamis J. Evaluation of TLICS for thoracolumbar fractures. *Eur Spine J* 2016; **25**: 1123-1127 [PMID: 25808483 DOI: 10.1007/s00586-015-3889-y]
- 13 Weng Dennis HH, Tak HH. A Review of Thoracolumbar Spine Fracture Classifications. *JOT* 2011; **1** [DOI: 10.4303/jot/235406]
- 14 Chen R, Song Y, Kong Q, Zhou C, Liu L. Analysis of 78 patients with spinal injuries in the 2008 Sichuan, China, earthquake. *Orthopedics* 2009; **32**: 322 [PMID: 19472966 DOI: 10.3928/01477447-20090501-01]
- 15 Li T, Zhou C, Liu L, Gong Q, Zeng J, Liu H, Kong Q, Song Y. [Analysis of spinal injuries in Wenchuan earthquake]. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi* 2009; **23**: 415-418 [PMID: 19431977]
- 16 Dong ZH, Yang ZG, Chen TW, Feng YC, Wang QL, Chu ZG. Spinal injuries in the Sichuan earthquake. *N Engl J Med* 2009; **361**: 636-637 [PMID: 19657135 DOI: 10.1056/NEJMc0903047]
- 17 Ghabili K, Golzari SE, Salehpour F, Imani T, Bazzazi AM, Ghaffari A, Khanli HM, Tizro P, Taghizade S, Shakouri SK. Spinal injuries in the

- 2012 twin earthquakes, northwest iran. *PLoS Curr* 2013; **5** [PMID: 23568085 DOI: 10.1371/currents.dis.39b14d88c93fe04ef1a2ce180b24f8d1]
- 18 **Giotta Lucifero A**, Bruno N, Luzzi S. Surgical management of thoracolumbar junction fractures: An evidence-based algorithm. *World Neurosurg X* 2023; **17**: 100151 [PMID: 36793355 DOI: 10.1016/j.wnsx.2022.100151]
- 19 **Zheng Z**, Hu W, Tian T, Liu W, Wang X, Gao M, You Y, Zhang X. Dynamic Spine Hyperflexion Is Related to Vertebral Compression Fractures in Postmenopausal Women. *Orthop Surg* 2023; **15**: 111-117 [PMID: 36394084 DOI: 10.1111/os.13473]
- 20 **Li WJ**, Guo LX. Influence of different postures under vertical impact load on thoracolumbar burst fracture. *Med Biol Eng Comput* 2020; **58**: 2725-2736 [PMID: 32880092 DOI: 10.1007/s11517-020-02254-1]
- 21 **Aydin S**, Kazci O, Ece B, Kantarci M. Earthquakes from a radiological perspective: what is demanded from the radiologists, and what can we do? A pictorial review. *Diagn Interv Radiol* 2024; **30**: 30-41 [PMID: 37095695 DOI: 10.4274/dir.2023.232157]
- 22 **Erdemir AG**, İdilman İS, Çifçi GÇ, Yıldız AE, Demirkazık F, Onur MR, Akpınar E, Aydingöz Ü. Imaging in crush injury: a spectrum of findings in survivors of the twin earthquakes on February 6, 2023. *Emerg Radiol* 2023; **30**: 513-523 [PMID: 37270438 DOI: 10.1007/s10140-023-02147-4]
- 23 **Wei Y**, Tian W, Zhang GL, Lv YW, Cui GY. Thoracolumbar kyphosis is associated with compressive vertebral fracture in postmenopausal women. *Osteoporos Int* 2017; **28**: 1925-1929 [PMID: 28251286 DOI: 10.1007/s00198-017-3971-x]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

