World Journal of *Orthopedics*

World J Orthop 2024 November 18; 15(11): 1001-1111





Published by Baishideng Publishing Group Inc

World Journal of Orthopedics

Contents

Monthly Volume 15 Number 11 November 18, 2024

EDITORIAL

- 1001 Adipose-derived stem cells and knee osteoarthritis: New perspectives, old concerns de Sousa EB, Gabbi Filho JPA, Gameiro VS, Baptista LS
- 1007 Cardiotoxicity concerns in total joint arthroplasty Cheng CH, Hao WR, Cheng TH

ORIGINAL ARTICLE

Retrospective Study

1015 Effectiveness of negative pressure wound therapy in complex surgical treatment of necrotizing fasciitis of the upper limb

Lipatov KV, Asatryan A, Melkonyan G, Kazantcev AD, Solov'eva EI, Krivikhin DV, Gorbacheva IV, Cherkasov UE

Observational Study

1023 Artificial intelligence awareness and perceptions among pediatric orthopedic surgeons: A cross-sectional observational study

Alomran AK, Alomar MF, Akhdher AA, Al Qanber AR, Albik AK, Alumran A, Abdulwahab AH

Predictive factors for coronal and sagittal graft extrusion length after using tendon autograft for medial 1036 meniscus reconstruction

Zhu TW, Xiang XX, Li CH, Li RX, Zhang N

1047 Presentation and management outcome of foot drop with tibialis posterior tendon transfer Saaiq M

Basic Study

1056 Impacts of mesenchymal stem cells and hyaluronic acid on inflammatory indicators and antioxidant defense in experimental ankle osteoarthritis

Hagag UI, Halfaya FM, Al-Muzafar HM, Al-Jameel SS, Amin KA, Abou El-Kheir W, Mahdi EA, Hassan GANR, Ahmed OM

1075 Gene expression analysis of cytokines and MMPs in melatonin and rhBMP-2 enhanced bone remodeling Paulini MR, Montarele LF, Pitol DL, Giannocco G, Pereira BF, Buchaim DV, Reis CHB, Buchaim RL, Mardegan Issa JP

CASE REPORT

1088 Ilizarov technique for treatment of a giant aneurysmal bone cyst at the distal femur: A case report Long XY, Sun F, Wang T, Li P, Tian Z, Wu XW

1095 Metagenomic next-generation sequencing may assist diagnosis of osteomyelitis caused by Mycobacterium houstonense: A case report

Lin HY, Tan QH



Conter	World Journal of Orthopedics Its Monthly Volume 15 Number 11 November 18, 2024
1101	Arthroscopic synovectomy for synovial hyperplasia in chronic knee gouty arthritis: A case report <i>Utoyo GA, Calvin C</i>
1109	LETTER TO THE EDITOR Insights and implications from the study on meniscus reconstruction using tendon autograft
	Nguyen PD, Lam TK



Contents

Monthly Volume 15 Number 11 November 18, 2024

ABOUT COVER

Peer Reviewer of World Journal of Orthopedics, Jie Wen, PhD, Assistant Professor, Associate Chief Physician, Department of Pediatric Orthopedics, Hunan Provincial People's Hospital, The First Affiliated Hospital of Hunan Normal University, Changsha 410013, Hunan Province, China. cashwj@qq.com

AIMS AND SCOPE

The primary aim of World Journal of Orthopedics (WJO, World J Orthop) is to provide scholars and readers from various fields of orthopedics with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJO mainly publishes articles reporting research results and findings obtained in the field of orthopedics and covering a wide range of topics including arthroscopy, bone trauma, bone tumors, hand and foot surgery, joint surgery, orthopedic trauma, osteoarthropathy, osteoporosis, pediatric orthopedics, spinal diseases, spine surgery, and sports medicine.

INDEXING/ABSTRACTING

WJO is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for WJO as 2.0; JIF Quartile: Q2. The WJO's CiteScore for 2023 is 3.1.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yn-Qing Zhao; Production Department Director: Xiao-Mei Zheng; Cover Editor: Jin-Lei Wang.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS				
World Journal of Orthopedics	https://www.wjgnet.com/bpg/gerinfo/204				
ISSN	GUIDELINES FOR ETHICS DOCUMENTS				
ISSN 2218-5836 (online)	https://www.wjgnet.com/bpg/GerInfo/287				
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH				
November 18, 2010	https://www.wjgnet.com/bpg/gerinfo/240				
FREQUENCY	PUBLICATION ETHICS				
Monthly	https://www.wjgnet.com/bpg/GerInfo/288				
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT				
Massimiliano Leigheb, Xiao-Jian Ye	https://www.wjgnet.com/bpg/gerinfo/208				
EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF	POLICY OF CO-AUTHORS				
Xin Gu	https://www.wjgnet.com/bpg/GerInfo/310				
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE				
http://www.wjgnet.com/2218-5836/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242				
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS				
November 18, 2024	https://www.wjgnet.com/bpg/GerInfo/239				
COPYRIGHT	ONLINE SUBMISSION				
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com				
PUBLISHING PARTNER	PUBLISHING PARTNER'S OFFICIAL WEBSITE				
The Minimally Invasive Spine Surgery Research Center Of Shanghai Jiaotong University	https://www.shtrhospital.com/zkjs/info_29.aspx?itemid=647				
© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA					

E-mail: office@baishideng.com https://www.wjgnet.com



WJD

World Journal of **Orthopedics**

Submit a Manuscript: https://www.f6publishing.com

World J Orthop 2024 November 18; 15(11): 1036-1046

DOI: 10.5312/wjo.v15.i11.1036

Observational Study

ISSN 2218-5836 (online)

ORIGINAL ARTICLE

Predictive factors for coronal and sagittal graft extrusion length after using tendon autograft for medial meniscus reconstruction

Tian-Wang Zhu, Xian-Xiang Xiang, Chun-Hui Li, Rui-Xin Li, Nan Zhang

Specialty type: Orthopedics

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade C Novelty: Grade B Creativity or Innovation: Grade B Scientific Significance: Grade B

P-Reviewer: Gao Y

Received: July 31, 2024 Revised: August 30, 2024 Accepted: September 9, 2024 Published online: November 18, 2024 Processing time: 107 Days and 2.2

Hours



Tian-Wang Zhu, Xian-Xiang Xiang, Chun-Hui Li, Rui-Xin Li, Department of Sports Medicine, Dalian University Affiliated Xinhua Hospital, Dalian 116021, Liaoning Province, China

Nan Zhang, Department of Spine Surgery, Dalian University Affiliated Xinhua Hospital, Dalian 116021, Liaoning Province, China

Co-corresponding authors: Rui-Xin Li and Nan Zhang.

Corresponding author: Rui-Xin Li, MD, PhD, Associate Chief Physician, Surgeon, Department of Sports Medicine, Dalian University Affiliated Xinhua Hospital, 156 Wansui Road, Shahekou District, Dalian 116021, Liaoning Province, China. m-cc@163.com

Abstract

BACKGROUND

Meniscus extrusion occurs in most elderly individuals and most patients after meniscus allograft transplantation. The risk factors and correlative factors of meniscus extrusion have been extensively studied. However, for using tendon autograft for meniscus reconstruction, both graft type and surgical method are different from those in previous studies on meniscus extrusion.

AIM

To identify predictive factors for coronal and sagittal graft extrusion length after using tendon autograft for medial meniscus reconstruction.

METHODS

Ten patients who underwent medial meniscus reconstruction with tendon autograft were selected for this retrospective observational study. The graft extrusions and potential factors were measured and correlation and regression analyses were performed to analyze their relationships.

RESULTS

The medial graft extrusion correlated with the preoperative bilateral hip-kneeankle angle difference, preoperative Kellgren-Lawrence grade, preoperative relative joint space width, and preoperative bilateral medial edge incline angle difference. The anterior graft correlated with the anterior tunnel edge distance at 1 week after operation. The posterior graft extrusion correlated with the preoperative bilateral hip-knee-ankle angle difference, preoperative relative joint space width, and posterior tunnel edge distance at 1 week after operation. The mean



WJO | https://www.wjgnet.com

graft extrusion correlated with the preoperative bilateral hip-knee-ankle angle difference and preoperative relative joint space width. The preoperative joint space width and anterior and posterior tunnel edge distance at 1 week can be used to predict the medial, anterior, posterior, and mean graft extrusion length.

CONCLUSION

The preoperative joint space width and tunnel position can be used to predict the coronal and sagittal graft extrusion length after using tendon autograft for medial meniscus reconstruction.

Key Words: Meniscus; Extrusion; Reconstruction; Autograft; Graft

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Meniscus extrusion in the natural meniscus and meniscus allograft transplantation have been extensively studied but there are no studies on meniscus extrusion after using tendon for meniscus reconstruction. This study found that the preoperative joint space width and tunnel position at 1 week after operation could be used to predict the coronal and sagittal graft extrusion length at 1 week and 8 mo after using tendon autograft for medial meniscus reconstruction. This knowledge offers information for patient selection and surgical points of using tendon autograft for meniscus reconstruction.

Citation: Zhu TW, Xiang XX, Li CH, Li RX, Zhang N. Predictive factors for coronal and sagittal graft extrusion length after using tendon autograft for medial meniscus reconstruction. *World J Orthop* 2024; 15(11): 1036-1046 URL: https://www.wjgnet.com/2218-5836/full/v15/i11/1036.htm DOI: https://dx.doi.org/10.5312/wjo.v15.i11.1036

INTRODUCTION

After subtotal or total meniscectomy, the onset and progression of osteoarthritis are hard to avoid because of the absence of the meniscus[1]. To delay the process of degeneration of the articular cartilage and restore the function of the meniscus, materials for meniscal replacement, including meniscus allografts and tendon autografts, have been used[2].

Meniscus extrusion occurs in more than 70% of the elderly individuals, and it is considered a risk factor for osteoarthritis[3,4]. The risk factors of natural meniscus extrusion may include female sex, meniscal tear, and higher baseline value of extrusion[5]. Similarly, graft extrusions occur in half or even most meniscus allografts after meniscus allograft transplantation although the relationship between graft extrusion and clinical outcome remains unknown[6]. Studies found that incorrect bone block position, relatively large meniscus allograft, and large osteophyte in peripheral tibial plateau could contribute to graft extrusion[7-9]. For using tendon autograft for meniscus reconstruction, both the graft and surgical method are different from those used for meniscus allograft transplantation[2]. To our knowledge, there are no studies on graft extrusion after using tendon for meniscus reconstruction, thus the potential risk factors of graft extrusion are unclear.

The aim of this study was to explore the potential predictive factors for coronal and sagittal graft extrusion after using tendon autograft for medial meniscus reconstruction. It was hypothesized that factors that are pre-existing and can directly influence the force of the graft can be used to predict the coronal and sagittal graft extrusion length.

MATERIALS AND METHODS

Patients

This retrospective observational study was approved by the Ethics Committee of Dalian University Affiliated Xinhua Hospital (document number: 2024-43-02). Ten patients who underwent medial meniscus reconstruction with tendon autograft between July 2021 and February 2023 were selected for the analysis. The inclusion criteria were medial meniscus reconstruction with tendon autograft, a preoperative anteroposterior knee radiograph, a preoperative anteroposterior lower limb radiograph, a preoperative lateral tibia radiograph, and postoperative magnetic resonance imaging (MRI) images of the knee at 1 week and 8 mo. The exclusion criteria were integrity, resolution, contrast, or position of the bone, bone tunnel, or tendon graft in a radiograph or on MRI images that did not meet the standard required for the accurate measurement in the study. Meniscus reconstructions were performed in selected patients. Indications included meniscus deficiency, pain, and < 50 years of age. Contraindications included irreparable chondral injury, malalignment, ligamentous laxity, arthrofibrosis, or active infection.

MR VB17A (Siemens) with a magnetic field strength of 1.5T was used to acquire proton-density weighted turbo spinecho fat-suppressed coronal and sagittal MRI images and proton-density weighted BLADE fat-suppressed axial MRI images. CXDI NE 213014 (Canon) was used to acquire radiographs.

Zaishideng® WJO | https://www.wjgnet.com



Figure 1 The absolute value of the degree of graft extrusion, measured as the horizontal distance from the most peripheral aspect of the graft to the edge of the tibia measured on the slice that the medial joint space is the narrowest on coronal and sagittal knee magnetic resonance imaging. A: Medial extrusion. a: The absolute value of medial extrusion; b: Meniscus remnant; c: Tendon graft; B: Anterior and posterior extrusiona. a: The absolute value of anterior extrusion; b: Tendon graft; c: Meniscus remnant; d: The absolute value of posterior extrusion; e: Tendon graft; f: Meniscus remnant.

Surgical technique

All surgeries were performed by a senior surgeon. The tendon autograft was harvested and threaded. The anatomic anterior and posterior meniscus root tunnels for medial meniscus reconstruction were drilled. Subtotal or total meniscectomy was carried out. The corresponding part of the joint capsule or remnant meniscus was debrided to obtain a bleeding surface. The prepared tendon was pulled into the tibial tunnels and sutured to the joint capsule or meniscus remnant. The tension of the tendon was adjusted and the threads were fixed.

Evaluation of MRI images and radiographs

The x and y coordinates of the anchor points for the measurement were accurate to at least 1 pixel. The blind method was used for the evaluation.

The slices where the joint space width was the narrowest on postoperative coronal and sagittal knee MRI images were used to measure the medial extrusion and anterior and posterior extrusion (Figure 1). The absolute value of the degree of graft extrusion was measured as the horizontal distance from the most peripheral aspect of the graft to the edge of the tibia. The mean extrusion was calculated as the arithmetic mean of the medial, anterior, and posterior extrusion. The relative posterior graft position ratio was calculated by dividing the value of subtracting the anterior extrusion from the posterior extrusion by the value of adding the posterior anterior extrusion and the anterior extrusion.

 $relative \ posterior \ graft \ position \ ratio = \frac{posterior \ extrusion - anterior \ extrusion}{posterior \ extrusion + anterior \ extrusion}$

The slices where the anterior and posterior meniscus root tunnel entrances were the widest on the sagittal knee MRI images at 1 week after operation were used to measure the anterior and posterior tunnel edge distance, respectively (Figure 2). The tunnel edge distance was measured as the horizontal distance from the intraarticular tunnel entrance to the edge of the tibia.

The preoperative anteroposterior knee radiograph was used to measure the medial edge incline angle and absolute joint space width and assess the Kellgren-Lawrence grade (Figure 3). The medial edge incline angle was measured as the degree of the incline angle formed by the common tangent to the medial edge of the femur and tibia and the vertical line. If the medial edge of the femur was medial to that of the tibia, the value was determined as positive and *vice versa*. The bilateral medial edge incline angle difference was calculated by subtracting the contralateral medial edge incline angle from the ipsilateral medial edge incline angle.

bilateral medial edge incline angle difference

- = ipsilateral medial edge incline angle
- contralateral medial edge incline angle

The absolute joint space width was measured as the length of the vertical line segment from the medial edge of the tibial plateau to the femoral condyle. The relative joint space width was calculated by dividing the ipsilateral absolute joint space width by the contralateral absolute joint space width.

Raishideng® WJO | https://www.wjgnet.com



Figure 2 Tunnel edge distance, measured as the horizontal distance from the intraarticular tunnel entrance to the edge of the tibia measured on the slice where the tunnel entrance is the widest on sagittal knee magnetic resonance imaging. A: Anterior tunnel edge distance (indicated by the letter "a"); B: Posterior tunnel edge distance (indicated by the letter "a").



Figure 3 Medial edge incline angle and medial joint space measured on anteroposterior knee radiograph. a: Medial edge incline angle, measured as the degree of the incline angle formed by the common tangent to the medial edge of the femur and tibia and the vertical line; b: Medial joint space, measured as the length of the vertical line segment from the medial edge of the tibia plateau to the femoral condyle.

 $relative \ joint \ space \ width = \frac{ipsilateral \ absolute \ joint \ space \ width}{contralateral \ absolute \ joint \ space \ width}$

The preoperative anteroposterior lower limb radiograph was used to measure the hip-knee-ankle angle, which was measured as the degree of the medial angle formed by the line from the hip center to the knee center and the line from the knee center to the ankle center. The center of the femoral head was regarded as the hip center. The center of the bases of the tibial spines was regarded as the knee center. The center of the talus was regarded as the ankle center. The bilateral hip-knee-ankle angle difference was calculated by subtracting the ipsilateral hip-knee-ankle angle from the contralateral hip-knee-ankle angle.

Baisbideng® WJO https://www.wjgnet.com

Zhu TW et al. Predicting graft extrusion after meniscus reconstruction

bilateral hip-knee-ankle angle difference

= contralateral hip-knee-ankle angle

- ipsilateral hip-knee-ankle angle

The preoperative lateral tibia radiograph was used to measure the medial posterior tibial slope, which was measured as the degree of the posterior inferior angle formed by the line from the anterior edge to the posterior edge of the medial tibia plateau and the long axis of the tibia.

Statistical analysis

GraphPad Prism 9 was used for all statistical analyses. For the description of the demographics, the mean and standard deviation were calculated for measurement data and the number of each portion was calculated for enumeration and ranked data. The correlations between the graft extrusions and the potential factors, between the sagittal graft extrusions and the tunnel edge distances, and between the relative posterior graft position ratio and the preoperative medial posterior tibial slope were computed. Two patients were excluded from the correlation analysis between the relative posterior graft position ratio and the preoperative medial posterior tibial slope because of the additional anterior closing wedge high tibial osteotomy. The factors that are preexisting and can directly influence the force of the graft were selected for the regressions. The regressions between the graft extrusions and the potential factors were computed by simple linear regression and least squares multiple linear regression to explore good predictive models for the medial, anterior, posterior, and mean extrusions. If the *P* value of the analysis of variance ≤ 0.05 , the *R* squared value ≥ 0.6 , and the variance inflation factor of the multicollinearity \leq 5, the model was regarded as a good predictive model for graft extrusion.

RESULTS

Demographics

The medial, anterior, posterior, and mean extrusion of the tendon grafts at 1 week after operation were 1.00 ± 1.90 , $-3.42 \pm$ 2.34, 0.13 ± 2.90 , and -0.76 ± 2.05 mm, respectively. The medial, anterior, posterior, and mean extrusions of the tendon grafts at 8 mo after operation were 1.73 ± 2.38 , 2.90 ± 2.39 , 0.07 ± 2.72 , and 0.49 ± 0.76 mm, respectively (Table 1).

Correlations between graft extrusions and potential factors

The medial extrusion at 1 week after operation correlated with the preoperative Kellgren-Lawrence grade, preoperative bilateral hip-knee-ankle angle difference, preoperative absolute joint space width, preoperative relative joint space width, preoperative bilateral medial edge incline angle difference, anterior extrusion, and posterior extrusion. The medial extrusion at 8 mo after operation correlated with the preoperative Kellgren-Lawrence grade, preoperative hip-knee-ankle angle, preoperative bilateral hip-knee-ankle angle difference, preoperative relative joint space width, preoperative bilateral medial edge incline angle difference, and posterior extrusion (Table 2). The anterior extrusion at 1 week after operation correlated with the medial extrusion and anterior tunnel edge distance. The anterior extrusion at 8 mo after operation correlated with the anterior tunnel edge distance (Tables 2 and 3). The posterior extrusion at 1 week after operation correlated with the preoperative bilateral hip-knee-ankle angle difference, preoperative absolute joint space width, preoperative relative joint space width, preoperative bilateral medial edge incline angle difference, medial extrusion, and posterior tunnel edge distance. The posterior extrusion at 8 mo after operation correlated with the preoperative bilateral hip-knee-ankle angle difference, preoperative relative joint space width, medial extrusion, and posterior tunnel edge distance (Tables 2 and 3). The mean extrusion at 1 week after operation correlated with the preoperative bilateral hip-knee-ankle angle difference, preoperative relative joint space width, medial extrusion, anterior extrusion, and posterior extrusion. The mean extrusion at 8 mo after operation correlated with the preoperative bilateral hip-knee-ankle angle difference, preoperative relative joint space width, medial extrusion, and posterior extrusion. The relative posterior graft position ratio at 1 week after operation correlated with the preoperative medial posterior tibial slope (Table 4).

Regressions between graft extrusions and potential factors

The preoperative relative joint space width could be used to predict the medial and mean extrusion length at 1 week and 8 mo after operation. The anterior tunnel edge distance at 1 week after operation could be used to predict the anterior extrusion length at 1 week and 8 mo after operation. The preoperative relative joint space and the posterior tunnel edge distance could be used to predict the posterior extrusion length at 1 week and 8 mo after operation (Table 5).

DISCUSSION

The most important finding of the study was that the preoperative joint space width and tunnel position at 1 week after operation could be used to predict the coronal and sagittal graft extrusion length at 1 week and 8 mo after using tendon autograft for medial meniscus reconstruction. The hypothesis of the study was confirmed. Additionally, the medial, anterior, posterior, or mean extrusion at 1 week or 8 mo correlated with the preoperative Kellgren-Lawrence grade; preoperative hip-knee-ankle angle or bilateral hip-knee-ankle angle difference; preoperative absolute or relative joint



WJO | https://www.wjgnet.com

Table 1 Demographics	
Variable	Description
Age (year)	36.10 ± 4.98
Gender, male/female	6/4
Body mass index	28.30 ± 4.80
Preoperative medial posterior tibial slope (°)	14.42 ± 2.39
Preoperative Kellgren-Lawrence grade, grade 0/grade 1/grade 2/grade 3	2/3/2/3
Preoperative hip-knee-ankle angle (°)	175.40 ± 3.86
Preoperative bilateral hip-knee-ankle angle difference (°)	2.13 ± 3.00
Preoperative absolute joint space width (mm)	3.09 ± 0.77
Preoperative relative joint space width	0.68 ± 0.12
Preoperative medial edge incline angle (°)	6.68 ± 9.68
Preoperative bilateral medial edge incline angle difference (°)	4.88 ± 7.87
Anterior tunnel edge distance (mm)	5.93 ± 3.42
Posterior tunnel edge distance (mm)	11.23 ± 6.15
1 wk after operation	
Medial extrusion (mm)	1.00 ± 1.90
Anterior extrusion (mm)	-3.46 ± 2.29
Posterior extrusion (mm)	0.13 ± 2.90
Mean extrusion (mm)	-0.76 ± 2.05
Relative posterior graft position ratio	-0.08 ± 1.09
8 mo after operation	
Medial extrusion (mm)	1.73 ± 2.38
Anterior extrusion (mm)	-2.90 ± 2.39
Posterior extrusion (mm)	1.38 ± 4.43
Mean extrusion (mm)	0.07 ± 2.72
Relative posterior graft position ratio	0.49 ± 0.76

Measurement data are given as the mean ± SD. Enumeration and ranked data are given as numbers of each portion.

space width; preoperative bilateral medial edge incline angle difference; or anterior or posterior tunnel edge distance at 1 week after operation. This knowledge offers information for patient selection and surgical points of using tendon autograft for meniscus reconstruction.

The joint space width reflects the ability of joint space to accommodate the meniscus because of the direct contacts between the meniscus and femoral condyle and tibial plateau. A study compared the preoperative absolute and relative joint space width between patients with graft extrusion (\geq 3 mm) and graft nonextrusion (< 3 mm) at 1 year after meniscal allograft transplantation. The study found no difference in joint space width between patients with graft extrusion and those with graft nonextrusion[10]. However, this study found that the preoperative joint space width can be used to predict medial, posterior, and mean graft extrusion at 1 week and 8 mo after operation. There are several possible reasons for this difference. First, the tendon autograft is softer than the meniscus allograft and it is therefore more susceptible to the contact area of femoral condyle and tibial plateau. In a study using semitendinosus tendon for meniscus reconstruction in rabbits, the tendon autografts might need 6 weeks to generate as many chondrocytes as the natural meniscus. For the hardness, the time might be 24 weeks[11]. Considering that the life cycle of rabbits is shorter than that of humans, it may take more time for tendon autograft in humans to construct the mechanical structure like the natural meniscus so as to achieve adequate hardness and avoid extrusion under the pressure of the femoral condyle and tibial plateau. In this study, the observation that general extrusion at 8 mo after operation was greater than that at 1 week after operation also demonstrated that enough time is needed for the tendon to be remodeled into the meniscus. Second, the joint space width was measured as the length of the vertical line segment from the medial edge of the tibial plateau to the femoral condyle in this study in order to ensure that the measure position was close to the graft midbody, since the measurement method of joint space width varied across studies. Third, that study did not differentiate the medial meniscus and lateral meniscus in study design, while our study concentrated on the medial meniscus.

Baishidena® WJO | https://www.wjgnet.com

Table 2 Correlations between graft extrusions and potential factors

	Medial extrusion		Anterior extrusion		Posterior extrusion		Mean extrusion	
	Pearson r	P value	Pearson r	P value	Pearson <i>r</i>	P value	Pearson r	P value
1 wk after operation								
Preoperative medial posterior tibial slope	0.259	0.575	0.375	0.360	0.381	0.399	0.362	0.425
Preoperative Kellgren-Lawrence grade	0.646	0.044 ^a	0.125	0.732	0.543	0.105	0.488	0.152
Preoperative hip-knee-ankle angle	-0.580	0.079	-0.127	0.727	-0.625	0.053	-0.518	0.125
Preoperative bilateral hip-knee-ankle angle difference	0.736	0.015 ^a	0.272	0.448	0.725	0.018 ^a	0.661	0.038 ^a
Preoperative absolute joint space width	-0.672	0.034 ^a	-0.218	0.545	-0.668	0.035 ^a	-0.591	0.072
Preoperative relative joint space width	-0.934	0.000 ^a	-0.575	0.082	-0.795	0.006 ^a	-0.873	0.001 ^a
Preoperative medial edge incline angle	0.332	0.349	-0.006	0.987	0.455	0.186	0.311	0.382
Preoperative bilateral medial edge incline angle difference	0.678	0.031 ^a	0.133	0.713	0.756	0.011 ^a	0.604	0.064
Medial extrusion			0.711	0.021 ^a	0.733	0.016 ^a	0.914	0.000 ^a
Anterior extrusion	0.711	0.021 ^a			0.480	0.160	0.818	0.004 ^a
Posterior extrusion	0.733	0.016 ^a	0.480	0.160			0.872	0.001 ^a
8 mo after operation								
Preoperative medial posterior tibial slope	0.643	0.357	-0.265	0.735	0.469	0.531	0.415	0.585
Preoperative Kellgren-Lawrence grade	0.832	0.040 ^a	0.142	0.789	0.575	0.233	0.596	0.212
Preoperative hip-knee-ankle angle	-0.890	0.018 ^a	-0.142	0.789	-0.799	0.056	-0.735	0.096
Preoperative bilateral hip-knee-ankle angle difference	0.962	0.002 ^a	0.398	0.434	0.867	0.025 ^a	0.868	0.025 ^a
Preoperative absolute joint space width	-0.723	0.105	-0.208	0.692	-0.675	0.142	-0.638	0.173
Preoperative relative joint space width	-0.940	0.005 ^a	-0.266	0.610	-0.887	0.018 ^a	-0.834	0.039 ^a
Preoperative medial edge incline angle	0.509	0.302	-0.398	0.434	0.229	0.662	0.156	0.767
Preoperative bilateral medial edge incline angle difference	0.848	0.033 ^a	-0.052	0.921	0.625	0.185	0.571	0.236
Medial extrusion			0.324	0.531	0.844	0.035 ^a	0.854	0.031 ^a
Anterior extrusion	0.324	0.531			0.614	0.194	0.721	0.106
Posterior extrusion	0.844	0.035 ^a	0.614	0.194			0.985	0.000 ^a

^aP value ≤ 0.05 .

Table 3 Correlations between sagittal graft extrusions and tunnel edge distances

	Anterior extrusion		Posterior extrusion		
	Pearson <i>r</i>	<i>P</i> value	Pearson <i>r</i>	P value	
1 wk after operation					
Anterior tunnel edge distance	-0.753	0.012 ^a	-0.151	0.676	
Posterior tunnel edge distance	-0.349	0.323	-0.658	0.039 ^a	
8 mo after operation					
Anterior tunnel edge distance	-0.910	0.012 ^a	-0.634	0.177	
Posterior tunnel edge distance	-0.728	0.101	-0.942	0.005 ^a	

^aP value ≤ 0.05 .

Jaishideng® WJO | https://www.wjgnet.com

November 18, 2024 Volume 15 Issue 11

Table 4 Correlations between relative posterior graft position ratio and preoperative medial posterior tibial slope							
	Relative posterior graft position ratio						
	Pearson <i>r</i>	<i>P</i> value					
1 wk after operation							
Preoperative medial posterior tibial slope	0.800	0.031 ^a					
8 mo after operation							
Preoperative medial posterior tibial slope	0.147	0.854					

^aP value ≤ 0.05 .

Two patients were excluded from the analysis because of the additional anterior closing wedge high tibial osteotomy.

Table 5 Regressions between graft extrusions and potential factors

			<i>P</i> value					
v	Standardized	Equation		X ₁	X ₁	x ₁	- R ²	
y	otandardized	Equation	Regression	Preoperative relative joint space width	Anterior tunnel edge distance	Posterior tunnel edge distance		
1 wk after operation								
Medial	Yes	$y = -14.6x_1 + 68.9$	0.000	0.000			0.873	
extrusion	No	$y = -0.0604x_1 + 0.737$						
Anterior	Yes	$y = -9.755x_2 + 5.77$	0.006		0.006		0.632	
extrusion	No	y = -1.185x ₂ + 1.831						
Posterior extrusion	Yes	$y = -0.6192x_1 - 0.3481x_3 + 86.65$	0.009	0.024		0.132	0.741	
	No	$y = -14.75x_1 - 0.1744x_3 + 12.07$						
Mean	Yes	$y = -12.6x_1 + 44.6$	0.001	0.001			0.763	
extrusion	No	$y = -0.0523x_1 + 0.63$						
8 mo after operation								
Medial extrusion	Yes	y = -15.13x ₁ + 88.34	0.005	0.005			0.884	
	No	$y = -0.04563x_1 + 0.7183$						
Anterior	Yes	$y = -16.80x_2 - 6.641$	0.012		0.012		0.827	
extrusion	No	$y = -0.9522x_2 + 2.994$						
Posterior extrusion	Yes	$y = -0.3597x_1 - 0.6639x_3 + 99.15$	0.016	0.232		0.078	0.936	
	No	$y = -14.09x_1 - 0.5704x_3 + 17.02$						
Mean extrusion	Yes	$y = -11.74x_1 + 62.93$	0.039	0.039			0.696	
	No	$y = -0.0354x_1 + 0.6417$						

Jaishideng® WJO | https://www.wjgnet.com

The position of the meniscus root may affect meniscus extrusion because of the action and reaction force between adjacent tissues. The ratio of the distance between the edge of the lateral tibial plateau and the center of the bone bridge to the tibial plateau length and the distance between the interspinal groove and the middle of the bone bridge were demonstrated to correlate with the coronal extrusion after lateral meniscus transplantation[12]. Similarly, the axial trough angle, the angle between the long axis of the bone trough and the line perpendicular to the common tangent to the medial and lateral posterior tibial condylar cortexes on axial MRI images, was demonstrated to correlate with the coronal extrusion after lateral meniscus transplantation [13,14]. For medial meniscus transplantation, the distance between the anterior bone plug and the line perpendicular to the common tangent to the medial and lateral posterior tibial condylar cortexes and passing through the posterior cruciate ligament on axial MRI images was found to be associated with the coronal extrusion[15]. All these studies found that the closer the meniscus root is to the ipsilateral edge of the tibial plateau, the greater the likelihood that the coronal extrusion will occur. However, when it comes to using tendon for meniscus reconstruction, the intraarticular length of the tendon graft can be adjusted after the bone tunnels are drilled, hence the tunnel position hardly affects the magnitude but rather the direction of the force on the graft. Therefore, the relationship between the coronal tunnel position and the coronal graft extrusion was not investigated in this study. In addition, a study assessed the ratio of the anteroposterior distance between the tunnel center and the anatomic center to the anteroposterior length of the tibia at 1 week after medial meniscus posterior root pullout repair and posterior meniscus extrusion at 3 mo after operation, and no correlation was found[16]. However, the present study found that the tunnel edge distance, measured as the anteroposterior distance from the intraarticular tunnel entrance to the edge of the tibia, can be used to predict anterior and posterior graft extrusion at 1 week and 8 mo after operation. The possible reason for this difference is that the tendon autograft is softer than the meniscus allograft and its sagittal extrusion is therefore more susceptible to its sagittal tunnel position.

The hip-knee-ankle angle, the reflection of lower limb alignment, may influence meniscus extrusion because it determines the distribution of the load between medial and lateral knee joint. A study evaluated the hip-knee-ankle angle and coronal extrusion of the medial meniscus within 3 mo from the first consultation in those suspected of having knee disease or injury. A correlation between the hip-knee-ankle angle and meniscus extrusion was demonstrated [17]. Similarly, this study found that the preoperative bilateral hip-knee-ankle angle difference correlated with the medial, posterior, and mean graft extrusions at 1 week and 8 mo after operation. At first glance, the hip-knee-ankle angle is also a good factor to be chosen for the predictive model for graft extrusion in this study, since it correlates with the extrusion of both the natural meniscus and tendon graft. However, considering that the joint space can more directly influence the force of the graft, the hip-knee-ankle was not chosen.

The medial edge incline angle, measured as the degree of the incline angle formed by the common tangent to the medial edge of the femur and tibia and the vertical line, was used to reflect the relationship between the medial femoral condyle and the medial tibial plateau in the mediolateral direction. To our knowledge, there are no studies on this relationship in the field of meniscus extrusion. This study found that the preoperative bilateral medial edge incline angle difference correlated with the medial and posterior graft extrusion at 1 week and the medial graft extrusion at 8 mo after operation. Because the function that medial meniscus restricts lateral tibial translation, which is equal to medial femoral translation, implies that medial femoral condyle exerts a force toward the medial side on the medial meniscus, medial femoral translation may contribute to medial graft extrusion[18]. Since the posterior graft extrusion was measured on the slice that the joint space width was the narrowest of the sagittal knee on MRI images, the medial femoral translation made the corresponding tibial plateau at the relative medial portion of the medial tibial plateau. In this condition, the anteroposterior length of the tibial plateau might decrease, and the posterior graft extrusion might increase. As a result, the medial femoral translation might contribute to the posterior graft extrusion.

Kellgren-Lawrence grade has been extensively used to assess knee arthrosis according to bone morphology. Kellgren-Lawrence grade was found to correlate with the coronal extrusion of the medial meniscus in both early knee osteoarthritis and medial meniscus posterior horn tear [19,20]. Two studies analyzed the preoperative Kellgren-Lawrence grade and the coronal extrusion of the lateral meniscus at 1 year following meniscus allograft transplantation. One failed to find an association between them[21]. The other found that among the knees with grades 0, 1, and 2, the meniscus extrusion was more likely to occur in the knees with grade 1 compared with those with grade 0[14]. In this study, the preoperative Kellgren-Lawrence grade correlated with the medial graft extrusion at 1 week and 8 mo after operation. Since the tendon autograft is softer than the meniscus allograft and Kellgren-Lawrence grade involves joint space, which determined the degree of graft extrusion in this study, tendon autograft may be more susceptible to bone morphology than meniscus allograft.

The posterior tibial slope can influence the force on the femoral condyle and meniscus. A study found that the knees with a posterior tibial slope of 15° exerted greater force on the medial meniscus posterior roots with the joints loaded in compression compared with those with a posterior tibial slope of 10°[22]. Similarly, this study found that the preoperative posterior tibial slope correlated with the relative posterior graft position ratio at 1 week after operation.

There may be a relationship among medial, anterior, and posterior extrusions of the meniscus since they are adjacent tissues. However, a study found no correlation between the medial, anterior, and posterior extrusions of the medial meniscus on immediate postoperative MRI images after meniscus allograft transplantation[23]. Nevertheless, this study found that the medial extrusion correlated with the posterior extrusion at 1 week and 8 mo after operation. In addition, the medial extrusion correlated with the anterior extrusion at 1 week after operation. Interestingly, the correlation between the medial extrusion and posterior extrusion was stronger than that between the medial extrusion and anterior extrusion. Similarly, the most important risk factor for the medial and posterior extrusions was the preoperative joint space width while the most important risk factor for the anterior extrusion was the anterior tunnel edge distance in this study. The anatomical characteristic of the medial meniscus that anterior root attachment is anterior to the center of the anterior horn and posterior root attachment is anterior to the center of the posterior horn helps to explain these findings.

WJO https://www.wjgnet.com

In theory, the characteristic determines that radially inward tension and radially outward compression on the anterior horn are weaker than those on the posterior horn, which has been confirmed in a previous study [24]. In theory, the characteristic also determines that the influence of the position of anterior root attachment on anterior extrusion is stronger than that of posterior root attachment on posterior extrusion since the resistance of the femoral condyle is weaker.

This study had several strengths. First, the longitudinal study in which the images used to measure almost all the correlative factors were acquired ahead of the images used to measure the graft extrusions was helpful to explore the causal link while most studies on meniscus extrusion were cross-sectional. Second, this study not only found that many factors correlated with the coronal and sagittal graft extrusion, but also proposed models with good fitness that can be used to predict the medial, anterior, posterior, and mean extrusion length since no studies on graft extrusion after using tendon autograft for meniscus reconstruction could be found. Third, to our knowledge, the sample size was the largest among clinical studies on using tendon autograft for meniscus reconstruction since other studies included no more than several patients.

This study had several limitations. First, the sample size was small, which might contribute to random errors. Second, the respective slices where the joint space width was the narrowest on coronal and sagittal knee MRI images were used to measure the medial extrusion and anterior and posterior extrusions, respectively. The method that using graft extrusion measured on a particular slice to represent graft extrusion might contribute to systematic errors. Third, there might be some correlative factors or risk factors for graft extrusion that this study neglected.

CONCLUSION

The preoperative joint space width and tunnel position can be used to predict the coronal and sagittal graft extrusion length after using tendon autograft for medial meniscus reconstruction. This knowledge offers information for patient selection and surgical points of using tendon autograft for meniscus reconstruction.

FOOTNOTES

Author contributions: Zhu TW designed and conducted the study and wrote the paper; Xiang XX and Li CH contributed to the analysis; Li RX and Zhang N supervised the study.

Institutional review board statement: This study was reviewed and approved by the Ethics Committee of Dalian University Affiliated Xinhua Hospital (document number: 2024-43-02).

Informed consent statement: Patients were not required to give informed consent to the study because the analysis used anonymous clinical data that were obtained after each patient agreed to treatment by written consent.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: No additional data are available.

STROBE statement: The authors have read the STROBE Statement-checklist of items-and the manuscript was prepared and revised according to the STROBE Statement-checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country of origin: China

ORCID number: Tian-Wang Zhu 0009-0004-0941-2316; Xian-Xiang Xiang 0000-0001-8184-4878; Chun-Hui Li 0009-0000-4039-0320; Rui-Xin Li 0009-0007-8942-9063; Nan Zhang 0000-0002-6441-8022.

S-Editor: Qu XL L-Editor: Wang TQ P-Editor: Yu HG

REFERENCES

Papalia R, Del Buono A, Osti L, Denaro V, Maffulli N. Meniscectomy as a risk factor for knee osteoarthritis: a systematic review. Br Med Bull 2011; 99: 89-106 [PMID: 21247936 DOI: 10.1093/bmb/ldq043]



- Goble EM, Verdonk R, Kohn D. Arthroscopic and open surgical techniques for meniscus replacement--meniscal allograft transplantation and 2 tendon autograft transplantation. Scand J Med Sci Sports 1999; 9: 168-176 [PMID: 10380275 DOI: 10.1111/j.1600-0838.1999.tb00449.x]
- 3 Kaneko H, Ishijima M, Aoki T, Negishi Y, Liu L, Arita H, Momoeda M, Arepati A, Ji X, Someya Y, Tamura Y, Watada Y, Kawamori R, Kaneko K. Prevalence of medial meniscus extrusion in elderly persons -the bunkyo health study. Osteoarthritis Cartilage 2020; 28: S295 [DOI: 10.1016/j.joca.2020.02.464]
- Emmanuel K, Quinn E, Niu J, Guermazi A, Roemer F, Wirth W, Eckstein F, Felson D. Quantitative measures of meniscus extrusion predict 4 incident radiographic knee osteoarthritis--data from the Osteoarthritis Initiative. Osteoarthritis Cartilage 2016; 24: 262-269 [PMID: 26318658 DOI: 10.1016/j.joca.2015.08.003]
- Zhang F, Kumm J, Svensson F, Turkiewicz A, Frobell R, Englund M. Risk factors for meniscal body extrusion on MRI in subjects free of 5 radiographic knee osteoarthritis: longitudinal data from the Osteoarthritis Initiative. Osteoarthritis Cartilage 2016; 24: 801-806 [PMID: 26706701 DOI: 10.1016/j.joca.2015.12.003]
- 6 Lee DH. Incidence and Extent of Graft Extrusion following Meniscus Allograft Transplantation. Biomed Res Int 2018; 2018: 5251910 [PMID: 29770332 DOI: 10.1155/2018/5251910]
- Lee SY, Yoon J, Cho YS, Son RC, Kim SK, Ahn SH, Lee HK. Various factors contribute to graft extrusion in lateral meniscus allograft 7 transplantation. Knee Surg Sports Traumatol Arthrosc 2017; 25: 374-382 [PMID: 26607944 DOI: 10.1007/s00167-015-3882-3]
- Jang SH, Kim JG, Ha JG, Shim JC. Reducing the size of the meniscal allograft decreases the percentage of extrusion after meniscal allograft 8 transplantation. Arthroscopy 2011; 27: 914-922 [PMID: 21693346 DOI: 10.1016/j.arthro.2011.02.017]
- 9 Jeon B, Kim JM, Kim JM, Lee CR, Kim KA, Bin SI. An osteophyte in the tibial plateau is a risk factor for allograft extrusion after meniscus allograft transplantation. Am J Sports Med 2015; 43: 1215-1221 [PMID: 25690626 DOI: 10.1177/0363546515570462]
- 10 Lee SM, Bin SI, Kim JM, Lee BS, Lee CR, Son DW, Park JG. Long-term Outcomes of Meniscal Allograft Transplantation With and Without Extrusion: Mean 12.3-Year Follow-up Study. Am J Sports Med 2019; 47: 815-821 [PMID: 30763116 DOI: 10.1177/0363546518825251]
- 11 Li C, Hu X, Meng Q, Zhang X, Zhu J, Dai L, Cheng J, Zhong M, Shi W, Ren B, Zhang J, Fu X, Duan X, Ao Y. The potential of using semitendinosus tendon as autograft in rabbit meniscus reconstruction. Sci Rep 2017; 7: 7033 [PMID: 28765605 DOI: 10.1038/s41598-017-07166-z]
- Choi NH, Yoo SY, Victoroff BN. Position of the bony bridge of lateral meniscal transplants can affect meniscal extrusion. Am J Sports Med 12 2011; 39: 1955-1959 [PMID: 21642600 DOI: 10.1177/0363546511407614]
- Lee DH, Kim JM, Lee BS, Kim KA, Bin SI. Greater axial trough obliquity increases the risk of graft extrusion in lateral meniscus allograft 13 transplantation. Am J Sports Med 2012; 40: 1597-1605 [PMID: 22634211 DOI: 10.1177/0363546512447932]
- Ahn JH, Kang HW, Yang TY, Lee JY. Multivariate Analysis of Risk Factors of Graft Extrusion After Lateral Meniscus Allograft 14 Transplantation. Arthroscopy 2016; 32: 1337-1345 [PMID: 26970835 DOI: 10.1016/j.arthro.2015.12.050]
- 15 Yoon KH, Lee HW, Park SY, Kim JS, Park JY. Centralized anterior bone plug results in less graft extrusion in patients undergoing medial meniscus allograft transplantation following anterior cruciate ligament reconstruction. Knee 2020; 27: 884-890 [PMID: 32563438 DOI: 10.1016/j.knee.2020.02.014]
- Kamatsuki Y, Furumatsu T, Hiranaka T, Okazaki Y, Okazaki Y, Kodama Y, Hino T, Masuda S, Miyazawa S, Ozaki T. Placement of an 16 anatomic tibial tunnel significantly improves the medial meniscus posterior extrusion at 90° of knee flexion following medial meniscus posterior root pullout repair. Knee Surg Sports Traumatol Arthrosc 2021; 29: 1025-1034 [PMID: 32451621 DOI: 10.1007/s00167-020-06070-w]
- Goto N, Okazaki K, Akiyama T, Akasaki Y, Mizu-Uchi H, Hamai S, Nakamura S, Nakashima Y. Alignment factors affecting the medial 17 meniscus extrusion increases the risk of osteoarthritis development. Knee Surg Sports Traumatol Arthrosc 2019; 27: 2617-2623 [PMID: 30456569 DOI: 10.1007/s00167-018-5286-7]
- Zhang Y, Huang W, Yao Z, Ma L, Lin Z, Wang S, Huang H. Anterior Cruciate Ligament Injuries Alter the Kinematics of Knees With or 18 Without Meniscal Deficiency. Am J Sports Med 2016; 44: 3132-3139 [PMID: 27511793 DOI: 10.1177/0363546516658026]
- 19 Sekiya I, Katano H, Guermazi A, Miura Y, Okanouchi N, Tomita M, Masumoto J, Kitazume Y, Koga H, Ozeki N. Association of AIdetermined Kellgren-Lawrence grade with medial meniscus extrusion and cartilage thickness by AI-based 3D MRI analysis in early knee osteoarthritis. Sci Rep 2023; 13: 20093 [PMID: 37973855 DOI: 10.1038/s41598-023-46953-9]
- 20 Lee DH, Lee BS, Kim JM, Yang KS, Cha EJ, Park JH, Bin SI. Predictors of degenerative medial meniscus extrusion: radial component and knee osteoarthritis. Knee Surg Sports Traumatol Arthrosc 2011; 19: 222-229 [PMID: 20890696 DOI: 10.1007/s00167-010-1274-2]
- Lee BS, Kim JM, Kim JM, Kim KA, Bin SI. Patient-related risk factors for the extrusion of lateral meniscal allograft transplants. Arthroscopy 21 2015; **31**: 699-706 [PMID: 25530512 DOI: 10.1016/j.arthro.2014.10.016]
- Melugin HP, Brown JR, Hollenbeck JFM, Fossum BW, Whalen RJ, Ganokroj P, Provencher CMT. Increased Posterior Tibial Slope Increases 22 Force on the Posterior Medial Meniscus Root. Am J Sports Med 2023; 51: 3197-3203 [PMID: 37715505 DOI: 10.1177/03635465231195841]
- Lee DH, Lee CR, Jeon JH, Kim KA, Bin SI. Graft extrusion in both the coronal and sagittal planes is greater after medial compared with lateral 23 meniscus allograft transplantation but is unrelated to early clinical outcomes. Am J Sports Med 2015; 43: 213-219 [PMID: 25389368 DOI: 10.1177/0363546514555699]
- Walker PS, Arno S, Bell C, Salvadore G, Borukhov I, Oh C. Function of the medial meniscus in force transmission and stability. J Biomech 24 2015; 48: 1383-1388 [PMID: 25888013 DOI: 10.1016/j.jbiomech.2015.02.055]



WJO | https://www.wjgnet.com



Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

