

World Journal of *Gastrointestinal Surgery*

Monthly Volume 17 Number 3 March 27, 2025



EDITORIAL

Fan YH, Wang MW, Gao YN, Li WM, Jiao Y. Genetic and environmental factors influencing Crohn's disease. *World J Gastrointest Surg* 2025; 17(3): 98526 [DOI: [10.4240/wjgs.v17.i3.98526](https://doi.org/10.4240/wjgs.v17.i3.98526)]

Pandey CK, Kumar A. Perioperative neurocognitive dysfunction and role of dexmedetomidine in radical colon cancer surgery in elderly patients. *World J Gastrointest Surg* 2025; 17(3): 100126 [DOI: [10.4240/wjgs.v17.i3.100126](https://doi.org/10.4240/wjgs.v17.i3.100126)]

Wang Y, Xun X, Luan WY, Zhang Z, Xu ZX, Lin SX, Miao YD. Hyperthermia combined with opioid therapy: Enhancing cancer pain management and reducing surgical stress in gastrointestinal cancer patients. *World J Gastrointest Surg* 2025; 17(3): 101060 [DOI: [10.4240/wjgs.v17.i3.101060](https://doi.org/10.4240/wjgs.v17.i3.101060)]

Li LQ, Jiao Y. Risk and management of adverse events in minimally invasive esophagectomy. *World J Gastrointest Surg* 2025; 17(3): 103941 [DOI: [10.4240/wjgs.v17.i3.103941](https://doi.org/10.4240/wjgs.v17.i3.103941)]

MINIREVIEWS

Deng SS, Zhu YP, Chen ZT, Li W. Application progress of early nutrition intervention in patients with hepatocellular carcinoma after liver transplantation. *World J Gastrointest Surg* 2025; 17(3): 100321 [DOI: [10.4240/wjgs.v17.i3.100321](https://doi.org/10.4240/wjgs.v17.i3.100321)]

Feng LF, Li XW, Zhu XQ, Jin LN. Advances in management strategies for enteral nutrition-related gastric retention in adult patients with nasogastric tubes. *World J Gastrointest Surg* 2025; 17(3): 101751 [DOI: [10.4240/wjgs.v17.i3.101751](https://doi.org/10.4240/wjgs.v17.i3.101751)]

Wu L, Wu H, Mu S, Li XY, Zhen YH, Li HY. Surgical approaches for complete rectal prolapse. *World J Gastrointest Surg* 2025; 17(3): 102043 [DOI: [10.4240/wjgs.v17.i3.102043](https://doi.org/10.4240/wjgs.v17.i3.102043)]

Zhang XD, Zhang LY, Luo JL, Yu KH, Zhu KL. Neoadjuvant therapy: Dawn of reducing the high post-surgery recurrence rate of hepatocellular carcinoma. *World J Gastrointest Surg* 2025; 17(3): 103740 [DOI: [10.4240/wjgs.v17.i3.103740](https://doi.org/10.4240/wjgs.v17.i3.103740)]

ORIGINAL ARTICLE

Retrospective Cohort Study

Liu M, Feng B, He N, Yan R, Qin J. Efficacy of fluorouracil combined with paclitaxel and oxaliplatin for the treatment of advanced gastric signet ring cell carcinoma. *World J Gastrointest Surg* 2025; 17(3): 94286 [DOI: [10.4240/wjgs.v17.i3.94286](https://doi.org/10.4240/wjgs.v17.i3.94286)]

Zu QQ, You Y, Chen AZ, Wang XR, Zhang SH, Chen FL, Liu M. Combined application of the preclosure technique and traction approach facilitates endoscopic full-thickness resection of gastric submucosal tumors. *World J Gastrointest Surg* 2025; 17(3): 95704 [DOI: [10.4240/wjgs.v17.i3.95704](https://doi.org/10.4240/wjgs.v17.i3.95704)]

Zhao L, Wei L, Fei XL. Impact of diabetes on recovery after radical gastrectomy for gastric cancer: A retrospective cohort study. *World J Gastrointest Surg* 2025; 17(3): 100763 [DOI: [10.4240/wjgs.v17.i3.100763](https://doi.org/10.4240/wjgs.v17.i3.100763)]

Zhao SQ, Wang SY, Ge N, Guo JT, Liu X, Wang GX, Su L, Sun SY, Wang S. Endoscopic full-thickness resection vs surgical resection for gastric stromal tumors: Efficacy and safety using propensity score matching. *World J Gastrointest Surg* 2025; 17(3): 101002 [DOI: [10.4240/wjgs.v17.i3.101002](https://doi.org/10.4240/wjgs.v17.i3.101002)]

Salehi O, Gao WL, Kenfield C, Hebbard G. Roux-en-Y jejunostomy in gastroparesis: Insight into patient perspectives and outcomes. *World J Gastrointest Surg* 2025; 17(3): 102543 [DOI: 10.4240/wjgs.v17.i3.102543]

Shu Y, Li KJ, Sulayman S, Zhang ZY, Ababaik S, Wang K, Zeng XY, Chen Y, Zhao ZL. Predictive value of serum calcium ion level in patients with colorectal cancer: A retrospective cohort study. *World J Gastrointest Surg* 2025; 17(3): 102638 [DOI: 10.4240/wjgs.v17.i3.102638]

Retrospective Study

Yu Y, Wang XQ, Liu G, Li L, Chen LN, Zhang LJ, Xia Q. Impact of a visual mobile terminal-based continuity of care model on caregiver competence of children with enterostomies. *World J Gastrointest Surg* 2025; 17(3): 99099 [DOI: 10.4240/wjgs.v17.i3.99099]

Chen DX, Fang KX, Chen SX, Hou SL, Wen GH, Yang HK, Shi DP, Lu QX, Zhai YQ, Li MY. Optimal timing of endoscopic biliary drainage for bile duct leaks: A multicenter, retrospective, clinical study. *World J Gastrointest Surg* 2025; 17(3): 99425 [DOI: 10.4240/wjgs.v17.i3.99425]

Liu LN, Chang YF, Wang H. Correlations of three scoring systems with the prognosis of patients with liver cirrhosis complicated with sepsis syndrome. *World J Gastrointest Surg* 2025; 17(3): 99570 [DOI: 10.4240/wjgs.v17.i3.99570]

Lou QX, Xu KP. Analgesic effect and safety of dexmedetomidine-assisted intravenous-inhalation combined general anesthesia in laparoscopic minimally invasive inguinal hernia surgery. *World J Gastrointest Surg* 2025; 17(3): 99597 [DOI: 10.4240/wjgs.v17.i3.99597]

Shi JH, Yang H, Wang ST, Wang WJ, Shi Y, Huang SS, Jiang S. Retrospective analysis on Lou Bei Er Chen decoction and acupuncture in gastroesophageal reflux disease post-gastric cancer surgery. *World J Gastrointest Surg* 2025; 17(3): 99626 [DOI: 10.4240/wjgs.v17.i3.99626]

Fang ZH, Hao AH, Qi YG. Imaging features and correlation with short-term prognosis in laparoscopic radical resection of colorectal cancer. *World J Gastrointest Surg* 2025; 17(3): 99782 [DOI: 10.4240/wjgs.v17.i3.99782]

Li J, Chen JP, Lai CH, Fu L, Ji Y. Efficacy of water infusion combined with defoamers in colonoscopy. *World J Gastrointest Surg* 2025; 17(3): 99784 [DOI: 10.4240/wjgs.v17.i3.99784]

Chen L, Li BX, Gan QZ, Guo RG, Chen X, Shen X, Chen Y. Enhanced recovery after surgery-based evidence-based care plus ice stimulation for thirst management in convalescent patients following digestive surgery under general anesthesia. *World J Gastrointest Surg* 2025; 17(3): 100185 [DOI: 10.4240/wjgs.v17.i3.100185]

Ni WJ, Xi YX, Zhou YC. Efficacy of combined psychological and physical nursing in preventing peripherally inserted central catheter-related thrombosis in gastric cancer patients. *World J Gastrointest Surg* 2025; 17(3): 100430 [DOI: 10.4240/wjgs.v17.i3.100430]

Yang JL, Yang YJ, Xu L. Effect of forearm and posterior wall anastomosis on gastroesophageal reflux in proximal gastrectomy patients. *World J Gastrointest Surg* 2025; 17(3): 100799 [DOI: 10.4240/wjgs.v17.i3.100799]

Li M, Yuan DH, Yang Z, Lu TX, Zhang L. Retrospective analysis of preoperative tumor marker levels in rectal cancer patients: Implications for diagnosis. *World J Gastrointest Surg* 2025; 17(3): 100820 [DOI: 10.4240/wjgs.v17.i3.100820]

Lin YM, Yu C, Xian GZ. Retrospective analysis of delta hemoglobin and bleeding-related risk factors in pancreaticoduodenectomy. *World J Gastrointest Surg* 2025; 17(3): 100999 [DOI: 10.4240/wjgs.v17.i3.100999]

Liu JR, Zhang J, Duan XL. Risk factors influencing sphincter preservation in laparoscopic radical rectal cancer surgery. *World J Gastrointest Surg* 2025; 17(3): 101061 [DOI: 10.4240/wjgs.v17.i3.101061]

Wu PH, Ta ZQ. Clinical effect and prognosis of laparoscopic surgery on colon cancer complicated with intestinal obstruction patients. *World J Gastrointest Surg* 2025; 17(3): 101609 [DOI: 10.4240/wjgs.v17.i3.101609]

Li HS, Zhang XF, Fu J, Yuan B. Efficacy of microwave ablation vs laparoscopic hepatectomy for primary small liver cancer: A comparative study. *World J Gastrointest Surg* 2025; 17(3): 101786 [DOI: 10.4240/wjgs.v17.i3.101786]

Shin DW, Cho YA, Moon SH, Kim TH, Park JW, Lee JW, Choe JY, Kim MJ, Kim SE. High cellular prion protein expression in cholangiocarcinoma: A marker for early postoperative recurrence and unfavorable prognosis. *World J Gastrointest Surg* 2025; 17(3): 101940 [DOI: 10.4240/wjgs.v17.i3.101940]

Yang QS, Zhang M, Ma CS, Teng D, Li A, Dong JD, Wang XF, Liu FB. Analysis of risk factors for bile leakage after laparoscopic exploration and primary suture of common bile duct. *World J Gastrointest Surg* 2025; 17(3): 102190 [DOI: 10.4240/wjgs.v17.i3.102190]

Lu XY, Tan XD. Clinical outcomes of interlocking main pancreatic duct-jejunal internal bridge drainage in middle pancreatectomy: A comparative study. *World J Gastrointest Surg* 2025; 17(3): 102428 [DOI: 10.4240/wjgs.v17.i3.102428]

Xiao NJ, Chu JG, Ning SB, Wei BJ, Xia ZB, Han ZY. Successful management of bleeding ectopic small bowel varices secondary to portal hypertension: A retrospective study. *World J Gastrointest Surg* 2025; 17(3): 102589 [DOI: 10.4240/wjgs.v17.i3.102589]

Hu XS, Wang Y, Pan HT, Zhu C, Zhou S, Chen SL, Liu HC, Pang Q, Jin H. Initial experience with ultrafine choledochoscopy combined with low-dose atropine for the treatment of Oddi intersphincter stones. *World J Gastrointest Surg* 2025; 17(3): 102998 [DOI: 10.4240/wjgs.v17.i3.102998]

Yuan J, Liu Q, Wu BY. Therapeutic effectiveness and influencing factors of laparoscopic appendectomy with mesoappendix dissection in the treatment of acute appendicitis. *World J Gastrointest Surg* 2025; 17(3): 103516 [DOI: 10.4240/wjgs.v17.i3.103516]

Eray IC, Topal U, Gumus S, Isiker K, Yavuz B, Aydin I. Comparative analysis of Ferguson hemorrhoidectomy combined with doppler-guided hemorrhoidal artery ligation and Ferguson hemorrhoidectomy in hemorrhoidal disease treatment. *World J Gastrointest Surg* 2025; 17(3): 103953 [DOI: 10.4240/wjgs.v17.i3.103953]

Clinical Trials Study

Zhu LL, Shen RZ. Follow-up of elderly gastric cancer post-radical surgery: Trauma, complications, and prognosis. *World J Gastrointest Surg* 2025; 17(3): 100143 [DOI: 10.4240/wjgs.v17.i3.100143]

Observational Study

Hu G, Ma J, Qiu WL, Mei SW, Zhuang M, Xue J, Liu JG, Tang JQ. Patient selection and operative strategies for laparoscopic intersphincter resection without diverting stoma. *World J Gastrointest Surg* 2025; 17(3): 95983 [DOI: 10.4240/wjgs.v17.i3.95983]

Randomized Controlled Trial

Tan XQ, Huang XL. Effects of postoperative quantitative assessment strategy-based nursing in patients with colorectal cancer. *World J Gastrointest Surg* 2025; 17(3): 100302 [DOI: 10.4240/wjgs.v17.i3.100302]

SYSTEMATIC REVIEWS

Isah AD, Wang X, Shaibu Z, Yuan X, Dang SC. Systematic review and meta-analysis comparing extraperitoneal and transperitoneal routes of colostomy-related complications. *World J Gastrointest Surg* 2025; 17(3): 98947 [DOI: 10.4240/wjgs.v17.i3.98947]

SCIENTOMETRICS

Wang XY, Chen HY, Sun Q, Li MH, Xu MN, Sun T, Huang ZH, Zhao DL, Li BR, Ning SB, Fan CX. Global trends and research hotspots in esophageal strictures: A bibliometric study. *World J Gastrointest Surg* 2025; 17(3): 100920 [DOI: [10.4240/wjgs.v17.i3.100920](https://doi.org/10.4240/wjgs.v17.i3.100920)]

CASE REPORT

Chen JT, Li YP, Guo SQ, Huang JS, Wang YG. Nonsurgical treatment of postoperative intestinal obstruction caused by heterotopic ossification of the mesentery: A case report. *World J Gastrointest Surg* 2025; 17(3): 99015 [DOI: [10.4240/wjgs.v17.i3.99015](https://doi.org/10.4240/wjgs.v17.i3.99015)]

Wang CD, Liu RD, Liu MJ, Song J. Lung metastasis following temporary discontinuation of lenvatinib and tislelizumab in hepatocellular carcinoma: A case report. *World J Gastrointest Surg* 2025; 17(3): 100951 [DOI: [10.4240/wjgs.v17.i3.100951](https://doi.org/10.4240/wjgs.v17.i3.100951)]

Xu F, Kong J, Dong SY, Xu L, Wang SH, Sun WB, Gao J. Laparoscopic microwave ablation for giant cavernous hemangioma coexistent with diffuse hepatic hemangiomatosis: Two case reports. *World J Gastrointest Surg* 2025; 17(3): 101697 [DOI: [10.4240/wjgs.v17.i3.101697](https://doi.org/10.4240/wjgs.v17.i3.101697)]

Tian ZS, Ma XP, Ruan HX, Yang Y, Zhao YL. Rare large sigmoid hamartomatous polyp in an elderly patient with atypical Peutz-Jeghers syndrome: A case report. *World J Gastrointest Surg* 2025; 17(3): 102174 [DOI: [10.4240/wjgs.v17.i3.102174](https://doi.org/10.4240/wjgs.v17.i3.102174)]

LETTER TO THE EDITOR

Deng HZ, Liu YF, Zhang HW. Role of two-dimensional shear wave elastography in predicting post-hepatectomy liver failure: A step forwards in hepatic surgery. *World J Gastrointest Surg* 2025; 17(3): 98454 [DOI: [10.4240/wjgs.v17.i3.98454](https://doi.org/10.4240/wjgs.v17.i3.98454)]

Rao AG, Nashwan AJ. Enhancing endoscopic retrograde cholangiopancreatography safety: Predictive insights into gastric retention. *World J Gastrointest Surg* 2025; 17(3): 98898 [DOI: [10.4240/wjgs.v17.i3.98898](https://doi.org/10.4240/wjgs.v17.i3.98898)]

Munini M, Fodor M, Corradi A, Frena A. Clinical benefits and controversies of jejunostomy feeding in patients undergoing gastrectomy for gastric cancer. *World J Gastrointest Surg* 2025; 17(3): 100384 [DOI: [10.4240/wjgs.v17.i3.100384](https://doi.org/10.4240/wjgs.v17.i3.100384)]

Pavlidis ET, Galanis IN, Pavlidis TE. Current opinions on the use of prophylactic antibiotics in patients undergoing laparoscopic cholecystectomy. *World J Gastrointest Surg* 2025; 17(3): 101938 [DOI: [10.4240/wjgs.v17.i3.101938](https://doi.org/10.4240/wjgs.v17.i3.101938)]

Xie Y, Xie H, Wang RL. Enhancing palliative care in malignant obstructive jaundice: A critical care perspective on endoscopic biliary stenting. *World J Gastrointest Surg* 2025; 17(3): 103431 [DOI: [10.4240/wjgs.v17.i3.103431](https://doi.org/10.4240/wjgs.v17.i3.103431)]

ABOUT COVER

Peer Reviewer of *World Journal of Gastrointestinal Surgery*, Adrienn Biró, MD, PhD, Assistant Professor, Surgeon, Department of Surgery, Somogy County Kaposi Mor Teaching Hospital, Kaposvár 7400, Hungary.
b.adrienn5@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (*WJGS, World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

INDEXING/ABSTRACTING

The *WJGS* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for *WJGS* as 1.8; JIF without journal self cites: 1.7; 5-year JIF: 1.9; JIF Rank: 126/292 in surgery; JIF Quartile: Q2; and 5-year JIF Quartile: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Eva Lieto

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

March 27, 2025

COPYRIGHT

© 2025 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Efficacy of combined psychological and physical nursing in preventing peripherally inserted central catheter-related thrombosis in gastric cancer patients

Wei-Jing Ni, Yu-Xiu Xi, Yong-Chao Zhou

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade B, Grade C

Novelty: Grade B, Grade C

Creativity or Innovation: Grade B, Grade B

Scientific Significance: Grade C, Grade C

P-Reviewer: Han SU; Jeong O

Received: November 28, 2024

Revised: December 23, 2024

Accepted: January 6, 2025

Published online: March 27, 2025

Processing time: 88 Days and 2.1 Hours



Wei-Jing Ni, Yu-Xiu Xi, Yong-Chao Zhou, Venous Therapy Nursing Clinic, Affiliated Hospital of Jiangnan University, Wuxi 214000, Jiangsu Province, China

Co-corresponding authors: Yu-Xiu Xi and Yong-Chao Zhou.

Corresponding author: Yu-Xiu Xi, Venous Therapy Nursing Clinic, Affiliated Hospital of Jiangnan University, No. 1000 Hefeng Road, Binhu District, Wuxi 214000, Jiangsu Province, China. xiyuxiu@163.com

Abstract

BACKGROUND

Long-term chemotherapy for patients with gastric cancer (GC), facilitated by peripherally inserted central catheter (PICC) catheterization, reduces vascular damage and enhances drug delivery efficiency but carries risks of catheter-related complications. A combination of group psychological nursing and physical movement care significantly mitigates the risk of venous thrombosis and improves psychological well-being, and enhances motor function, underscoring its clinical importance.

AIM

To assess group psychological and physical movement nursing in preventing venous thrombosis in patients with PICC GC.

METHODS

Sixty-five GC patients with PICC, admitted from January 2022 to January 2023, were randomly divided into two groups using the lottery method: A control group ($n = 35$, routine nursing) and an observation group ($n = 30$, routine nursing plus psychological nursing and physical movement nursing). Both groups received continuous care for 2 weeks. Pre-nursing and post-nursing data on psychological state, physical function, chemotherapy-related thrombosis incidence, and cancer-related fatigue were analyzed using SPSS 26.0 and GraphPad Prism 8.0.

RESULTS

After nursing, both groups showed reduced Hamilton Anxiety Scale scores and increased General Perceived Self-Efficacy Scale scores, with the observation group performing better ($P < 0.05$). The Functional Comprehensive Assessment score for

the observation group after nursing was (65.42 ± 2.35) points, lower than the control group's (62.19 ± 4.33) points ($P < 0.05$). Although no significant difference was observed in the incidence of venous thrombosis between the two groups ($\chi^2 = 0.815$, $P = 0.367$), the observation group had lower incidence. Both groups showed decreased Revised Piper Fatigue Scale scores, with the observation group scoring lower ($P < 0.05$).

CONCLUSION

Group psychological and physical movement nursing for patients with PICC reduces venous thrombosis risk, improves psychological well-being, cancer-related fatigue, and physical function, making it highly promotable.

Key Words: Group psychological nursing; Physical movement nursing; Peripherally inserted central catheter; Gastric cancer; Venous thrombosis; Cancer-related fatigue

©The Author(s) 2025. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: In a comprehensive care approach that integrates psychological counseling with physical activity, there is a significant reduction in the risk of thrombosis among gastric cancer patients, particularly those who have undergone percutaneous central venous catheterization. This strategy not only enhances the overall health status of patients but also specifically targets individuals requiring long-term venous therapy due to gastric cancer, offering an effective method for thromboprophylaxis.

Citation: Ni WJ, Xi YX, Zhou YC. Efficacy of combined psychological and physical nursing in preventing peripherally inserted central catheter-related thrombosis in gastric cancer patients. *World J Gastrointest Surg* 2025; 17(3): 100430

URL: <https://www.wjgnet.com/1948-9366/full/v17/i3/100430.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v17.i3.100430>

INTRODUCTION

Gastric cancer (GC), one of the most common malignant tumors, requires prolonged chemotherapy. Repeated vascular punctures increase the risk of vascular damage and complications such as subcutaneous tissue necrosis due to drug extravasation[1]. In recent years, with the continuous promotion of peripherally inserted central catheter (PICC) catheterization, advantages such as protecting blood vessels, rapidly delivering drugs to central veins, and accelerating drug distribution have led to significant therapeutic effects in clinical practice[2]. However, complications such as catheter dislodgement, phlebitis, and venous thrombosis are inevitable during the catheterization process and can affect subsequent treatments. Therefore, exploring measures to reduce venous thrombosis risk in GC patients with PICC is a key focus of clinical research[3,4]. Physical movement can effectively promote local blood circulation in patients with tumors and help prevent thrombosis. However, the psychological state of several patients with GC, affected by fear of cancer and a lack of confidence in treatment, can affect the effectiveness of daily treatment. Compared to routine nursing, group psychological nursing is a novel intervention strategy guided by a psychologist and carried out by a professional team through systematic observation, expression, and communication. This approach aims to transform patients' cognitive structures and effectively alleviate or eliminate negative emotions. It emphasizes teamwork and professional guidance to achieve positive effects on the patients' psychological well-being. Currently, there is limited research on combining group psychological nursing with physical movement nursing for GC patients with PICC. This study observed 65 GC patients with PICC admitted to our hospital and analyzed the application value of this combined nursing approach.

MATERIALS AND METHODS

General data

Sixty-five patients with PICC, admitted between January 2022 and January 2023, were selected. Diagnostic criteria were in accordance with "The Chinese Society of Clinical Oncology: Clinical guidelines for the diagnosis and treatment of GC, 2021"[3]. The inclusion criteria were as follows: (1) Meeting the aforementioned diagnostic criteria and undergoing PICC treatment; (2) Concurrent vitamin K deficiency; (3) No mental disorder; (4) Suitable vessels for puncture; and (5) Informed consent obtained from patients and family members. The exclusion criteria included: (1) Poor compliance; (2) Cellulitis; (3) Active infection at the catheter site; (4) Communication or hearing impairment; and (5) Coagulation mechanism disorders. Patients were randomly divided into two groups using the ball-drawing method: A control group ($n = 35$) and an observation group ($n = 30$), with no significant difference in general data between the two groups ($P > 0.05$) (Table 1).

Table 1 Comparison of general information of two groups of patients

General information	Observation (n = 30)	Control (n = 35)	F/t/ χ^2	P value
Average age (years), mean \pm SD	52.33 \pm 5.84	53.18 \pm 6.42	0.559	0.578
Gender (n)				
Male	19	20	0.258	0.612
Female	11	15	-	-
Catheter vein (n)				
Median cubital vein	11	13	0.511	0.775
Subclavian vein	10	14	-	-
Venae magnalis	9	8	-	-
Average BMI (kg/m ²), mean \pm SD	21.11 \pm 3.11	22.02 \pm 3.12	1.174	0.245
TNM staging				
T1 stage	15	20	1.389	0.499
T2 stage	6	5		
T3 stage	9	6		

BMI: Body mass index; TNM: Tumor-node-metastasis.

Methods

Control group (standard care): During the treatment period, nursing staff were required to assess various vital signs of the patients, including their heart rate and blood pressure. They provided guidance on a scientific diet, avoided cold and greasy foods, and emphasized the safety of catheter placement.

Observation group: In addition to the standard care provided to the control group, the observation group received an integrated program of psychological and physical movement nursing: (1) Formation of the integrated nursing team: The team consisted of one psychological physician, one attending oncology physician, one head nurse, and several nursing staff members. Prior to the study, team members underwent training and assessment on the content of group psychological and physical movement nursing; (2) Psychological intervention: The psychological physician and nursing staff developed intervention content tailored to the actual situation of patients with GC with PICC. Preparation activity: Before the start of group psychological nursing, the physician conducted one-on-one conversations with patients to build rapport and foster confidence in treatment. The WeChat group was established by the nursing staff prior to the first group's psychological activity. Getting to know each other activity: This activity aimed to clarify intervention goals and facilitate familiarity among team members through self-introductions. The physician initiated the process, helping each member define their expected goals and outcomes, encouraging patients to share their situations and concerns, and providing a summary. Emotional expression activity: Starting with the physician, each participant described themselves with an adjective, and subsequent participants built on the previous introduction until all members had introduced themselves. The physician explained the rules, played relevant songs to enliven the atmosphere, and the members expressed emotions such as happiness or sadness until everyone had finished, marking the end of one round. After the two rounds, the members reflected on their experiences. Self-understanding activity: Patients were guided to create self-portraits and share the meanings of different elements within them. The nursing staff prepared cards for patients to record their self-awareness and hopes for their future progress. All cards were collected and placed in a box from which the patients drew cards and read the content aloud, offering support and encouragement to the card owner. Facing life activity: The physician instructed patients to draw a coordinate axis on A4 paper, with the horizontal axis representing age from birth to the present and the vertical axis representing life satisfaction. The patients were guided to recall and mark significant turning points and trends in satisfaction changes by sharing these points with the group. The physician provided targeted advice based on shared experiences, highlighting potential future challenges. Messages for the future activity: Group members wrote down reflections on "progress made", "happiest moments", "greatest help received", and "future expectations", and composed messages for peers, summarizing their gains and feedback on goal achievement. Implementation frequency: Each session lasts for 1.5 hours and was conducted once every 2 weeks; and (3) Physical movement nursing: This included finger flexion and extension, first clenching, and wrist rotation. Patients were instructed to flex and extend their fingers 25 times per set, rest for 10-15 seconds, and then grip a stress ball forcefully for 15 seconds, with a 5-second rest per set. The training session lasted for 15 minutes, 2-3 times a day. Nursing staff also instructed the patients to clench their fists with palms facing outward, extend laterally, and then rotate internally for 1 minute each. These exercises were done for 10-15 minutes per session, 2-3 times daily.

Observational indicators

Psychological state: The psychological state was assessed before and after nursing using the Hamilton Anxiety Scale (HAMA) and General Perceived Self-Efficacy Scale (GSES). The HAMA uses a 0-4 scoring system, with higher scores indicating more severe anxiety symptoms. The GSES consists of 10 items measuring self-confidence in overcoming difficulties or setbacks, scored from 1 to 4. Higher total scores indicate stronger self-efficacy.

Physical movement function: Functional comprehensive assessment (FCA) was used to assess physical movement function before and after nursing. The FCA includes two dimensions, physical and cognitive function, with a total score of 108. Higher scores indicate stronger overall physical function.

Thrombosis incidence during chemotherapy: The incidence of venous thrombosis during chemotherapy was recorded for both groups of patients.

Cancer-related fatigue: The Revised Piper Fatigue Scale (PFE-R) was used to evaluate cancer-related fatigue, with 22 items and five open-ended questions scored 0-10, with higher scores indicating more severe fatigue.

Statistical analysis

All data collected in this study were statistically analyzed using SPSS 26.0. Quantitative data are described as mean \pm SD, with inter-group comparisons made using independent samples *t*-tests and intra-group comparisons made using paired samples *t*-tests. Count data are presented as percentages and compared using χ^2 tests, with $P < 0.05$ indicating significant differences.

RESULTS

Psychological state

Following the nursing intervention, there was a decrease in HAMA scores and an increase in GSES scores in the patients, with the intervention group showing superior psychological state scores compared to the control group ($P < 0.05$), as shown in [Table 2](#).

Motor function

Post-nursing care, both groups exhibited an increase in FCA scores, with the intervention group demonstrating higher scores ($P < 0.05$), as detailed in [Table 3](#).

Thrombosis incidence during chemotherapy

The incidence of thrombosis was 6.67% (2/30) and 17.14% (6/35) in the intervention and control groups, respectively. There was no significant difference in thrombosis incidence between the two groups ($\chi^2 = 0.815$, $P = 0.367$), although the intervention group had a lower incidence.

Cancer-related fatigue

After the nursing intervention, there was a decrease in all PFE-R scores in both groups, with the intervention group showing significantly lower PFE-R scores ($P < 0.05$), as presented in [Table 4](#).

DISCUSSION

The PICC is a relatively safe vascular access device typically used for patients requiring prolonged intravenous infusions or frequent venipuncture for therapy[5,6]. Patients with GC who undergo long-term chemotherapy often present with poor vascular conditions, where repeated punctures can lead to drug extravasation. This not only compromises treatment efficacy but also exacerbates vascular strain[7]. Consequently, effective catheter care interventions are essential to prevent PICC-related venous thrombosis[8]. Studies have suggested that patients with cancer, driven by the fear of the disease, weak confidence in treatment, and concerns about treatment costs, often experience heightened levels of anxiety, depression, and other negative emotions[9]. Although psychological interventions in routine nursing can effectively alleviate symptoms, their ability to significantly improve patients' psychological state is limited due to patients' misconceptions about GC and low trust in treatment methods[10]. Group psychological nursing, as a form of collective intervention, helps address individual and shared issues by leveraging team building, enhanced communication, and emotional support for patients[11]. Additionally, physical movement nursing can promote local blood circulation and dilate local venous vessels through physical activity, contributing to improved vascular.

After nursing, both groups showed a decrease in HAMA scores and an increase in GSES scores, with the observation group's psychological state scores being significantly better than those of the control group ($P < 0.05$). The analysis indicated that group nursing interventions, through collective activities, can reduce patients' feelings of loneliness and anxiety by providing a sense of support and belonging. A psychologist's targeted guidance can help patients develop a positive and proactive mindset[12]. During physical movement nursing, patients can promote blood circulation, prevent thrombosis through physical activities, and master the methods of activity training, thereby building confidence in their

Table 2 Pre-nursing and post-nursing psychological state score comparison between two groups

Groups	Cases	HAMA (scores)		<i>t</i>	<i>P</i> value	GSES (scores)		<i>t</i>	<i>P</i> value
		Before	After			Before	After		
Observation, mean ± SD	30	36.44 ± 3.42	20.53 ± 1.84	22.439	< 0.001	28.75 ± 4.12	36.58 ± 6.22	5.748	< 0.001
Control, mean ± SD	35	36.39 ± 3.30	21.61 ± 2.13	22.262	< 0.001	28.75 ± 4.22	31.25 ± 4.78	2.320	< 0.001
<i>t</i>	-	0.060	2.168	-	-	0.000	3.902	-	-
<i>P</i> value	-	0.952	0.034	-	-	1.000	< 0.001	-	-

HAMA: Hamilton Anxiety Scale; GSES: General Perceived Self-Efficacy Scale.

Table 3 Pre-nursing and post-nursing physical movement function score comparison between two groups

Groups	Cases	FCA score		<i>t</i>	<i>P</i> value
		Before	After		
Observation, mean ± SD	30	40.46 ± 6.58	65.42 ± 2.35	19.566	< 0.001
Control, mean ± SD	35	40.21 ± 6.13	62.19 ± 4.33	17.326	< 0.001
<i>t</i>	-	0.158	3.648	-	-
<i>P</i>	-	0.875	0.001	-	-

FCA: Functional comprehensive assessment.

Table 4 Pre-nursing and post-nursing cancer-related fatigue score comparison between two groups

Items	Time	Observation (<i>n</i> = 30)	Control (<i>n</i> = 35)	<i>t</i>	<i>P</i> value
Emotion (scores), mean ± SD	Before	7.33 ± 1.12	7.22 ± 1.18	0.384	0.703
	After	2.41 ± 0.64	4.56 ± 0.77	12.118	< 0.001
<i>t</i>	-	20.891	11.169	-	-
<i>P</i> value	-	< 0.001	< 0.001	-	-
Cognition (points)	Before	6.68 ± 1.44	6.71 ± 1.51	0.082	0.935
	After	1.87 ± 0.44	2.64 ± 0.74	4.990	< 0.001
<i>t</i>	-	17.497	14.319	-	-
<i>P</i> value	-	< 0.001	< 0.001	-	-
Behavior (points)	Before	8.44 ± 1.51	8.51 ± 1.46	0.190	0.850
	After	2.02 ± 0.43	3.02 ± 0.65	7.183	< 0.001
<i>t</i>	-	22.397	20.323	-	-
<i>P</i> value	-	< 0.001	< 0.001	-	-
Body (points)	Before	7.46 ± 1.13	7.33 ± 1.24	0.439	0.662
	After	2.03 ± 0.41	2.45 ± 0.55	3.441	0.001
<i>t</i>	-	24.742	21.283	-	-
<i>P</i> value	-	< 0.001	< 0.001	-	-

treatment. One study reported that after the intervention, the incidence of mechanical phlebitis, infection, catheter obstruction, and venous thrombosis in the observation group was lower than in the control group ($P < 0.05$). This indicates that group psychological nursing not only improves patients' psychological state but also reduces the occurrence of complications[13]. These findings align with our results, demonstrating that physical movement nursing promotes blood circulation, prevents thrombosis, and helps patients build confidence in treatment[14].

Furthermore, post-nursing, the observation group's PFE-R scores were lower than those of the control group ($P < 0.05$). The analysis showed that compared with routine nursing, the combination of group psychological nursing and physical movement nursing can effectively alleviate cancer-related fatigue in patients. This was attributed to the group's psychological nursing ability to enhance patients' self-awareness and self-acceptance, thereby improving their emotional control capabilities. Physical movement nursing helps patients move their limb extremities, promotes peripheral nerves, and improves blood circulation through activities such as fist clenching and finger flexion. The combined nursing approach enabled patients to correctly understand PICC-related venous thrombosis in GC and chemotherapy, thereby alleviating the psychological fatigue caused by cancer[15]. One study investigated the application of psychological nursing combined with exercise therapy in diabetes care and analyzed its impact on the quality of life. The results demonstrated that compared with routine nursing, the integration of psychological nursing and exercise therapy significantly improved patients' glycemic control and quality of life ($P < 0.05$). This finding is congruent with our observation that physical movement nursing can promote blood circulation, prevent thrombosis, and bolster patient confidence in treatment[16].

The comparison of thrombosis incidence between the two groups showed no statistically significant difference ($\chi^2 = 0.815$, $P = 0.367$), but the incidence was lower in the observation group. The analysis revealed that during physical movement nursing, guidance from nursing staff on exercises, such as slow finger flexion and extension, fist clenching, and wrist rotation, aims to improve peripheral blood circulation and promote local blood flow recovery. This enhances vascular strength and reduces the incidence of local venous thrombosis. Concurrently, the combination of group psychological nursing allowed patients to engage in local physical activities during the group activities, further promoting blood circulation improvement[17]. A study found that combining ankle pump exercises with intermittent pneumatic compression therapy significantly reduced the incidence of lower extremity deep vein thrombosis, with a statistically significant difference ($P < 0.05$)[18]. This finding suggests that limb movement nursing plays a beneficial role in decreasing the occurrence of venous thrombosis, consistent with our findings that physical movement nursing can promote blood circulation, prevent thrombosis, and foster patients' confidence in their treatment. However, this study was hospital-based, potentially introducing selection bias and limiting the representativeness of the sample for the general population. Future validation will involve a larger and more diverse sample to address these limitations and expand the generalizability of the findings.

CONCLUSION

In conclusion, this pioneering study integrated group psychological nursing with physical movement care for patients with GC undergoing PICC therapy. Our findings revealed that this novel, comprehensive nursing approach significantly reduced the risk of venous thrombosis, markedly ameliorated cancer-related fatigue, and enhanced the patients' motor function. These results introduce an innovative strategy for the clinical care of patients with GC and hold significant implications for guiding clinical practice, advocating for the broader adoption of this nursing model in clinical settings.

FOOTNOTES

Author contributions: Ni WJ designed the research study; Ni WJ and Xi YX contributed new reagents and analytical tools; Xi YX and Zhou YC jointly designed the research, performed the research, wrote the manuscript, visited the entire process, interpreted the results, and were responsible for manuscript revision and communication with the journal, ensuring the integrity and quality of the study, they contributed equally to this article, they are the co-corresponding authors of this manuscript; and all authors have read and approved the final manuscript.

Institutional review board statement: This study was approved by the Medical Ethics Committee of Affiliated Hospital of Jiangnan University, approval No. LS2024073.

Informed consent statement: All study participants and their legal guardians provided written informed consent before recruitment.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: No additional data are available.

Open Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: China

ORCID number: Wei-Jing Ni 0009-0008-2116-9346; Yu-Xiu Xi 0009-0006-0510-9031.

S-Editor: Bai Y

L-Editor: A

P-Editor: Xu ZH

REFERENCES

- 1 **Xiang L**, Jin S, Yu Y, Wang D, Chen H. Risk of venous thromboembolism in patients undergoing gastric cancer surgery: a systematic review and meta-analysis. *BMC Cancer* 2023; **23**: 933 [PMID: 37789268 DOI: 10.1186/s12885-023-11424-x]
- 2 **Mielke D**, Wittig A, Teichgräber U. Peripherally inserted central venous catheter (PICC) in outpatient and inpatient oncological treatment. *Support Care Cancer* 2020; **28**: 4753-4760 [PMID: 31970514 DOI: 10.1007/s00520-019-05276-0]
- 3 **Song X**, Lu H, Chen F, Bao Z, Li S, Li S, Peng Y, Liu Q, Chen X, Li J, Zhang W. A longitudinal observational retrospective study on risk factors and predictive model of PICC associated thrombosis in cancer patients. *Sci Rep* 2020; **10**: 10090 [PMID: 32572092 DOI: 10.1038/s41598-020-67038-x]
- 4 **Wan L**, Yin L, Zhang W, Yang J, Yi X, Guo L. Application value of modified skin expansion in PICC catheterization under the guidance of B-ultrasound in gastrointestinal cancer patients with chemotherapy. *Am J Transl Res* 2022; **14**: 7932-7941 [PMID: 36505288]
- 5 **Wang FH**, Zhang XT, Li YF, Tang L, Qu XJ, Ying JE, Zhang J, Sun LY, Lin RB, Qiu H, Wang C, Qiu MZ, Cai MY, Wu Q, Liu H, Guan WL, Zhou AP, Zhang YJ, Liu TS, Bi F, Yuan XL, Rao SX, Xin Y, Sheng WQ, Xu HM, Li GX, Ji JF, Zhou ZW, Liang H, Zhang YQ, Jin J, Shen L, Li J, Xu RH. The Chinese Society of Clinical Oncology (CSCO): Clinical guidelines for the diagnosis and treatment of gastric cancer, 2021. *Cancer Commun (Lond)* 2021; **41**: 747-795 [PMID: 34197702 DOI: 10.1002/cac2.12193]
- 6 **Bahoush G**, Salajegheh P, Anari AM, Eshghi A, Aski BH. A review of peripherally inserted central catheters and various types of vascular access in very small children and pediatric patients and their potential complications. *J Med Life* 2021; **14**: 298-309 [PMID: 34377194 DOI: 10.25122/jml-2020-0011]
- 7 **Chen P**, Teng S, Chen MG, Yao TT. Comparison of the effect of two-way evaluation model on the whole process management of PICC catheterization in patients with lung cancer. *Asian J Surg* 2023; **46**: 2592-2594 [PMID: 36641275 DOI: 10.1016/j.asjsur.2022.12.125]
- 8 **Xu YP**, Shang ZR, Dorazio RM, Shi LP. Risk factors for peripherally inserted central catheterization-associated bloodstream infection in neonates. *Zhongguo Dang Dai Er Ke Za Zhi* 2022; **24**: 141-146 [PMID: 35209978 DOI: 10.7499/j.issn.1008-8830.2109147]
- 9 **Johansson E**, Hammar skjöld F, Lundberg D, Arnlind MH. Advantages and disadvantages of peripherally inserted central venous catheters (PICC) compared to other central venous lines: a systematic review of the literature. *Acta Oncol* 2013; **52**: 886-892 [PMID: 23472835 DOI: 10.3109/0284186X.2013.773072]
- 10 **He F**, He RX. Systematic nursing interventions in gastric cancer: A randomized controlled study. *World J Clin Cases* 2022; **10**: 1843-1851 [PMID: 35317163 DOI: 10.12998/wjcc.v10.i6.1843]
- 11 **Liu F**, Yao K, Liu X. Analysis on effect of psychological nursing combined with extended care for improving negative emotions and self-care ability in patients with colorectal cancer and enterostomy: A retrospective study. *Medicine (Baltimore)* 2024; **103**: e38165 [PMID: 38788010 DOI: 10.1097/MD.00000000000038165]
- 12 **Finsrud I**, Nissen-Lie HA, Ulvenes P, Melsom L, Vrabel K, Wampold B. Confidence in the therapist and confidence in the treatment predict symptomatic improvement week by week in therapy: A latent curve modeling approach. *J Couns Psychol* 2022; **69**: 823-834 [PMID: 36136794 DOI: 10.1037/cou0000640]
- 13 **Yin L**, Zhang W, Liu L, Guo L, Guo M, He X, Zhu L. Application of nursing intervention based on the IKAP model in self-management of patients with gastric cancer. *Am J Transl Res* 2022; **14**: 6389-6398 [PMID: 36247277]
- 14 **Engineering JOH**. Retracted: Prevention and Nursing Research of PICC Catheter-Related Complications in Patients with Digestive System Malignant Tumor Based on Smart Medical Block Chain. *J Healthc Eng* 2022; **2022**: 9864161 [PMID: 36465252 DOI: 10.1155/2022/9864161]
- 15 **Hou A**, Zhao C. Structured Psychological Nursing Combined with Group Health Education for Patients Receiving Blood Purification. *Altern Ther Health Med* 2024 [PMID: 38870493]
- 16 **Zhang X**, Bai X, Yan Z. Application of psychological nursing combined with exercise therapy in nursing of gestational diabetes mellitus. *Minerva Med* 2023; **114**: 117-118 [PMID: 33242942 DOI: 10.23736/S0026-4806.20.07104-9]
- 17 **Tian X**, Zhou X, Sun M, Yu NX, Peng Y, Zheng X, Xiao W. The effectiveness of positive psychological interventions for patients with cancer: A systematic review and meta-analysis. *J Clin Nurs* 2024; **33**: 3752-3774 [PMID: 38979929 DOI: 10.1111/jocn.17358]
- 18 **Sun Y**, Ren T, Ji X. Influence of hyperbaric oxygen therapy-specialized care on limb motor function and mental state of cerebral infarction patients with hemiplegia. *Technol Health Care* 2024; **32**: 1967-1976 [PMID: 38393863 DOI: 10.3233/THC-231643]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

