

Name of Journal: *World Journal of Hepatology*

ESPS Manuscript NO: 29805

Manuscript Type: Original Article

Case Control Study

Egg consumption and risk of Non-alcoholic fatty liver disease

Zeinab Mokhtari, Hossein Poustchi, Tannaz Eslamparast, Azita Hekmatdoost

Abstract

AIM

To evaluate the association between egg consumption and risk of non-alcoholic fatty liver disease (NAFLD) development.

Match Overview

1	Crossref 127 words Hekmatdoost, Azita, Ali Shamsipour, Mohammad Meibodi, Newsha Gheibizadeh, Tannaz Eslamparast, and Hos	5%
2	Crossref 23 words He, J., J. H. Lee, M. Febbraio, and W. Xie. "The emerging roles of fatty acid translocase/CD36 and the aryl hydro...	1%
3	Crossref 22 words Mokhtari, Z., S. Hosseini, R. Miri, A. R. Baghestani, M. Zahedirad, M. Rismanchi, and J. Nasrollahzadeh. "Relation	1%
4	Internet 15 words crawled on 22-Aug-2012 www.wjgnet.com	1%
5	Internet 15 words crawled on 27-Feb-2014 www.ncbi.nlm.nih.gov	1%
6	Internet 12 words crawled on 18-Mar-2016 www.mdpi.com	<1%
7	Publications 12 words	<1%

[All](#)[News](#)[Images](#)[Shopping](#)[Videos](#)[More ▾](#)[Search tools](#)

About 123,000 results (1.04 seconds)

Scholarly articles for **Egg consumption and risk of Non-alcoholic fatty liver disease (NAFLD)**

... content: prevalence of hepatic **steatosis** in the general ... - [Szczepaniak](#) - Cited by 1021

... postprandial lipemia in nonalcoholic **steatohepatitis** - [Musso](#) - Cited by 558

Pathogenesis of **non-alcoholic fatty liver disease** - [Dowman](#) - Cited by 324

Dietary habits and behaviors associated with nonalcoholic fatty liver ...

www.ncbi.nlm.nih.gov/PMC39309 ... ▾ National Center for Biotechnology Information ▾
by K Yasutake - 2014 - Cited by 33 - Related articles

Feb 21, 2014 - **Nonalcoholic fatty liver disease (NAFLD)** is one of the most frequent causes of ... **risks** were higher for patients with **NAFLD** (OR = 1.69) and **NASH** (OR **intake** of food high in cholesterol, such as **eggs**, fish **eggs**, liver, and ...

Dietary recommendations for patients with nonalcoholic fatty liver ...

www.ncbi.nlm.nih.gov/PMC40278 ... ▾ National Center for Biotechnology Information ▾
by A Kargulewicz - 2014 - Cited by 11 - Related articles

Mar 1, 2014 - Moderate **intake** of protein is optimum for patients with **NAFLD**, due to the alleviation of insulin resistance. ... Ingestion of monounsaturated fatty acids (MUFAs) lowers the **risk of** ... The source of omega-6 fatty acids are: cereals, **eggs** and animal fat [2] ... Protein content – diet in **nonalcoholic fatty liver disease**.

Dietary Patterns Modulate the Risk of Non-Alcoholic Fatty Liver ... - NCBI

www.ncbi.nlm.nih.gov/PMC44888 ... ▾ National Center for Biotechnology Information ▾
by CQ Yang - 2015 - Cited by 3 - Related articles

Jun 15, 2015 - **Non-alcoholic fatty liver disease (NAFLD)**, including simple by a high **consumption** of staple food, coarse grains, fruits, **eggs**, fish and shrimp, ...

Comparative review of diets for the metabolic syndrome: implications ...

[全部](#) [新闻](#) [图片](#) [视频](#) [购物](#) [更多](#)

[设置](#) [工具](#)

找到约 6,650,000 条结果 (用时 0.80 秒)

Google 学术: Egg consumption and risk of non-alcoholic fatty liver disease

... novel insights into non-alcoholic fatty liver disease and ... - Corbin - 被引用次数: 90

Hepatocellular carcinoma: epidemiology, risk factors ... - Gomaa - 被引用次数: 514

Alcoholic fatty liver in man on a high protein and low fat ... - Lieber - 被引用次数: 215

Dietary habits and behaviors associated with nonalcoholic fatty liver ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: K Yasutake - 2014 - 被引用次数: 37 - [相关文章](#)

2014年2月21日 - Nonalcoholic fatty liver disease (NAFLD) is one of the most frequent causes of intake of food high in cholesterol, such as eggs, fish eggs, liver, and cakes, ... Although saturated fatty acids increase the likelihood of NAFLD ...

Implications of Diet on Nonalcoholic Fatty Liver Disease - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: S Sullivan - 2010 - 被引用次数: 39 - [相关文章](#)

It is reasonable to advise patients with NAFLD to reduce calorie intake with either low fat or ... Nonalcoholic Fatty Liver Disease (NAFLD) is a spectrum of liver diseases ... with an increased risk of type 2 diabetes mellitus, coronary heart disease, ... in IHTG associated with high calorie/high fat feeding; however meat, but not ...

Soft drinks consumption and nonalcoholic fatty liver disease - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: W Nseir - 2010 - 被引用次数: 136 - [相关文章](#)

2010年6月7日 - Nonalcoholic fatty liver disease (NAFLD) is a common clinical ... This form of liver injury carries a 20%-50% risk for progressive intake of SD and meat and a tendency towards a lower intake of fish rich in omega-3[36]. Go to: ...

[全部](#) [新闻](#) [图片](#) [视频](#) [购物](#) [更多](#)

[设置](#) [工具](#)

找到约 1,790,000 条结果 (用时 0.74 秒)

Google 学术: Egg consumption and risk of non-alcoholic fatty liver disease

... novel insights into non-alcoholic fatty liver disease and ... - Corbin - 被引用次数: 92

Hepatocellular carcinoma: epidemiology, risk factors ... - Gomaa - 被引用次数: 523

Alcoholic fatty liver in man on a high protein and low fat ... - Lieber - 被引用次数: 215

Dietary Patterns Modulate the Risk of Non-Alcoholic Fatty Liver ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: CQ Yang - 2015 - 被引用次数: 5 - 相关文章

2015年6月15日 - Non-alcoholic fatty liver disease (NAFLD), including simple cooked meat, and beans), and the "high-salt" pattern (high intake of rice, pickled ...

Dietary habits and behaviors associated with nonalcoholic fatty liver ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: K Yasutake - 2014 - 被引用次数: 39 - 相关文章

2014年2月21日 - Nonalcoholic fatty liver disease (NAFLD) is one of the most frequent causes of intake of food high in cholesterol, such as eggs, fish eggs, liver, and cakes, ... Although saturated fatty acids increase the likelihood of NAFLD ...

Nonalcoholic Fatty Liver Disease: A Nutritional Approach

online.liebertpub.com/doi/full/10.1089/met.2011.0145 ▾ 翻译此页

作者: Y Colak - 2012 - 被引用次数: 15 - 相关文章

2012年3月6日 - Nonalcoholic fatty liver disease (NAFLD) is among the most common the consumption of high amounts of red meat causes insulin resistance, increased Long term nutritional intake and the risk for non-alcoholic fatty liver