

World Journal of *Hepatology*

World J Hepatol 2017 April 8; 9(10): 491-532



REVIEW

- 491 Drug-induced liver injury: Do we know everything?

Alempijevic T, Zec S, Milosavljevic T

ORIGINAL ARTICLE**Case Control Study**

- 503 Egg consumption and risk of non-alcoholic fatty liver disease

Mokhtari Z, Poustchi H, Eslamparast T, Hekmatdoost A

Retrospective Study

- 510 Serum 25-hydroxyvitamin D deficiency and hepatic encephalopathy in chronic liver disease

Vidot H, Potter A, Cheng R, Allman-Farinelli M, Shackel N

Prospective Study

- 519 Hepatic encephalopathy before and neurological complications after liver transplantation have no impact on the employment status 1 year after transplantation

Pflugrad H, Tryc AB, Goldbecker A, Strassburg CP, Barg-Hock H, Klempnauer J, Weissenborn K

ABOUT COVER

Editorial Board Member of *World Journal of Hepatology*, Dr. Henning Pflugrad, Department of Neurology, Hannover Medical School, 30625 Hannover, Germany

AIM AND SCOPE

World Journal of Hepatology (*World J Hepatol*, *WJH*, online ISSN 1948-5182, DOI: 10.4254), is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJH covers topics concerning liver biology/pathology, cirrhosis and its complications, liver fibrosis, liver failure, portal hypertension, hepatitis B and C and inflammatory disorders, steatohepatitis and metabolic liver disease, hepatocellular carcinoma, biliary tract disease, autoimmune disease, cholestatic and biliary disease, transplantation, genetics, epidemiology, microbiology, molecular and cell biology, nutrition, geriatric and pediatric hepatology, diagnosis and screening, endoscopy, imaging, and advanced technology. Priority publication will be given to articles concerning diagnosis and treatment of hepatology diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, functional diagnostics, and physical diagnosis; and comprehensive therapy, drug therapy, surgical therapy, interventional treatment, minimally invasive therapy, and robot-assisted therapy.

We encourage authors to submit their manuscripts to *WJH*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

World Journal of Hepatology is now indexed in PubMed, PubMed Central, and Scopus.

FLYLEAF

I-IV Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Dan Li*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Fang-Fang Ji*
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL
World Journal of Hepatology

ISSN
 ISSN 1948-5182 (online)

LAUNCH DATE
 October 31, 2009

FREQUENCY
 36 Issues/Year (8th, 18th, and 28th of each month)

EDITORS-IN-CHIEF
Clara Balsano, PhD, Professor, Department of Biomedicine, Institute of Molecular Biology and Pathology, Rome 00161, Italy

Wan-Long Chuang, MD, PhD, Doctor, Professor, Hepatobiliary Division, Department of Internal Medicine, Kaohsiung Medical University Hospital, Kaohsiung Medical University, Kaohsiung 807, Taiwan

EDITORIAL BOARD MEMBERS
 All editorial board members resources online at <http://www.wjgnet.com>

www.wjgnet.com/1948-5182/editorialboard.htm

EDITORIAL OFFICE
 Xiu-Xia Song, Director
World Journal of Hepatology
 Baishideng Publishing Group Inc
 8226 Regency Drive, Pleasanton, CA 94588, USA
 Telephone: +1-925-2238242
 Fax: +1-925-2238243
 E-mail: editorialoffice@wjgnet.com
 Help Desk: <http://www.fjpublishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
 Baishideng Publishing Group Inc
 8226 Regency Drive,
 Pleasanton, CA 94588, USA
 Telephone: +1-925-2238242
 Fax: +1-925-2238243
 E-mail: bpgoffice@wjgnet.com
 Help Desk: <http://www.fjpublishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
 April 8, 2017

COPYRIGHT
 © 2017 Baishideng Publishing Group Inc. Articles published by this Open Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
 All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.fjpublishing.com>

Case Control Study

Egg consumption and risk of non-alcoholic fatty liver disease

Zeinab Mokhtari, Hossein Poustchi, Tannaz Eslamparast, Azita Hekmatdoost

Zeinab Mokhtari, Tannaz Eslamparast, Azita Hekmatdoost, Department of Clinical Nutrition and Dietetics, Faculty of Nutrition Sciences and Food Technology, National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, Tehran 1981619573, Iran

Hossein Poustchi, Liver and Pancreatobiliary Diseases Research Group, Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran 14117, Iran

Author contributions: Mokhtari Z, Poustchi H, Eslamparast T and Hekmatdoost A conceptualized and designed the study and wrote the manuscript; Mokhtari Z analyzed data; Eslamparast T and Poustchi H collected data; Hekmatdoost A and Poustchi H interpreted the data, provided professional comments, and critically revised the manuscript for intellectual content and data accuracy.

Institutional review board statement: The study was approved by the ethics committee of National Nutrition and Food Technology Research Institute, Tehran, Iran.

Informed consent statement: All patients signed the informed consent form.

Conflict-of-interest statement: None of the authors had any personal or financial conflicts of interest to report.

Data sharing statement: Technical appendix, statistical code, and dataset available from the corresponding author at a_hekmat2000@yahoo.com.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Azita Hekmatdoost, MD, PhD, RD,

Department of Clinical Nutrition and Dietetics, Faculty of Nutrition Sciences and Food Technology, National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, West Arghavan St., Farahzadi Blvd., P.O. Box: 19395-4741, Tehran 1981619573, Iran. a_hekmat2000@yahoo.com
Telephone: +98-21-22360658
Fax: +98-21-22360657

Received: August 27, 2016

Peer-review started: August 29, 2016

First decision: September 27, 2016

Revised: January 4, 2017

Accepted: March 14, 2017

Article in press: March 17, 2017

Published online: April 8, 2017

Abstract

AIM

To evaluate the association between egg consumption and risk of non-alcoholic fatty liver disease (NAFLD) development.

METHODS

This case-control study was conducted on individuals who were referred to two hepatology clinics in Tehran, Iran in 2015. The study included 169 patients with NAFLD and 782 controls. Egg consumption was estimated using a validated food frequency questionnaire. The participants were categorized according to the frequency of their egg consumption during the previous year: Less than two eggs per week, two to three eggs per week, and four or more eggs per week.

RESULTS

In the crude model, participants who consumed 2 to 3 eggs per week, were 3.56 times more likely to have NAFLD in comparison to those who consumed less than 2 eggs per week (OR: 3.56; 95%CI: 2.35-5.31). Adjustment for known risk factors of NAFLD strengthened

this significant association so that individuals have consumed two to three eggs per week had 3.71 times higher risk of NAFLD than those who have eaten less than two eggs per week (OR: 3.71; 95%CI: 1.91, 7.75).

CONCLUSION

Our data indicate that higher egg consumption in common amount of usage is associated with higher risk of NAFLD.

Key words: Egg; Diet; Non-alcoholic fatty liver disease; Dietary cholesterol

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The data indicate that egg consumption in common amount of usage is associated with risk of non-alcoholic fatty liver disease. According to the case-control design of this study, it can not show the causality effect; thus, these findings should be confirmed in future prospective studies with separate parts of eggs to find the etiological relationships.

Mokhtari Z, Poustchi H, Eslamparast T, Hekmatdoost A. Egg consumption and risk of non-alcoholic fatty liver disease. *World J Hepatol* 2017; 9(10): 503-509 Available from: URL: <http://www.wjgnet.com/1948-5182/full/v9/i10/503.htm> DOI: <http://dx.doi.org/10.4254/wjh.v9.i10.503>

INTRODUCTION

Non-alcoholic fatty liver disease (NAFLD) includes a spectrum of liver disorders from simple steatosis to non-alcoholic steatohepatitis (NASH), fibrosis, cirrhosis, and even hepatocellular carcinoma^[1]. NAFLD is the most common cause of chronic liver diseases around the world^[2] and may be considered as hepatic manifestation of metabolic syndrome^[3]. The increasing prevalence of obesity, together with insulin resistance, hypertension, dyslipidemia, and eventually the metabolic syndrome dispose many people to the risk of NAFLD development in the future years^[4].

Increasing evidence showed that dietary factors contribute to the pathophysiology and treatment of NAFLD^[5-7]. Among the known dietary factors that involved in the development of NAFLD, dietary cholesterol has drawn a great deal of attention. Current studies of animal models propose that excess dietary cholesterol is regarded as the key factor related to the risk of steatohepatitis and hepatic inflammation^[8-10]. Addition of cholesterol to the diet of obese, diabetic mice increased the accumulation of hepatic free cholesterol, hepatocyte apoptosis, and liver fibrosis^[11]. Moreover, an association between raised cholesterol intake and the risk or severity of NAFLD has been addressed by epidemiological studies^[12-14].

Among single foods, eggs are regarded as a main

source of dietary cholesterol, with one large egg containing almost 210 mg of cholesterol; on the other hand, eggs are rich in proteins, and other nutrients^[15], which can improve human health. There is limited evidence on the relationship between egg consumption and NAFLD and its risk factors with controversial results^[16-18]. Therefore, the present study was designed to examine the association between egg consumption and risk of NAFLD development.

MATERIALS AND METHODS

Participants

The present case-control study was conducted on individuals who were undertaken a liver Ultrasound, and were referred to two Hepatology clinics in Tehran, Iran in 2015. The study included 169 patients with NAFLD and 782 controls. The cases were patients with NAFLD, which was diagnosed by a gastroenterologist according to the presence of hepatic steatosis in Ultrasound exam within previous month, and referred to our clinics to be examined by Fibroscan[®], and the Fibroscan result showed a Controlled Attenuation Parameter score of more than 263, and fibrosis score of more than 7. These patients were selected with the convenience-sampling procedure. Controls were randomly selected age- and sex-matched subjects from the same clinic among patients with pancreatobiliary disorders who had been undertaken an Ultrasound showing no hepatic steatosis. The age ranges for matching were 20-40, 40-60 and > 60 years old. Data on each pair of cases and controls were collected at the same time. The participation rate in the study was 94% for cases and 98% for controls. Written informed consent was obtained from all the participants. The study protocol was approved by the local Ethics Review Committee.

Assessment of dietary intake

Dietary intake of patients was assessed using a valid and reliable semi-quantitative food frequency questionnaire (FFQ), which included 168 items of foods with standard servingsizes, as commonly consumed by Iranians^[19]. The consumption frequency of each food item was questioned on a daily, weekly or monthly basis and converted to daily intakes. In the case of egg consumption, the participants were categorized according to the frequency of their egg consumption during the previous year: Less than two eggs per week, two to three eggs per week, and four or more eggs per week. Dietary nutrients intakes were calculated using NUTRITIONIST V (First Databank, Hearst Corp, San Bruno, CA, United States). The patients who had completed less than 90% of dietary questionnaires and subjects who reported extremely low or high energy intakes (< 500 or > 5000 kcal/d) were excluded from the study^[20].

Assessment of other variables

Physical activity was evaluated using the metabolic

Table 1 Baseline characteristics, biochemical parameters and dietary intakes of study participants based on the patients with non-alcoholic fatty liver disease and control group

	Cases (<i>n</i> = 169)	Controls (<i>n</i> = 782)	<i>P</i> value ^a
Age (yr), mean ± SD	42.65 ± 12.21	43.71 ± 14.52	0.373
Male <i>n</i> (%)	81 (47.9)	314 (40.2)	0.063
BMI (kg/m ²), mean ± SD	33.19 ± 8.71	27.74 ± 4.495	< 0.001
Physical activity (MET), mean ± SD	31.89 ± 3.15	34.33 ± 2.85	< 0.001
Current smokers, <i>n</i> (%)	151 (89.9)	145 (18.5)	< 0.001
Drank alcohol in past year, <i>n</i> (%)	22 (13.1)	68 (8.7)	0.077
Diabetes type 2, <i>n</i> (%)	26 (15.6)	53 (6.8)	< 0.001
FBS (mg/dL), mean ± SD	109.29 ± 39.39	90.09 ± 29.24	< 0.001
Total cholesterol (mg/dL), mean ± SD	184.79 ± 54.94	177.72 ± 38.74	0.221
LDL (mg/dL), mean ± SD	121.17 ± 43.04	104.26 ± 31.65	< 0.001
HDL (mg/dL), mean ± SD	41.26 ± 16.72	47.72 ± 10.51	0.001
Triglycerides (mg/dL), mean ± SD	180.40 ± 123.81	131.97 ± 81.59	< 0.001
Total energy (kcal), mean ± SEM	2627.67 ± 61.39	2746.69 ± 27.23	0.068
Carbohydrate (% of total energy), mean ± SEM	58.12 ± 0.95	59.82 ± 0.44	0.001
Protein (% of total energy), mean ± SEM	15.84 ± 0.18	14.07 ± 0.08	< 0.001
Fat (% of total energy), mean ± SEM	29.23 ± 0.30	33.78 ± 0.20	< 0.001
Dietary cholesterol (mg/d), mean ± SEM	315.31 ± 11.50	263.41 ± 5.35	< 0.001
Saturated fat (g/d), mean ± SEM	30.62 ± 5.72	62.67 ± 2.67	< 0.001
Monounsaturated fat (g/d) (mg/d), mean ± SEM	29.85 ± 0.48	32.00 ± 0.23	< 0.001
Polyunsaturated fat (g/d) (mg/d), mean ± SEM	18.51 ± 5.74	59.58 ± 2.67	< 0.001
Dietary fiber (g/d), mean ± SEM	19.21 ± 0.50	14.68 ± 0.23	< 0.001
Red/processed meats (g/d), mean ± SEM	70.95 ± 2.66	36.00 ± 1.24	< 0.001

^aIndependent *t*-test for quantitative variables and χ^2 test for qualitative variables. Dietary intakes (except total energy) were adjusted for age and total energy intake. BMI: Body mass index; MET: Metabolic equivalent task; FBS: Fasting blood sugar; LDL: Low-density lipoprotein cholesterol; HDL: High density lipoprotein cholesterol.

equivalent task (MET) questionnaire^[21,22]. Other covariate information including age, gender, smoking habits, alcohol consumption, medical history, and current use of medications were assessed using questionnaires. Weight and height of all participants were measured.

Statistical analysis

Baseline characteristics and dietary intakes were compared between cases and controls using *t*-test for continuous variables and χ^2 for categorical variables. Egg consumption was divided into three ascending categories on an ordinal scale. Mean or prevalence of baseline characteristics was computed for each category. Baseline characteristics were also compared using ANOVA for continuous variables and χ^2 for categorical variables. The relationship between NAFLD and egg consumption was assessed using multiple regression analysis. Estimates were presented in three models; the first model was adjusted for age (continuous), and total energy intake (kcal/d). In the second model, we further controlled for body mass index (BMI), history of diabetes and smoking (non-smoker, current smoker). Finally, we further adjusted for physical activity (MET) and gender. All models were conducted by treating the first category of egg consumption (< 2/wk) as a reference. All the statistical analyses were done using SPSS for Windows (version 19; SPSS Inc., Chicago, IL).

RESULTS

Baseline characteristics, biochemical parameters and

dietary intakes of the cases and controls are shown in Table 1. Mean age of the total study population was 43.54 ± 14.13 years and 41.5% (395) of participants were male. By design, cases and controls had the similar age and sex distribution. Patients with NAFLD had significantly more BMI, lower physical activity, lower consumption of alcohol, and were more likely to be smoker, and have diabetes in comparison to controls. Furthermore, the cases had elevated fasting blood glucose (FBS), low-density lipoprotein cholesterol (LDL), Triglycerides, and reduced high density lipoprotein cholesterol (HDL) levels and increased intake of protein, cholesterol, fiber and red/processed meats compared with the controls (Table 1).

Basic characteristics and dietary intakes of the studied participants by categories of egg consumption are presented in Table 2. Compared to egg consumption of lower than two per week, higher egg consumption was associated with a lower average age, male sex, current smoking, higher energy intake, lower percent of total energy from carbohydrate and fat. Additionally, the subjects with higher egg consumption tended to consume more protein, cholesterol, monounsaturated fat and red/processed meats, but less saturated and polyunsaturated fatty acids (Table 2).

In secondary analysis, there was a similar egg-NAFLD association in women (*P*-trend 0.001) and men (*P*-trend 0.048) (Table 3).

Multivariate adjusted odds ratios for NAFLD based on egg consumption categories are indicated in Figure 1. In the crude model, participants that consumed 2 to 3 eggs per week, were 3.56 times more likely to have NAFLD in

Table 2 Basic characteristics and dietary intakes of study participants by frequency of egg consumption *n* (%)

	Egg consumption categories			<i>P</i> value ^a
	< 2/wk (<i>n</i> = 589)	2-3/wk (<i>n</i> = 142)	≥ 4/wk (<i>n</i> = 220)	
Age (yr)	45.65 ± 12.26	39.73 ± 13.18	40.35 ± 13.30	< 0.001
Male gender	218 (37.0)	56 (39.4)	121 (55)	< 0.001
BMI (kg/m ²), mean ± SD	28.58 ± 5.44	29.60 ± 7.34	28.51 ± 5.87	0.150
Physical activity (MET), mean ± SD	33.99 ± 3.05	33.42 ± 3.21	33.94 ± 2.95	0.136
Current smokers	155 (26.3)	59 (41.8)	82 (37.3)	< 0.001
Total energy (kcal), mean ± SEM	2580.59 ± 30.68	2744.94 ± 57.45	3101.07 ± 51.20	< 0.001
Carbohydrate (% of total energy), mean ± SEM	60.44 ± 0.67	59.48 ± 0.63	58.14 ± 0.85	0.001
Protein (% of total energy), mean ± SEM	14.09 ± 0.10	14.71 ± 0.20	14.95 ± 0.17	0.001
Fat (% of total energy), mean ± SEM	33.06 ± 0.24	32.56 ± 0.49	32.97 ± 0.40	< 0.001
Dietary cholesterol (mg/d)	226.40 ± 5.75	291.95 ± 11.60	383.90 ± 9.53	< 0.001
Saturated fat (g/d)	56.70 ± 3.16	64.70 ± 6.38	52.57 ± 5.24	< 0.001
Monounsaturated fat (g/d) (mg/d), mean ± SEM	31.20 ± 0.26	31.32 ± 0.53	32.91 ± 0.44	< 0.001
Polyunsaturated fat (g/d) (mg/d), mean ± SEM	53.10 ± 3.20	57.26 ± 6.45	46.71 ± 5.30	< 0.001
Dietary fiber (g/d)	15.65 ± 0.28	16.25 ± 0.57	14.60 ± 0.47	< 0.001
Red/processed meats (g/d)	37.76 ± 1.53	47.79 ± 3.10	50.51 ± 2.54	< 0.001

^aDietary intakes (except total energy) were adjusted for age and total energy intake. BMI: Body mass index; MET: Metabolic equivalent task.

Table 3 Odds ratio for non-alcoholic fatty liver disease according to egg consumption stratified by gender

Egg consumption	Multivariate adjusted model ^a	
	Female	Male
< 2/wk	1.00	1.00
2-3/wk	5.55 (2.30-13.37)	1.90 (0.50-7.16)
≥ 4/wk	1.67 (0.68-4.10)	0.25 (0.06-1.01)
<i>P</i> for trend	0.001	0.048

^aAdjusted for age, energy intake, body mass index, history of diabetes, smoking, and physical activity.

comparison to those who consumed less than 2 eggs per week (OR: 3.56; 95%CI: 2.35-5.31). After controlling for age and total energy intake, consuming 2 to 3 eggs per week was positively associated with the risk of NAFLD (OR: 3.83; 95%CI: 2.49-5.89). These associations remained significant even after additionally controlling for BMI, history of diabetes and smoking (OR: 3.57; 95%CI: 1.89-6.75). Further adjustment for physical activity, and gender strengthened this significant association so that individuals who have consumed two to three eggs per week had 3.71 times higher risk of NAFLD than those who have eaten less than two eggs per week (OR: 3.71; 95%CI: 1.91-7.75). Egg consumption more than four per week was not significantly associated with the NAFLD risk.

DISCUSSION

The results of the present study showed that the egg consumption increases the risk of NAFLD in common range of its consumption (two to three eggs per week). This relationship was also significant after adjustment for age, gender, BMI, history of diabetes, smoking, and physical activity.

The role of diet and dietary supplements on the pathogenesis of NAFLD have been shown previously^[23-36]; however, to our knowledge, no study has yet evaluated the

association of egg consumption and NAFLD risk. It is well established that eggs contain a wide variety of essential nutrients and bioactive compounds that can affect human health. Their high quality protein, fats and micronutrients and low price make them an important part of many people's diet^[37]; despite the nutritional benefits of egg consumption, there are concerns about their high content of cholesterol and saturated fat and their influences on metabolic disorders^[38]. Thus, one possible explanation for the inverse association between egg consumption and risk of NAFLD development may be due to the high cholesterol content of egg. Previous studies have shown that a higher consumption of cholesterol is associated with NAFLD and its exacerbation^[12,13,39,40]. In addition, the presence of high amount of cholesterol in diet is necessary for development of NAFLD^[41]. Baumgartner *et al.*^[39] have shown that daily egg consumption increases serum cholesterol and LDL-C concentrations in women; however, there was no effects on markers for inflammation, endothelial activity, and liver function. Interestingly, the consumption of egg white hydrolyzed with pepsin considerably improved hepatic steatosis^[42]. Thus, it seems that the association between egg consumption and NAFLD is mainly due to high cholesterol content of it, and might not be seen when people consume only the white part of it. Therefore, more studies are recommended to evaluate the effects of consumption of different parts of egg on NAFLD risk^[13].

An unexpected finding of the present study was that more than 4 eggs consumption per week was not significantly associated with risk of NAFLD. This may be explained by the fact that nutritional factors are correlated with each other, and determining of the effect of particular nutrients or particular foods on a risk factor is difficult. The effects of egg cholesterol on serum cholesterol concentrations depends on the content of individuals' diet specially the fiber content of it^[43,44]. It is possible that those who ate more than 4 eggs per week, consumed it in mixed dishes containing vegetables, which reduces the absorption of cholesterol. Thus, we




Figure 1 Multivariate-adjusted odds ratio for non-alcoholic fatty liver disease according to egg consumption. A: Crude model; B: Model 2, multivariate adjusted for age and energy intake; C: Model 3, further controlled for, body mass index, history of diabetes and smoking; D: Model 4, additionally adjusted for physical activity, and gender. Data are presented as the odds ratio (95%CI).

suggest that future studies assess the type of dishes with egg to find the possible interactions of different constituent of them.

It has been reported that dietary intake of patients with NAFLD was richer in saturated fat, cholesterol and was poorer in polyunsaturated fat^[12]. Subramanian *et al.*^[40] have concluded that dietary cholesterol confers in progression of NAFLD to NASH. Furthermore, Zelber-Sagi *et al.*^[18] found that NAFLD patients have a higher intake of meat, which is another source of dietary cholesterol; however, some other studies only found a significant association between NAFLD and high dietary intake of carbohydrate and simple sugars^[45,46], and some studies did find an association only between NAFLD and low intake of n-3 fatty acids and some antioxidants^[16]. These dietary habits may accelerate the development of NAFLD by directly affecting steatosis of liver and oxidative injury^[12].

This study was the first study that examined the relationship between egg consumption and risk of NAFLD in newly diagnosed patients who have not probably changed their diet due to the disease diagnosis; other strengths of this study includes its relatively large sample size, the high participation rate of participants, and socioeconomic differences of participants, which affects their dietary intakes.

Although we used a validated FFQ for measurement of dietary intakes, measurement error, and recall bias are unavoidable errors. Moreover, there might be some unknown risk factors that affect our results. Therefore, we recommend this analysis to be done in other populations.

In conclusion, our data indicate that egg consumption

in common amount of usage is associated with risk of NAFLD. According to the case-control design of this study, it can not show the causality effect; thus, these findings should be confirmed in future prospective studies with separate parts of eggs to find the etiological associations.

ACKNOWLEDGMENTS

This work was supported by a grant from "National Nutrition and Food Technology Research Institute" of Shahid Beheshti University of Medical Sciences, and Digestive Disease Research Institute, Tehran University of Medical Sciences, Tehran, Iran.

COMMENTS

Background

Among the known dietary factors that affect the pathogenesis of non-alcoholic fatty liver disease (NAFLD), dietary cholesterol has drawn a great deal of attention. Current studies propose that excess dietary cholesterol is regarded as the key factor related to the risk of steatohepatitis and hepatic inflammation. Among individual foods, eggs are regarded as a main source of dietary cholesterol; on the other hand, eggs are rich in proteins, and other nutrients. Limited research has assessed the relationship between egg consumption and risk of (NAFLD) development.

Research frontiers

Understanding of the association between egg consumption and risk of NAFLD development can contribute to clarify how intake of special food groups correlate with the disease and could lead to more particular guidelines for NAFLD prevention.

Innovations and breakthroughs

This study showed that egg consumption in common amount of usage is associated with risk of NAFLD. It seems that this association is mainly due to

high cholesterol content of it, and might not be seen when people consume only the white part of it.

Applications

According to the results of this study, the authors recommend low intake of eggs specially the yolk part of it for prevention of NAFLD; however, further studies are recommended to reach to a consensus in this regard.

Peer-review

This is an interesting paper evaluating the association between egg consumption and NAFLD.

REFERENCES

- 1 **Angulo P.** Nonalcoholic fatty liver disease. *N Engl J Med* 2002; **346**: 1221-1231 [PMID: 11961152 DOI: 10.1056/NEJMra011775]
- 2 **Vernon G, Baranova A, Younossi ZM.** Systematic review: the epidemiology and natural history of non-alcoholic fatty liver disease and non-alcoholic steatohepatitis in adults. *Aliment Pharmacol Ther* 2011; **34**: 274-285 [PMID: 21623852 DOI: 10.1111/j.1365-2036.2011.04724.x]
- 3 **Marchesini G, Brizi M, Bianchi G, Tomassetti S, Bugianesi E, Lenzi M, McCullough AJ, Natale S, Forlani G, Melchionda N.** Nonalcoholic fatty liver disease: a feature of the metabolic syndrome. *Diabetes* 2001; **50**: 1844-1850 [PMID: 11473047 DOI: 10.2337/diabetes.50.8.1844]
- 4 **Marchesini G, Bugianesi E, Forlani G, Cerrelli F, Lenzi M, Manini R, Natale S, Vanni E, Villanova N, Melchionda N, Rizzetto M.** Nonalcoholic fatty liver, steatohepatitis, and the metabolic syndrome. *Hepatology* 2003; **37**: 917-923 [PMID: 12668987 DOI: 10.1053/jhep.2003.50161]
- 5 **Papandreou D, Andreou E.** Role of diet on non-alcoholic fatty liver disease: An updated narrative review. *World J Hepatol* 2015; **7**: 575-582 [PMID: 25848481 DOI: 10.4254/wjh.v7.i3.575]
- 6 **Nseir W, Nassar F, Assy N.** Soft drinks consumption and nonalcoholic fatty liver disease. *World J Gastroenterol* 2010; **16**: 2579-2588 [PMID: 20518077 DOI: 10.3748/wjg.v16.i21.2579]
- 7 **Zelber-Sagi S, Godos J, Salomone F.** Lifestyle changes for the treatment of nonalcoholic fatty liver disease: a review of observational studies and intervention trials. *Therap Adv Gastroenterol* 2016; **9**: 392-407 [PMID: 27134667 DOI: 10.1177/1756283x16638830]
- 8 **Wouters K, van Gorp PJ, Bieghs V, Gijbels MJ, Duimel H, Lütjohann D, Kerksiek A, van Kruchten R, Maeda N, Staels B, van Bilsen M, Shiri-Sverdlov R, Hofker MH.** Dietary cholesterol, rather than liver steatosis, leads to hepatic inflammation in hyperlipidemic mouse models of nonalcoholic steatohepatitis. *Hepatology* 2008; **48**: 474-486 [PMID: 18666236 DOI: 10.1002/hep.22363]
- 9 **Kleemann R, Verschuren L, van Erk MJ, Nikolsky Y, Cnubben NH, Verheij ER, Smilde AK, Hendriks HF, Zadelaar S, Smith GJ, Kaznacheev V, Nikolskaya T, Melnikov A, Hurt-Camejo E, van der Greef J, van Ommen B, Kooistra T.** Atherosclerosis and liver inflammation induced by increased dietary cholesterol intake: a combined transcriptomics and metabolomics analysis. *Genome Biol* 2007; **8**: R200 [PMID: 17892536 DOI: 10.1186/gb-2007-8-9-r200]
- 10 **Savard C, Tartaglione EV, Kuver R, Haigh WG, Farrell GC, Subramanian S, Chait A, Yeh MM, Quinn LS, Ioannou GN.** Synergistic interaction of dietary cholesterol and dietary fat in inducing experimental steatohepatitis. *Hepatology* 2013; **57**: 81-92 [PMID: 22508243 DOI: 10.1002/hep.25789]
- 11 **Van Rooyen DM, Larter CZ, Haigh WG, Yeh MM, Ioannou G, Kuver R, Lee SP, Teoh NC, Farrell GC.** Hepatic free cholesterol accumulates in obese, diabetic mice and causes nonalcoholic steatohepatitis. *Gastroenterology* 2011; **141**: 1393-1403, 1403.e1-5 [PMID: 21703998 DOI: 10.1053/j.gastro.2011.06.040]
- 12 **Musso G, Gambino R, De Micheli F, Cassader M, Rizzetto M, Durazzo M, Fagà E, Silli B, Pagano G.** Dietary habits and their relations to insulin resistance and postprandial lipemia in non-alcoholic steatohepatitis. *Hepatology* 2003; **37**: 909-916 [PMID: 12668986 DOI: 10.1053/jhep.2003.50132]
- 13 **Yasutake K, Nakamuta M, Shima Y, Ohyama A, Masuda K, Haruta N, Fujino T, Aoyagi Y, Fukuizumi K, Yoshimoto T, Takemoto R, Miyahara T, Harada N, Hayata F, Nakashima M, Enjoji M.** Nutritional investigation of non-obese patients with non-alcoholic fatty liver disease: the significance of dietary cholesterol. *Scand J Gastroenterol* 2009; **44**: 471-477 [PMID: 19058085 DOI: 10.1080/00365520802588133]
- 14 **Enjoji M, Nakamuta M.** Is the control of dietary cholesterol intake sufficiently effective to ameliorate nonalcoholic fatty liver disease? *World J Gastroenterol* 2010; **16**: 800-803 [PMID: 20143458]
- 15 **Song WO, Kerver JM.** Nutritional contribution of eggs to American diets. *J Am Coll Nutr* 2000; **19**: 556S-562S [PMID: 11023007 DOI: 10.1080/07315724.2000.10718980]
- 16 **Han JM, Jo AN, Lee SM, Bae HS, Jun DW, Cho YK, Suk KT, Yoon JH, Ahn SB, Cho YJ, Kim SW, Jang EC.** Associations between intakes of individual nutrients or whole food groups and non-alcoholic fatty liver disease among Korean adults. *J Gastroenterol Hepatol* 2014; **29**: 1265-1272 [PMID: 24955455 DOI: 10.1111/jgh.12520]
- 17 **Shi L, Liu ZW, Li Y, Gong C, Zhang H, Song LJ, Huang CY, Li M.** The prevalence of nonalcoholic fatty liver disease and its association with lifestyle/dietary habits among university faculty and staff in Chengdu. *Biomed Environ Sci* 2012; **25**: 383-391 [PMID: 23026517 DOI: 10.3967/0895-3988.2012.04.002]
- 18 **Zelber-Sagi S, Nitzan-Kaluski D, Goldsmith R, Webb M, Blendis L, Halpern Z, Oren R.** Long term nutritional intake and the risk for non-alcoholic fatty liver disease (NAFLD): a population based study. *J Hepatol* 2007; **47**: 711-717 [PMID: 17850914 DOI: 10.1016/j.jhep.2007.06.020]
- 19 **Esfahani FH, Asghari G, Mirmiran P, Azizi F.** Reproducibility and relative validity of food group intake in a food frequency questionnaire developed for the Tehran Lipid and Glucose Study. *J Epidemiol* 2010; **20**: 150-158 [PMID: 20154450]
- 20 **Jaceldo-Siegl K, Knutsen SF, Sabaté J, Beeson WL, Chan J, Herring RP, Butler TL, Haddad E, Bennett H, Montgomery S, Sharma SS, Oda K, Fraser GE.** Validation of nutrient intake using an FFQ and repeated 24 h recalls in black and white subjects of the Adventist Health Study-2 (AHS-2). *Public Health Nutr* 2010; **13**: 812-819 [PMID: 19968897 DOI: 10.1017/s1368980009992072]
- 21 **Aadahl M, Jørgensen T.** Validation of a new self-report instrument for measuring physical activity. *Med Sci Sports Exerc* 2003; **35**: 1196-1202 [PMID: 12840642 DOI: 10.1249/01.mss.0000074446.02192.14]
- 22 **Kelishadi R, Rabiei K, Khosravi A, Famouri F, Sadeghi M, Rouhafza H, Shirani S.** Assessment of physical activity of adolescents in Isfahan. 2001
- 23 **Askari F, Rashidkhani B, Hekmatdoost A.** Cinnamon may have therapeutic benefits on lipid profile, liver enzymes, insulin resistance, and high-sensitivity C-reactive protein in nonalcoholic fatty liver disease patients. *Nutr Res* 2014; **34**: 143-148 [PMID: 24461315 DOI: 10.1016/j.nutres.2013.11.005]
- 24 **Emamat H, Foroughi F, Eimi-Zinab H, Taghizadeh M, Rismanchi M, Hekmatdoost A.** The effects of onion consumption on treatment of metabolic, histologic, and inflammatory features of nonalcoholic fatty liver disease. *J Diabetes Metab Disord* 2015; **15**: 25 [PMID: 27453880 DOI: 10.1186/s40200-016-0248-4]
- 25 **Eslamparast T, Eghtesad S, Hekmatdoost A, Poustchi H.** Probiotics and Nonalcoholic Fatty liver Disease. *Middle East J Dig Dis* 2013; **5**: 129-136 [PMID: 24829682]
- 26 **Eslamparast T, Eghtesad S, Poustchi H, Hekmatdoost A.** Recent advances in dietary supplementation, in treating non-alcoholic fatty liver disease. *World J Hepatol* 2015; **7**: 204-212 [PMID: 25729475 DOI: 10.4254/wjh.v7.i2.204]
- 27 **Eslamparast T, Poustchi H, Zamani F, Sharafkhan M, Malekzadeh R, Hekmatdoost A.** Synbiotic supplementation in nonalcoholic fatty liver disease: a randomized, double-blind, placebo-controlled pilot study. *Am J Clin Nutr* 2014; **99**: 535-542 [PMID: 24401715]

- DOI: 10.3945/ajcn.113.068890]
- 28 **Faghihzadeh F**, Adibi P, Hekmatdoost A. The effects of resveratrol supplementation on cardiovascular risk factors in patients with non-alcoholic fatty liver disease: a randomised, double-blind, placebo-controlled study. *Br J Nutr* 2015; **114**: 796-803 [PMID: 26234526 DOI: 10.1017/S0007114515002433]
 - 29 **Faghihzadeh F**, Adibi P, Rafiei R, Hekmatdoost A. Resveratrol supplementation improves inflammatory biomarkers in patients with nonalcoholic fatty liver disease. *Nutr Res* 2014; **34**: 837-843 [PMID: 25311610 DOI: 10.1016/j.nutres.2014.09.005]
 - 30 **Ghaemi A**, Taleban FA, Hekmatdoost A, Rafiei A, Hosseini V, Amiri Z, Homayounfar R, Fakheri H. How Much Weight Loss is Effective on Nonalcoholic Fatty Liver Disease? *Hepat Mon* 2013; **13**: e15227 [PMID: 24358045 DOI: 10.5812/hepatmon.15227]
 - 31 **Hekmatdoost A**, Shamsipour A, Meibodi M, Gheibizadeh N, Eslamparast T, Poustchi H. Adherence to the Dietary Approaches to Stop Hypertension (DASH) and risk of Nonalcoholic Fatty Liver Disease. *Int J Food Sci Nutr* 2016; **67**: 1024-1029 [PMID: 27436528 DOI: 10.1080/09637486.2016.1210101]
 - 32 **Mofidi F**, Yari Z, Poustchi H, Merat S, Nourinayyer B, Malekzadeh R, Hekmatdoost A. Effects of Synbiotics Supplementation in Lean Patients with Nonalcoholic Fatty Liver Disease: Study Protocol of a Pilot Randomized Double-blind Clinical Trial. *Arch Iran Med* 2016; **19**: 282-284 [PMID: 27041524]
 - 33 **Rahimlou M**, Ahmadnia H, Hekmatdoost A. Dietary supplements and pediatric non-alcoholic fatty liver disease: Present and the future. *World J Hepatol* 2015; **7**: 2597-2602 [PMID: 26557952 DOI: 10.4254/wjh.v7.i25.2597]
 - 34 **Rahimlou M**, Yari Z, Hekmatdoost A, Alavian SM, Keshavarz SA. Ginger Supplementation in Nonalcoholic Fatty Liver Disease: A Randomized, Double-Blind, Placebo-Controlled Pilot Study. *Hepat Mon* 2016; **16**: e34897 [PMID: 27110262 DOI: 10.5812/hepatmon.34897]
 - 35 **Shavakhi A**, Minakari M, Firouzian H, Assali R, Hekmatdoost A, Ferns G. Effect of a Probiotic and Metformin on Liver Amino-transferases in Non-alcoholic Steatohepatitis: A Double Blind Randomized Clinical Trial. *Int J Prev Med* 2013; **4**: 531-537 [PMID: 23930163]
 - 36 **Yari Z**, Rahimlou M, Eslamparast T, Ebrahimi-Daryani N, Poustchi H, Hekmatdoost A. Flaxseed supplementation in non-alcoholic fatty liver disease: a pilot randomized, open labeled, controlled study. *Int J Food Sci Nutr* 2016; **67**: 461-469 [PMID: 26983396 DOI: 10.3109/09637486.2016.1161011]
 - 37 **Miranda JM**, Anton X, Redondo-Valbuena C, Roca-Saavedra P, Rodriguez JA, Lamas A, Franco CM, Cepeda A. Egg and egg-derived foods: effects on human health and use as functional foods. *Nutrients* 2015; **7**: 706-729 [PMID: 25608941 DOI: 10.3390/nu7010706]
 - 38 **Weggemans RM**, Zock PL, Katan MB. Dietary cholesterol from eggs increases the ratio of total cholesterol to high-density lipoprotein cholesterol in humans: a meta-analysis. *Am J Clin Nutr* 2001; **73**: 885-891 [PMID: 11333841]
 - 39 **Baumgartner S**, Kelly ER, van der Made S, Berendschot TT, Husche C, Lütjohann D, Plat J. The influence of consuming an egg or an egg-yolk buttermilk drink for 12 wk on serum lipids, inflammation, and liver function markers in human volunteers. *Nutrition* 2013; **29**: 1237-1244 [PMID: 23911216 DOI: 10.1016/j.nut.2013.03.020]
 - 40 **Subramanian S**, Goodspeed L, Wang S, Kim J, Zeng L, Ioannou GN, Haigh WG, Yeh MM, Kowdley KV, O'Brien KD, Pennathur S, Chait A. Dietary cholesterol exacerbates hepatic steatosis and inflammation in obese LDL receptor-deficient mice. *J Lipid Res* 2011; **52**: 1626-1635 [PMID: 21690266 DOI: 10.1194/jlr.M016246]
 - 41 **Emamat H**, Noori M, Foroughi F, Rismanchi M, Eini-Zinab H, Hekmatdoost A. An Accessible and Pragmatic Experimental Model of Nonalcoholic Fatty Liver Disease. *Middle East J Dig Dis* 2016; **8**: 109-115 [PMID: 27252817 DOI: 10.15171/mejdd.2016.15]
 - 42 **Garcés-Rimón M**, González C, Uranga JA, López-Miranda V, López-Fandiño R, Miguel M. Pepsin Egg White Hydrolysate Ameliorates Obesity-Related Oxidative Stress, Inflammation and Steatosis in Zucker Fatty Rats. *PLoS One* 2016; **11**: e0151193 [PMID: 26985993 DOI: 10.1371/journal.pone.0151193]
 - 43 **Cohn JS**, Kamili A, Wat E, Chung RW, Tandy S. Reduction in intestinal cholesterol absorption by various food components: mechanisms and implications. *Atheroscler Suppl* 2010; **11**: 45-48 [PMID: 20439167 DOI: 10.1016/j.atherosclerosis.2010.04.004]
 - 44 **Gunness P**, Gidley MJ. Mechanisms underlying the cholesterol-lowering properties of soluble dietary fibre polysaccharides. *Food Funct* 2010; **1**: 149-155 [PMID: 21776465 DOI: 10.1039/c0fo00080a]
 - 45 **Solga S**, Alkhuraishe AR, Clark JM, Torbenson M, Greenwald A, Diehl AM, Magnuson T. Dietary composition and nonalcoholic fatty liver disease. *Dig Dis Sci* 2004; **49**: 1578-1583 [PMID: 15573908 DOI: 10.1023/B: DDAS.0000043367.69470.b7]
 - 46 **Toshimitsu K**, Matsuura B, Ohkubo I, Niiya T, Furukawa S, Hiasa Y, Kawamura M, Ebihara K, Onji M. Dietary habits and nutrient intake in non-alcoholic steatohepatitis. *Nutrition* 2007; **23**: 46-52 [PMID: 17140767 DOI: 10.1016/j.nut.2006.09.004]

P- Reviewer: Bellanti F, Tziomalos K **S- Editor:** Kong JX

L- Editor: A **E- Editor:** Li D





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.f6publishing.com/helpdesk>

<http://www.wjgnet.com>

