

Mechanisms of hepatic ischemia-reperfusion injury and protective effects of nitric oxide

Lian-Yue Guan, Pei-Yao Fu, Pei-Dong Li, Zhuo-Nan Li, Hong-Yu Liu, Min-Gang Xin, Wei Li

Lian-Yue Guan, Pei-Yao Fu, Pei-Dong Li, Zhuo-Nan Li, Hong-Yu Liu, Wei Li, Department of Hepatobiliary-Pancreatic Surgery, Third Hospital (China-Japan Union Hospital) of Jilin University, Changchun 130033, Jilin Province, China

Min-Gang Xin, Departments of Anesthesiology, Third Hospital (China-Japan Union Hospital) of Jilin University, Changchun 130033, Jilin Province, China

Author contributions: Guan LY and Fu PY contributed equally to the paper; Guan LY primarily drafted the article; Fu PY performed the intensive revision and drew the two figures; Li PD drafted the nitric oxide portion of the manuscript; Li ZN, Liu HY and Xin MG conducted the data collection and interpretation, selected the references, and drafted the discussion; Li W contributed to the concept design, critical revision and finalization of the manuscript.

Supported by National Natural Science Foundation of China, No. 81170416 and No. 81273264; Doctoral Fund of Ministry of Education of China, No. 20100061110069; Jilin Province Science and Technology Bureau International Cooperation Fund, No. 2011742; Techpool Research Fund, No. 01201046; Jilin Province Nature Science Foundation, No. 201015178

Correspondence to: Wei Li, MD, PhD, Department of Hepatobiliary-Pancreatic Surgery, Third Hospital (China-Japan Union Hospital) of Jilin University, 126 Xiantai Street, Changchun 130033, Jilin Province, China. weili888@hotmail.com

Telephone: +86-431-89876816 Fax: +86-431-84641026

Received: January 4, 2014 Revised: May 26, 2014

Accepted: June 20, 2014

Published online: July 27, 2014

Abstract

Hepatic ischemia-reperfusion injury (IRI) is a pathological event post liver surgery or transplantation and significantly influences the prognosis of liver function. The mechanisms of IRI remain unclear, and effective methods are lacking for the prevention and therapy of IRI. Several factors/pathways have been implicated in the hepatic IRI process, including anaerobic metabolism, mitochondria, oxidative stress, intracellular calcium overload, liver Kupffer cells and neutrophils, and cytokines and chemokines. The role of nitric oxide (NO)

in protecting against liver IRI has recently been reported. NO has been found to attenuate liver IRI through various mechanisms including reducing hepatocellular apoptosis, decreasing oxidative stress and leukocyte adhesion, increasing microcirculatory flow, and enhancing mitochondrial function. The purpose of this review is to provide insights into the mechanisms of liver IRI, indicating the potential protective factors/pathways that may help to improve therapeutic regimens for controlling hepatic IRI during liver surgery, and the potential therapeutic role of NO in liver IRI.

© 2014 Baishideng Publishing Group Inc. All rights reserved.

Key words: Liver; Ischemia-reperfusion injury; Cytokine; Chemokine; Kupffer cells; Mitochondria; Nitric oxide

Core tip: This review provides insights into several key mechanisms of liver ischemia-reperfusion injury, including the effects of anaerobic metabolism and the role of mitochondria, oxidative stress, intracellular calcium overload, liver Kupffer cells and neutrophils, and cytokines and chemokines; and summarizes the protective effects of nitric oxide.

Guan LY, Fu PY, Li PD, Li ZN, Liu HY, Xin MG, Li W. Mechanisms of hepatic ischemia-reperfusion injury and protective effects of nitric oxide. *World J Gastrointest Surg* 2014; 6(7): 122-128 Available from: URL: <http://www.wjgnet.com/1948-9366/full/v6/i7/122.htm> DOI: <http://dx.doi.org/10.4240/wjgs.v6.i7.122>

INTRODUCTION

In recent years, liver resection and liver transplantation have been widely adopted in clinical practice for the treatment of liver diseases. Hepatic ischemia-reperfusion injury (IRI) occurs substantially during liver resection or

transplantation and remains a major cause of liver non-function or functional failure following liver surgery. This non-negligible injury has become a bottleneck which has restricted the use of marginal liver donors and the development of extensive liver resection. Hepatic IRI includes both warm and cold IRI - two types that share similar pathophysiological processes. The mechanisms of liver IRI have been widely investigated, but nevertheless remain largely unclear. The factors/pathways have been implicated in the hepatic IRI process include anaerobic metabolism, mitochondria, oxidative stress, intracellular calcium overload, liver Kupffer cells (KCs) and neutrophils, and cytokines and chemokines. More importantly, an effective prevention or treatment method is still lacking. Therefore, an effective method for preventing or minimizing hepatic IRI during liver surgery is urgently needed. A better understanding of the mechanisms in the development of IRI will provide insights into improving the treatment regimen for IRI. In this review, the authors comprehensively discuss the mechanisms of liver IRI and describe the role of nitric oxide (NO) in protecting the liver from IRI.

ANAEROBIC METABOLISM AND ACIDOSIS

IRI exerts wide-ranging metabolic effects on the body. During the state of hepatic ischemia, the metabolic pattern is shifted from aerobic to anaerobic, the redox process of the hepatocytes is blocked, adenosine triphosphate (ATP)-dependent cellular metabolic activities are gradually stopped, and intracellular ATP is rapidly depleted. Conversely, there is accumulation of acidic metabolites, such as lactic acid and ketone bodies, which is caused by enhanced anaerobic glycolysis. This is accompanied by hypofunction of mitochondrial oxidative phosphorylation, resulting in the decrease of pH values between tissues and cells, known as metabolic acidosis. Studies have shown that this change plays a role in protecting the liver cells^[1,2]. However, the pH values restore to normal after reperfusion, and further enhance pH-dependent enzyme activation, such as activation of proteases and phospholipases, further worsening the damage of tissues and organs. This is called the pH paradox^[3]. The toxicity of acidic metabolites caused by a lower ATP supply mainly impairs the cellular functions of homeostasis, signaling interactions, and sodium/potassium ATPase (Na^+/K^+ -ATPase), causing mitochondrial damage and resulting in microcirculation failure and cellular destruction^[4].

ROLE OF MITOCHONDRIA

IRI exerts effects not only on the body as a whole, but also at the cellular level. The mitochondria are the location where oxidative phosphorylation mainly takes place, and the mitochondria participate in multiple pathophysiological processes of IRI. A large number of reactive

oxygen species (ROS) and reactive nitrogen species are generated in the mitochondria during the state of ischemia. Hypoxia undermines the process of oxidative phosphorylation in cells and obstructs the production of ATP, causing disorders of the cytoplasmic ions such as Ca^{2+} , Na^+ , and H^+ in the mitochondria, and finally leads to mitochondrial membrane permeability transition (MMPT)^[5]. MMPT is manifested primarily by mitochondrial swelling and the decline of membrane potential^[6], which allows soluble molecules of a molecular weight less than 1500 kDa to freely pass through the inner mitochondrial membrane, the so-called "mitochondrial megachannel"^[7]. Many studies have indicated that MMPT is related to the process of hepatocyte damage after IRI^[5,8].

OXIDATIVE STRESS

IRI has many biochemical ramifications. It has been shown that oxidative stress plays a key role in reperfusion injury. Many highly reactive molecules, such as ROS, are induced during the period of hepatic IRI. ROS include superoxide anions, hydroxyl radicals, and peroxide hydrogen, and mainly act on proteins, enzymes, nucleic acids, cytoskeleton, and lipid peroxides, leading to mitochondrial dysfunction and lipid peroxidation^[9]. ROS can also damage endothelial cells and destroy the integrity of the microvasculature. ROS can be reduced or overcome by reducing the blood flow and applying endogenous antioxidants, such as superoxide dismutase, catalase, glutathione, vitamin E, or beta-carotene^[10]. On the other hand, application of recombinant adenovirus superoxide has been shown to effectively reduce hepatic IRI in mice^[11].

INTRACELLULAR CALCIUM OVERLOAD

Among the biochemical factors affected by IRI, calcium has an especially important role. The electrochemical gradient of the calcium ion plays an important role in maintaining homeostasis of physical calcium (Ca^{2+}). If the calcium level is elevated when ischemia or hypoxia, oxidative stress, toxic substance release or other harmful events occur, this is called Ca^{2+} overload. Intracellular Ca^{2+} overload can activate Ca^{2+} -dependent enzymes such as calpains, protein kinase C, and phospholipase C, and ultimately leads to cell death or apoptosis. Recent studies have shown that the increased amount of intracellular Ca^{2+} is not uniform, but is a local phenomenon. Non-specific calcium channel blockers can inhibit the elevation of intracellular Ca^{2+} and reduce cellular damage, demonstrating that Ca^{2+} influx may play a major role in the IRI process^[12,13].

KCS AND NEUTROPHILS

It has been demonstrated that liver KCs and neutrophils are involved in the hepatic IRI process. The KCs mainly mediate liver ischemic injury in the earlier stage of reperfusion (within 2 h) by synthesizing and releasing




Figure 1 Cytokine network on the regulation of liver ischemia-reperfusion injury. IRI: Ischemia-reperfusion injury; IL: Interleukin; IFN- γ : Interferon-gamma; HGF: Hepatocyte growth factor; MIP: Macrophage inflammatory protein; ICAM-1: Intercellular adhesion molecule 1; NF: Nuclear factor; MAPK: Mitogen-activated protein kinase.

ROS and the pro-inflammatory cytokines tumor necrosis factor- α (TNF- α) and interleukin (IL)-1 β to further activate liver sinusoidal endothelial cells, enhance the expression of the adhesion molecules intercellular adhesion molecule 1 (ICAM-1)/vascular cell adhesion molecule 1 (VCAM-1), further promote the adhesion, migration, and chemotaxis of neutrophils and endothelial cells, and accumulate and activate neutrophils, resulting in subsequent liver cell damage^[14]. Studies have shown that endotoxins are also involved in the process of liver IRI^[10,15]. Blocking KC activation by the use of gadolinium chloride or methyl palmitate can reduce acute liver cell injury significantly. Activation of neutrophils can directly damage liver cells by the release of oxidants and proteases after reperfusion. Ultimately, myeloperoxidase (halide form, such as Cl) released from neutrophils changes hydrogen peroxide (H₂O₂) into hypochlorous acid (HOCl), which is a potent oxidant. These oxidants can directly cause liver cell damage and/or induce protease-mediated injury through inactivation of the endogenous anti-protease system^[15,16], suggesting that anti-oxidant or anti-protease therapy would be helpful for preventing IRI.

ROLE OF CYTOKINES AND CHEMOKINES

Cytokines play a dual role of anti-inflammatory and pro-inflammatory responses in the process of liver IRI (Figure 1). TNF- α is a key member of the group of endogenous pro-inflammatory and anti-inflammatory molecules, and is a critical factor in triggering the inflammatory cascade.

It is secreted by activated KCs and impacts liver tissue and distant organs through paracrine signaling and the endocrine system^[17]. TNF- α can bind to the receptors on the surface of liver cells to induce overproduction of the chemokine epithelial neutrophil activating protein-78 (ENA-78) and ROS, activate nuclear factor (NF)- κ B, mitogen-activated protein kinase, and c-Jun N-terminal kinase (JNK), and cause liver injury directly^[18]. In addition, TNF- α also can upregulate expression of the chemokines ICAM-1, VCAM-1 and P-selectin^[19]. Moreover, JNK and ROS can directly act on liver cells to cause liver damage.

In addition to TNF- α , the other important cytokines involved in liver IRI are interferon-gamma (IFN- γ), IL-1 β , IL-6, IL-12, IL-23, IL-10, IL-13, vascular endothelial growth factor (VEGF), and hepatocyte growth factor (HGF). These cytokines promote leukocyte activation in the liver after ischemia through various pathways. IFN- γ is mainly produced by T cells and natural killer T cells, and activated by toll-like receptor-4 and IL-12. IFN- γ can either aggravate liver damage or reduce liver damage through enhancing or downregulating neutrophil accumulation and activation in a dose-dependent manner^[20]. IL-1 β , IL-6, IL-12, and IL-23 are mainly produced by KCs and hepatocytes. IL-1 β can upregulate NO synthesis through the protein kinase B (Akt), NF- κ B, and inducible nitric oxide synthase (iNOS) pathways. IL-1 β can further upregulate leukocyte aggregation and adhesion by activating NF- κ B and macrophage inflammatory protein (MIP)-2, thus damaging the liver cells^[21]. IL-12 and IL-23 can also increase TNF- α production by activating




Figure 2 The protective effects of nitric oxide on liver ischemia-reperfusion injury. ATP: Adenosine triphosphate; IL: Interleukin.

NF- κ B and signal transducer and activator of transcription (STAT)-4, and further stimulating CD4 T cells to produce IL-17, ensuring the accumulation of neutrophils and aggravating liver damage^[22].

On the contrary, IL-6 can activate STAT-3, upregulate glutathione (GSH) expression, and downregulate oxidative stress markers, thus reducing hepatocyte damage and promoting hepatocyte proliferation^[23]. IL-10 and IL-13 are mainly produced by KCs and T lymphocytes, and also play a role in alleviating liver damage and promoting liver regeneration. The protective role of IL-10 and IL-13 is mainly mediated by upregulation of heme oxygenase (HO)-1, B-cell lymphoma (Bcl)-2/bcl-x, and downregulation of NF- κ B, IL-1 β , IL-2, IFN- γ , MIP-2, cytokine-induced neutrophil chemotaxin, E-selectin, and neutrophil aggregation^[24,25].

VEGF can be produced by many types of cells including KCs, T cells, sinusoidal endothelial cells and hepatocytes. It plays dual functions in liver IRI. IRI triggers the VEGF receptor and Src tyrosine kinase activation, and upregulates the expression of TNF- α , INF- γ , monocyte chemoattractant protein-1 and E-selectin, all of which result in the accumulation of intrahepatic T lymphocytes, macrophages and neutrophils, producing liver damage. On the other hand, exogenous administration of VEGF can upregulate iNOS production and protect the liver from IRI^[26].

HGF is produced by liver non-parenchymal cells, mainly KCs. HGF can increase hepatocyte DNA synthesis, proliferation, and glutathione expression, downregulate the expression of the oxidative stress marker ICAM-1 in sinusoidal endothelial cells, and inhibit cytokine-induced neutrophil chemotaxin and neutrophil permeability, further reducing liver damage and promoting liver cell proliferation^[27].

PROTECTIVE ROLE OF NITRIC OXIDE

The effects of NO in protecting the liver from IRI have

been studied extensively in recent years. NO is a highly reactive free radical produced from L-arginine and oxygen by nitric oxide synthase (NOS) *in vivo*^[28]. Many studies have demonstrated that NO is a versatile signaling mediator involved in a multitude of critical cellular events, such as inhibition of platelet aggregation, regulation of the microcirculation, and inhibition of caspase activities to prevent cell apoptosis^[29,30]. It has been shown that both endogenously generated and exogenously administered NO plays an important role in protecting the liver from IRI^[31]. NO has been found to attenuate liver IRI through various mechanisms, including the protection of hepatocytes from apoptosis and the reduction of macrophage infiltration^[32]. Complicated mechanisms and numerous molecules are involved in exerting the protective effects of NO against liver IRI, including ATP molecules, endothelin, adhesion molecules, cytokines, free radical species, and antioxidants^[33] (Figure 2). NO has been shown to potentiate hepatic ATP levels, reduce oxidative damage, prevent the reduction of antioxidants such as glutathione, and reduce the adverse effects of endothelin during liver IRI^[33,34]. Studies have demonstrated that NO affects cellular decisions of life and death by either turning on or shutting off apoptotic pathways, suggesting that NO can function differently depending on the dose and duration of exposure^[35,36]. Large amounts of NO may in turn paradoxically damage liver tissue by forming nitrogen peroxide^[37], suggesting that the therapeutic safety window of NO is limited.

NO-based therapy has been applied for many years to patients with pulmonary hypertension or cardiopulmonary disorders. The therapeutic application of NO in protecting the liver from IRI has just been emerging. A prospective randomized small group trial with liver transplant patients has demonstrated that NO inhalation in liver recipients during the perioperative period of liver transplantation significantly protects hepatocytes from apoptotic death, accelerates the restoration of liver graft function, and reduces hospital length of stay^[38]. Since NO has a very short half-life *in vivo*, it is not an ideal gas




Figure 3 Mechanisms of hepatic ischemia reperfusion injury. ATP: Adenosine triphosphate; IL: Interleukin; ROS: Reactive oxygen species; IRI: Ischemia-reperfusion injury; IFN- γ : Interferon-gamma; ICAM: Intercellular adhesion molecule; VCAM: Vascular cell adhesion molecule; TNF: Tumor necrosis factor.

for the treatment of IRI. NO drugs administered to liver donors, such as organic nitrates and sodium nitropruside, are now being explored as an alternative choice for NO delivery.

Sodium nitrite, a storage form of NO, can release NO during hypoxia and acidosis^[39]. Sodium nitrite has now been identified as an important storage reservoir of bioavailable NO in the blood and tissues^[40]. The reduction of nitrite to NO has been demonstrated to confer cytoprotection against IRI in the heart, liver, brain, and kidney^[40]. Interventions that increase NO production by the use of sodium nitrite before the occurrence of ischemia, either through intraperitoneal injection or oral administration, can mediate significant cytoprotection. This strategy has been demonstrated to potently limit acute IRI in both the heart and liver in murine warm IRI models, with the ability to decrease myocardial infarction and hepatocyte apoptosis^[40-43].

NO is also an important effector molecule, produced by KCs and dendritic cells (DCs), and is involved in immune regulation and host innate and adaptive immunity^[44]. NO inhibits proinflammatory cytokines, including TNF- α , IL-1 β , IL-1 α and IL-12, which may induce the inflammatory cascade during liver IRI^[24-26,33]. It has been reported that NO exerts multiple effects on immune cells, decreasing the number of T helper (Th)1 cells and augmenting Th2 cell proliferation and their cytokine synthesis, regulating leukocyte adhesion and recruitment to the site of infection^[45-47], inhibiting Th1 proliferation, and promoting T cell apoptosis^[48,49]. Moreover, NO also contributes to the immunosuppressive function of induced T regulatory cells (Treg)^[50]. Therefore, NO is involved in the regulation of liver IRI-associated immune responses. The underlying mechanisms are largely unknown and warrant further investigation.

CONCLUSION

Hepatic IRI is not only a pathophysiological process

involving the liver itself, but also a complex systemic process affecting multiple tissues and organs. Hepatic IRI can seriously impair liver function, even producing irreversible damage, which causes a cascade of multiple organ dysfunction. Many factors, including anaerobic metabolism, mitochondrial damage, oxidative stress, intracellular Ca²⁺ overload, cytokines and chemokines produced by KCs and neutrophils, and NO, are all involved in the regulation of liver IRI processes. The most important pathways of liver IRI are initiated by oxidative stress, anaerobic metabolism and acidosis, further resulting in the cellular damage through induction of apoptosis, immune responses, and cytokine regulations (Figure 3). Inhaled NO or NO-producing drugs have shown positive effects on IRI protection in clinical practice, and may be a good choice for liver IRI therapy in the future. Therefore, further exploration of the mechanisms of IRI on animal models focusing on the regulatory pathway of IRI development, with concomitant development of a more effective method of controlling IRI, will help overcome the challenges in the prevention of IRI and therapeutic strategies.

REFERENCES

- 1 **Kanoria S**, Glantzounis G, Quaglia A, Dinesh S, Fusai G, Davidson BR, Seifalian AM. Remote preconditioning improves hepatic oxygenation after ischaemia reperfusion injury. *Transpl Int* 2012; **25**: 783-791 [PMID: 22533545 DOI: 10.1111/j.1432-2277.2012.01481.x]
- 2 **Guan YF**, Pritts TA, Montrose MH. Ischemic post-conditioning to counteract intestinal ischemia/reperfusion injury. *World J Gastrointest Pathophysiol* 2010; **1**: 137-143 [PMID: 21607154 DOI: 10.4291/wjgp.v1.i4.137]
- 3 **Datta G**, Fuller BJ, Davidson BR. Molecular mechanisms of liver ischemia reperfusion injury: insights from transgenic knockout models. *World J Gastroenterol* 2013; **19**: 1683-1698 [PMID: 23555157 DOI: 10.3748/wjg.v19.i11.1683]
- 4 **Siriusawakul A**, Zaky A, Lang JD. Role of nitric oxide in hepatic ischemia-reperfusion injury. *World J Gastroenterol* 2010; **16**: 6079-6086 [PMID: 21182222 DOI: 10.3748/wjg.v16.i48.6079]

- 5 **Sastre J**, Serviddio G, Pereda J, Minana JB, Arduini A, Vendemiale G, Poli G, Pallardo FV, Vina J. Mitochondrial function in liver disease. *Front Biosci* 2007; **12**: 1200-1209 [PMID: 17127373 DOI: 10.2741/2138]
- 6 **Aguilar HI**, Botla R, Arora AS, Bronk SF, Gores GJ. Induction of the mitochondrial permeability transition by protease activity in rats: a mechanism of hepatocyte necrosis. *Gastroenterology* 1996; **110**: 558-566 [PMID: 8566604 DOI: 10.1053/gast.1996.v110.pm8566604]
- 7 **Bernardi P**, Broekemeier KM, Pfeiffer DR. Recent progress on regulation of the mitochondrial permeability transition pore; a cyclosporin-sensitive pore in the inner mitochondrial membrane. *J Bioenerg Biomembr* 1994; **26**: 509-517 [PMID: 7896766 DOI: 10.1007/BF00762735]
- 8 **Abu-Amara M**, Yang SY, Tapuria N, Fuller B, Davidson B, Seifalian A. Liver ischemia/reperfusion injury: processes in inflammatory networks--a review. *Liver Transpl* 2010; **16**: 1016-1032 [PMID: 20818739 DOI: 10.1002/lt.22117]
- 9 **Brass CA**, Roberts TG. Hepatic free radical production after cold storage: Kupffer cell-dependent and -independent mechanisms in rats. *Gastroenterology* 1995; **108**: 1167-1175 [PMID: 7698585 DOI: 10.1016/0016-5085(95)90216-3]
- 10 **Jaeschke H**. Reactive oxygen and ischemia/reperfusion injury of the liver. *Chem Biol Interact* 1991; **79**: 115-136 [PMID: 1884426 DOI: 10.1016/0009-2797(91)90077-K]
- 11 **McCord JM**. Oxygen-derived free radicals in postischemic tissue injury. *N Engl J Med* 1985; **312**: 159-163 [PMID: 2981404 DOI: 10.1056/NEJM198501173120305]
- 12 **Ikeda M**, Ariyoshi H, Sakon M, Kambayashi J, Yoshikawa N, Shinoki N, Kawasaki T, Monden M. A role for local calcium gradients upon hypoxic injury in human umbilical vein endothelial cells (HUVEC). *Cell Calcium* 1998; **24**: 49-57 [PMID: 9793688 DOI: 10.1016/S0143-4160(98)90088-4]
- 13 **Wang HG**, Pathan N, Ethell IM, Krajewski S, Yamaguchi Y, Shibasaki F, McKeon F, Bobo T, Franke TF, Reed JC. Ca²⁺-induced apoptosis through calcineurin dephosphorylation of BAD. *Science* 1999; **284**: 339-343 [PMID: 10195903 DOI: 10.1126/science.284.5412.339]
- 14 **Zhou W**, Zhang Y, Hosch MS, Lang A, Zwacka RM, Engelhardt JF. Subcellular site of superoxide dismutase expression differentially controls AP-1 activity and injury in mouse liver following ischemia/reperfusion. *Hepatology* 2001; **33**: 902-914 [PMID: 11283855 DOI: 10.1053/jhep.2001.23073]
- 15 **Boury NM**, Czuprynski CJ. Listeria monocytogenes infection increases neutrophil adhesion and damage to a murine hepatocyte cell line in vitro. *Immunol Lett* 1995; **46**: 111-116 [PMID: 7590905 DOI: 10.1016/0165-2478(95)00027-3]
- 16 **Nagendra AR**, Mickelson JK, Smith CW. CD18 integrin and CD54-dependent neutrophil adhesion to cytokine-stimulated human hepatocytes. *Am J Physiol* 1997; **272**: G408-G416 [PMID: 9124560]
- 17 **Gujral JS**, Bucci TJ, Farhood A, Jaeschke H. Mechanism of cell death during warm hepatic ischemia-reperfusion in rats: apoptosis or necrosis? *Hepatology* 2001; **33**: 397-405 [PMID: 11172341 DOI: 10.1053/jhep.2001.22002]
- 18 **Redaelli CA**, Tian YH, Schaffner T, Ledermann M, Baer HU, Dufour JF. Extended preservation of rat liver graft by induction of heme oxygenase-1. *Hepatology* 2002; **35**: 1082-1092 [PMID: 11981758 DOI: 10.1053/jhep.2002.33067]
- 19 **Peralta C**, Fernández L, Panés J, Prats N, Sans M, Piqué JM, Gelpí E, Roselló-Catafau J. Preconditioning protects against systemic disorders associated with hepatic ischemia-reperfusion through blockade of tumor necrosis factor-induced P-selectin up-regulation in the rat. *Hepatology* 2001; **33**: 100-113 [PMID: 11124826 DOI: 10.1053/jhep.2001.20529]
- 20 **Schwabe RF**, Brenner DA. Mechanisms of Liver Injury. I. TNF-alpha-induced liver injury: role of IKK, JNK, and ROS pathways. *Am J Physiol Gastrointest Liver Physiol* 2006; **290**: G583-G589 [PMID: 16537970]
- 21 **Oe S**, Hirotsu T, Fujii H, Yasuchika K, Nishio T, Iimuro Y, Morimoto T, Nagao M, Yamaoka Y. Continuous intravenous infusion of deleted form of hepatocyte growth factor attenuates hepatic ischemia-reperfusion injury in rats. *J Hepatol* 2001; **34**: 832-839 [PMID: 11451166 DOI: 10.1016/S0168-8278(01)00030-7]
- 22 **Hamada T**, Tsuchihashi S, Avanesyan A, Duarte S, Moore C, Busuttill RW, Coito AJ. Cyclooxygenase-2 deficiency enhances Th2 immune responses and impairs neutrophil recruitment in hepatic ischemia/reperfusion injury. *J Immunol* 2008; **180**: 1843-1853 [PMID: 18209082 DOI: 10.4049/jimmunol.180.3.1843]
- 23 **Welborn MB**, Moldawer LL, Seeger JM, Minter RM, Huber TS. Role of endogenous interleukin-10 in local and distant organ injury after visceral ischemia-reperfusion. *Shock* 2003; **20**: 35-40 [PMID: 12813366 DOI: 10.1097/01/SHK.0000071062.67193.b6]
- 24 **Teoh N**, Field J, Farrell G. Interleukin-6 is a key mediator of the hepatoprotective and pro-proliferative effects of ischaemic preconditioning in mice. *J Hepatol* 2006; **45**: 20-27 [PMID: 16600417 DOI: 10.1016/j.jhep.2006.01.039]
- 25 **Husted TL**, Blanchard J, Schuster R, Shen H, Lentsch AB. Potential role for IL-23 in hepatic ischemia/reperfusion injury. *Inflamm Res* 2006; **55**: 177-178 [PMID: 16830103 DOI: 10.1007/s00011-006-0073-1]
- 26 **Ke B**, Shen XD, Lassman CR, Gao F, Busuttill RW, Kupiec-Weglinski JW. Cytoprotective and antiapoptotic effects of IL-13 in hepatic cold ischemia/reperfusion injury are heme oxygenase-1 dependent. *Am J Transplant* 2003; **3**: 1076-1082 [PMID: 12919086 DOI: 10.1034/j.1600-6143.2003.00147.x]
- 27 **Shen XD**, Ke B, Zhai Y, Gao F, Tsuchihashi S, Lassman CR, Busuttill RW, Kupiec-Weglinski JW. Absence of toll-like receptor 4 (TLR4) signaling in the donor organ reduces ischemia and reperfusion injury in a murine liver transplantation model. *Liver Transpl* 2007; **13**: 1435-1443 [PMID: 17902130 DOI: 10.1002/lt.21251]
- 28 **Diesen DL**, Kuo PC. Nitric oxide and redox regulation in the liver: Part I. General considerations and redox biology in hepatitis. *J Surg Res* 2010; **162**: 95-109 [PMID: 20444470 DOI: 10.1016/j.jss.2009.09.019]
- 29 **Pacher P**, Beckman JS, Liaudet L. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* 2007; **87**: 315-424 [PMID: 17237348 DOI: 10.1152/physrev.00029.2006]
- 30 **Mocellin S**, Bronte V, Nitti D. Nitric oxide, a double edged sword in cancer biology: searching for therapeutic opportunities. *Med Res Rev* 2007; **27**: 317-352 [PMID: 16991100 DOI: 10.1002/med.20092]
- 31 **Abu-Amara M**, Yang SY, Seifalian A, Davidson B, Fuller B. The nitric oxide pathway--evidence and mechanisms for protection against liver ischaemia reperfusion injury. *Liver Int* 2012; **32**: 531-543 [PMID: 22316165 DOI: 10.1111/j.1478-3231.2012.02755.x]
- 32 **Koti RS**, Yang W, Dashwood MR, Davidson BR, Seifalian AM. Effect of ischemic preconditioning on hepatic microcirculation and function in a rat model of ischemia reperfusion injury. *Liver Transpl* 2002; **8**: 1182-1191 [PMID: 12474159 DOI: 10.1053/jlts.2002.36846]
- 33 **Liu P**, Xu B, Spokas E, Lai PS, Wong PY. Role of endogenous nitric oxide in TNF-alpha and IL-1beta generation in hepatic ischemia-reperfusion. *Shock* 2000; **13**: 217-223 [PMID: 10718379 DOI: 10.1097/00024382-200003000-00008]
- 34 **Hsu CM**, Wang JS, Liu CH, Chen LW. Kupffer cells protect liver from ischemia-reperfusion injury by an inducible nitric oxide synthase-dependent mechanism. *Shock* 2002; **17**: 280-285 [PMID: 11954827 DOI: 10.1097/00024382-200204000-00007]
- 35 **Brüne B**. Nitric oxide: NO apoptosis or turning it ON? *Cell Death Differ* 2003; **10**: 864-869 [PMID: 12867993 DOI: 10.1038/sj.cdd.4401261]
- 36 **Bolli R**. Cardioprotective function of inducible nitric oxide synthase and role of nitric oxide in myocardial ischemia and preconditioning: an overview of a decade of research.

- J Mol Cell Cardiol* 2001; **33**: 1897-1918 [PMID: 11708836 DOI: 10.1006/jmcc.2001.1462]
- 37 **Miyake T**, Yokoyama Y, Kokuryo T, Mizutani T, Imamura A, Nagino M. Endothelial nitric oxide synthase plays a main role in producing nitric oxide in the superacute phase of hepatic ischemia prior to the upregulation of inducible nitric oxide synthase. *J Surg Res* 2013; **183**: 742-751 [PMID: 23485075 DOI: 10.1016/j.jss.2013.01.048]
- 38 **Lang JD**, Teng X, Chumley P, Crawford JH, Isbell TS, Chacko BK, Liu Y, Jhala N, Crowe DR, Smith AB, Cross RC, Frenette L, Kelley EE, Wilhite DW, Hall CR, Page GP, Fallon MB, Bynon JS, Eckhoff DE, Patel RP. Inhaled NO accelerates restoration of liver function in adults following orthotopic liver transplantation. *J Clin Invest* 2007; **117**: 2583-2591 [PMID: 17717604 DOI: 10.1172/JCI31892]
- 39 **Bryan NS**, Rassaf T, Maloney RE, Rodriguez CM, Saijo F, Rodriguez JR, Feelisch M. Cellular targets and mechanisms of nitrosylation: an insight into their nature and kinetics in vivo. *Proc Natl Acad Sci USA* 2004; **101**: 4308-4313 [PMID: 15014175 DOI: 10.1073/pnas.0306706101]
- 40 **Sinha SS**, Shiva S, Gladwin MT. Myocardial protection by nitrite: evidence that this reperfusion therapeutic will not be lost in translation. *Trends Cardiovasc Med* 2008; **18**: 163-172 [PMID: 18790386 DOI: 10.1016/j.tcm.2008.05.001]
- 41 **Duranski MR**, Greer JJ, Dejam A, Jaganmohan S, Hogg N, Langston W, Patel RP, Yet SF, Wang X, Kevil CG, Gladwin MT, Lefer DJ. Cytoprotective effects of nitrite during in vivo ischemia-reperfusion of the heart and liver. *J Clin Invest* 2005; **115**: 1232-1240 [PMID: 15841216 DOI: 10.1172/JCI22493]
- 42 **Kumar D**, Branch BG, Pattillo CB, Hood J, Thoma S, Simpson S, Illum S, Arora N, Chidlow JH, Langston W, Teng X, Lefer DJ, Patel RP, Kevil CG. Chronic sodium nitrite therapy augments ischemia-induced angiogenesis and arteriogenesis. *Proc Natl Acad Sci USA* 2008; **105**: 7540-7545 [PMID: 18508974 DOI: 10.1073/pnas.0711480105]
- 43 **Li W**, Meng Z, Liu Y, Patel RP, Lang JD. The hepatoprotective effect of sodium nitrite on cold ischemia-reperfusion injury. *J Transplant* 2012; **2012**: 635179 [PMID: 22530108 DOI: 10.1155/2012/635179]
- 44 **Panjwani NN**, Popova L, Srivastava PK. Heat shock proteins gp96 and hsp70 activate the release of nitric oxide by APCs. *J Immunol* 2002; **168**: 2997-3003 [PMID: 11884472 DOI: 10.4049/jimmunol.168.6.2997]
- 45 **Chen C**, Lee WH, Zhong L, Liu CP. Regulatory T cells can mediate their function through the stimulation of APCs to produce immunosuppressive nitric oxide. *J Immunol* 2006; **176**: 3449-3460 [PMID: 16517713 DOI: 10.4049/jimmunol.176.6.3449]
- 46 **Taylor-Robinson AW**, Liew FY, Severn A, Xu D, McSorley SJ, Garside P, Padron J, Phillips RS. Regulation of the immune response by nitric oxide differentially produced by T helper type 1 and T helper type 2 cells. *Eur J Immunol* 1994; **24**: 980-984 [PMID: 8149966 DOI: 10.1002/eji.1830240430]
- 47 **Fox-Robichaud A**, Payne D, Hasan SU, Ostrovsky L, Fairhead T, Reinhardt P, Kubes P. Inhaled NO as a viable anti-adhesive therapy for ischemia/reperfusion injury of distal microvascular beds. *J Clin Invest* 1998; **101**: 2497-2505 [PMID: 9616221 DOI: 10.1172/JCI2736]
- 48 **Roland CR**, Walp L, Stack RM, Flye MW. Outcome of Kupffer cell antigen presentation to a cloned murine Th1 lymphocyte depends on the inducibility of nitric oxide synthase by IFN-gamma. *J Immunol* 1994; **153**: 5453-5464 [PMID: 7527442]
- 49 **Monsonogo A**, Imitola J, Zota V, Oida T, Weiner HL. Microglia-mediated nitric oxide cytotoxicity of T cells following amyloid beta-peptide presentation to Th1 cells. *J Immunol* 2003; **171**: 2216-2224 [PMID: 12928365 DOI: 10.4049/jimmunol.171.5.2216]
- 50 **Wood KJ**, Sawitzki B. Interferon gamma: a crucial role in the function of induced regulatory T cells in vivo. *Trends Immunol* 2006; **27**: 183-187 [PMID: 16527542 DOI: 10.1016/j.it.2006.02.008]

P- Reviewer: Chapel A, Cnossen WR **S- Editor:** Wen LL
L- Editor: Wang TQ **E- Editor:** Wang CH



