

World Journal of *Orthopedics*

World J Orthop 2024 November 18; 15(11): 1001-1111



EDITORIAL

- 1001** Adipose-derived stem cells and knee osteoarthritis: New perspectives, old concerns
de Sousa EB, Gabbi Filho JPA, Gameiro VS, Baptista LS
- 1007** Cardiotoxicity concerns in total joint arthroplasty
Cheng CH, Hao WR, Cheng TH

ORIGINAL ARTICLE**Retrospective Study**

- 1015** Effectiveness of negative pressure wound therapy in complex surgical treatment of necrotizing fasciitis of the upper limb
Lipatov KV, Asatryan A, Melkonyan G, Kazantsev AD, Solov'eva EI, Krivikhin DV, Gorbacheva IV, Cherkasov UE

Observational Study

- 1023** Artificial intelligence awareness and perceptions among pediatric orthopedic surgeons: A cross-sectional observational study
Alomran AK, Alomar MF, Akhdher AA, Al Qanber AR, Albik AK, Alumran A, Abdulwahab AH
- 1036** Predictive factors for coronal and sagittal graft extrusion length after using tendon autograft for medial meniscus reconstruction
Zhu TW, Xiang XX, Li CH, Li RX, Zhang N
- 1047** Presentation and management outcome of foot drop with tibialis posterior tendon transfer
Saaig M

Basic Study

- 1056** Impacts of mesenchymal stem cells and hyaluronic acid on inflammatory indicators and antioxidant defense in experimental ankle osteoarthritis
Hagag UI, Halfaya FM, Al-Muzafar HM, Al-Jameel SS, Amin KA, Abou El-Kheir W, Mahdi EA, Hassan GANR, Ahmed OM
- 1075** Gene expression analysis of cytokines and MMPs in melatonin and rhBMP-2 enhanced bone remodeling
Paulini MR, Montarele LF, Pitol DL, Giannocco G, Pereira BF, Buchaim DV, Reis CHB, Buchaim RL, Mardegan Issa JP

CASE REPORT

- 1088** Ilizarov technique for treatment of a giant aneurysmal bone cyst at the distal femur: A case report
Long XY, Sun F, Wang T, Li P, Tian Z, Wu XW
- 1095** Metagenomic next-generation sequencing may assist diagnosis of osteomyelitis caused by *Mycobacterium houstonense*: A case report
Lin HY, Tan QH

- 1101** Arthroscopic synovectomy for synovial hyperplasia in chronic knee gouty arthritis: A case report
Utoyo GA, Calvin C

LETTER TO THE EDITOR

- 1109** Insights and implications from the study on meniscus reconstruction using tendon autograft
Nguyen PD, Lam TK

ABOUT COVER

Peer Reviewer of *World Journal of Orthopedics*, Jie Wen, PhD, Assistant Professor, Associate Chief Physician, Department of Pediatric Orthopedics, Hunan Provincial People's Hospital, The First Affiliated Hospital of Hunan Normal University, Changsha 410013, Hunan Province, China. cashwj@qq.com

AIMS AND SCOPE

The primary aim of *World Journal of Orthopedics* (WJO, *World J Orthop*) is to provide scholars and readers from various fields of orthopedics with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJO mainly publishes articles reporting research results and findings obtained in the field of orthopedics and covering a wide range of topics including arthroscopy, bone trauma, bone tumors, hand and foot surgery, joint surgery, orthopedic trauma, osteoarthropathy, osteoporosis, pediatric orthopedics, spinal diseases, spine surgery, and sports medicine.

INDEXING/ABSTRACTING

WJO is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for WJO as 2.0; JIF Quartile: Q2. The WJO's CiteScore for 2023 is 3.1.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Yu-Qing Zhao*; Production Department Director: *Xiao-Mei Zheng*; Cover Editor: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Orthopedics

ISSN

ISSN 2218-5836 (online)

LAUNCH DATE

November 18, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Massimiliano Leigheb, Xiao-Jian Ye

EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF

Xin Gu

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/2218-5836/editorialboard.htm>

PUBLICATION DATE

November 18, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

PUBLISHING PARTNER

The Minimally Invasive Spine Surgery Research Center Of Shanghai Jiaotong University

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/bpg/GerInfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

PUBLISHING PARTNER's OFFICIAL WEBSITE

https://www.shtrhospital.com/zkjs/info_29.aspx?itemid=647



Retrospective Study

Effectiveness of negative pressure wound therapy in complex surgical treatment of necrotizing fasciitis of the upper limb

Konstantin V Lipatov, Arthur Asatryan, George Melkonyan, Aleksandr D Kazantcev, Ekaterina I Solov'eva, Denis V Krivikhin, Irina V Gorbacheva, Urii E Cherkasov

Specialty type: Orthopedics

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade C

Novelty: Grade B

Creativity or Innovation: Grade B

Scientific Significance: Grade B

P-Reviewer: Das A

Received: June 11, 2024

Revised: September 22, 2024

Accepted: October 10, 2024

Published online: November 18, 2024

Processing time: 156 Days and 12.7 Hours



Konstantin V Lipatov, Aleksandr D Kazantcev, Department of General Surgery, Institute of Clinical Medicine named after N.V. Sklifosovsky, Sechenov First Moscow State Medical University (Sechenov University), Moscow 119021, Russia

Arthur Asatryan, Denis V Krivikhin, Department of General Surgery, Wound and Wound Infection Surgery, State Budgetary Institution "City Clinical Hospital named after S.S. Yudin of Moscow Healthcare Department", Moscow 115446, Russia

George Melkonyan, Department of General Surgery, Physician of The Hospital for War Veterans No 3, Moscow 129336, Russia

Ekaterina I Solov'eva, Irina V Gorbacheva, Urii E Cherkasov, Department of General Surgery, I.M. Sechenov First Moscow State Medical University, Moscow 119048, Russia

Corresponding author: Konstantin V Lipatov, DSc, MD, Full Professor, Surgeon, Department of General Surgery, Institute of Clinical Medicine named after N.V. Sklifosovsky, Sechenov First Moscow State Medical University (Sechenov University), Ro-ssolimo Street 11-2, Moscow 119021, Russia. lipatov_k_v@staff.sechenov.ru

Abstract

BACKGROUND

Necrotizing fasciitis (NF) of the upper extremities is a severe surgical pathology, and the incidence of this disease has been steadily increasing in recent decades. Surgical treatment is accompanied by the formation of extensive wounds, which can be treated with significant difficulties. In recent years, negative pressure wound therapy (NPWT) has proven to be highly effective. It is also promising for the treatment of NF.

AIM

To explore the effectiveness of NPWT in the treatment of NF of the upper extremities.

METHODS

The results of the treatment of 36 patients with NF of the upper extremities in two groups (NPWT group and control group; 2022–2023) were retrospectively analyzed. In the NPWT group, the NPWT method (120 mmHg; constant mode) was used after surgical treatment. The number of vacuum-assisted dressings in pa-

tients ranged from 1 to 3, depending on the dynamics of the wound process. The duration of fixation of one bandage was up to 2–3 d. In the control group, conventional methods of local wound treatment were used. The following indicators were analyzed: The treatment delay, the prevalence of inflammation, the microbial landscape, the number of debridements, the duration of wound preparation for surgical closure, and the nature of skin plastic surgery.

RESULTS

Most patients experienced a significant treatment delay [4 d, interquartile range (IQR): 2–7 d], which led to the spread of the pathological process to the forearm and shoulder. The most common pathogens were *Staphylococcus aureus* (14; 38.9%) and *Streptococcus pyogenes* (22; 61.1%). The average number of debridements *per patient* was 5 (IQR: 3–7), with no difference between groups. The average time to prepare wounds for surgical closure was 11 ± 4 d in the NPWT group and 29 ± 10 d ($P = 0.00001$) in the control group. In the NPWT group, the wounds were more often closed with local tissues (15; 83.3%), and in the control group, split-thickness skin grafts were more often used (4; 50%).

CONCLUSION

The predominant isolation of *Staphylococcus aureus* and/or *Streptococcus pyogenes* from the lesions allowed us to classify these patients as NF type II. Multiple debridement procedures have become a feature of this disease treatment. The use of NPWT has significantly reduced the time required to prepare wounds for surgical closure. Early closure of wounds allows for more frequent use of local tissue repair, which ensures better results. NPWT is a highly effective way to prepare wounds for early surgical closure in patients with upper extremity NF.

Key Words: Necrotizing fasciitis; Upper limb; Negative pressure wound therapy; Vacuum-assisted closure; Surgical treatment

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Negative pressure wound therapy, performed after full surgical debridement for necrotizing fasciitis of the upper extremities, significantly reduces the duration of postnecrectomy wound preparation for surgical closure, thereby improving patient treatment outcomes.

Citation: Lipatov KV, Asatryan A, Melkonyan G, Kazantsev AD, Solov'eva EI, Krivikhin DV, Gorbacheva IV, Cherkasov UE. Effectiveness of negative pressure wound therapy in complex surgical treatment of necrotizing fasciitis of the upper limb. *World J Orthop* 2024; 15(11): 1015-1022

URL: <https://www.wjgnet.com/2218-5836/full/v15/i11/1015.htm>

DOI: <https://dx.doi.org/10.5312/wjo.v15.i11.1015>

INTRODUCTION

Necrotizing fasciitis (NF) was considered a fairly rare disease, and publications devoted to its diagnosis and treatment have rarely included more than 10 clinical cases. In recent years, the situation has changed significantly. The frequency of this pathology has increased, and the severity of its course, high mortality rate, and significant material costs make the search for effective treatment methods urgent[1,2].

Among all NF involvements, those in the upper limb account for 6%–27%, and the average in-hospital mortality/30-d mortality may range between 20% and 40%[3]. According to the classification of Choueka and De Tolla (2020), NF type II is the most common when it affects the extremities[4]. The causative agents of infection are *Staphylococcus aureus* and *Streptococcus pyogenes*, both as monoinfections and in combination with each other[5]. The disease usually develops after minor skin injuries or even after blunt trauma to soft tissues[6]. NF type II is more common in healthy people of working age, but cases of its occurrence are not uncommon in patients suffering from diabetes mellitus, drug addiction, or who is taking corticosteroid hormones[7]. Certain difficulties in diagnosing the disease at the initial stage can lead to untimely surgical intervention and extension of inflammation to surrounding tissues[8,9]. In these cases, as a result of delayed surgical treatment, extensive wound defects are formed, the further treatment of which can present significant difficulties [10]. This disease is characterized by the formation of secondary necrosis, which determines the need for repeated debridement, the number of which can reach 5 or more[11]. In addition, the wound process in NF patients is characterized by a complicated course with a prolonged phase of inflammation. All these lead to the need for long-term hospital treatment compared to other purulent diseases of soft tissues. The scarring process that develops over time in the tissues surrounding the wound leads to a decrease in plastic reserves. This, in turn, significantly reduces the possibility of closing wounds with local tissues and expands the indications for split-skin grafts[3,6,7]. However, the cosmetic and functional results are much worse, which is especially important for the upper limb. In this regard, it is difficult to overestimate the importance of methods that help reduce the time needed to prepare wounds for surgical closure[12].

Negative pressure wound therapy (NPWT) has proven to be highly effective and is widely used for treating purulent wounds of various origins and locations. NPWT (syn. vacuum-assisted closure-VAC) offers many benefits: Wound isolation, reduced edema, increased tissue perfusion, and reduced bacterial load[13,14]. All of these factors contribute to subsequent reconstructive surgery. Using VAC reduces the absorption of toxins and mitigates pain. Less frequent dressings not only have a positive effect on the patient's psycho-emotional state, but also have a significant economic effect, reducing the amount of material used in dressings and the labor costs of medical personnel. This is especially important for extensive wounds accompanying NF[15]. The effective use of NPWT requires strict adherence to a number of requirements and restrictions. Preliminary removal of the main mass of necrotic and nonviable tissue is considered a mandatory condition for the utilization of NPWT. It is also necessary to take into account that a vacuum dressing applied immediately after surgical treatment can provoke bleeding[14].

The choice of plastic wound closure method for the treatment of NF of the upper extremity is influenced by a number of factors. The most significant factors are: The size and location of the defect, the plastic reserves of the surrounding soft tissues, and the age and condition of the patient[11,13]. Additionally, the professional skills of surgeons should not be underestimated. When closing wound defects in the area of the upper limb, full-thickness skin grafting (with local tissue or displaced flaps-free or pedicled) is a priority[13]. Unfortunately, with significant skin loss, split-thickness skin grafting often becomes the only possible method[15].

MATERIALS AND METHODS

Methodology

The study was carried out in a single clinical center. The results of the examination and treatment of 36 patients with NF of the upper extremities were retrospectively analyzed.

Type of sampling: Non-repetitive subjects were consecutively recruited.

Inclusion criteria: NF of the upper limb.

Exclusion criteria: Extension of inflammation to the trunk, diabetes mellitus, intake of corticosteroid hormones, pregnancy, and poor compliance with the treatment.

Population

All patients were divided into two groups depending on the characteristics of the local treatment of the wounds after surgical treatment: NPWT group and control group. In the NPWT group (18 patients), NPWT (120 mmHg; constant mode) was used. The number of vacuum-assisted dressings used in patients ranged from 1 to 3, depending on the dynamics of the wound process. The duration of fixation of one bandage was up to 2–3 d. In the control group (18 patients), conventional methods of local wound treatment (antiseptic solutions) were used. According to the main parameters (age, gender, localization, and prevalence of the pathological process), patients of both groups corresponded to each other. All patients underwent urgent debridement supplemented by repeated treatments at intervals of 24 h until the wounds were cleared of nonviable tissue.

The following indicators were analyzed and compared in the groups: The treatment delay, the prevalence of inflammation, the microbial landscape, the number of debridements, the duration of wound preparation for surgical closure, and the nature of skin plastic surgery.

The choice of method for surgical closure of wounds was determined by the size, location, and plastic reserves of the surrounding skin. To assess the severity of the general inflammatory reaction of the body, the level of leukocytes in the peripheral blood and the concentrations of C-reactive protein and procalcitonin in the plasma were studied.

Compliance with ethical standards and consent to participate

The local institutional review board approved the use of patient data for research purposes prior to reviewing the data, and this study was deemed exempt from continued review.

Informed consent was not required due to the retrospective nature of this study. All study participants or their legal guardian provided informed written consent about personal and medical data collection prior to study enrolment. All procedures were performed in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2008.

Statistical analysis

Statistical processing was carried out by collecting and grouping factual material, with the median and interquartile range (IQR) calculated. The arithmetic mean and standard deviation (SD) were calculated to estimate the average duration of wound preparation for surgical closure. The normality of the distribution was determined using the Shapiro-Wilk normality test. The significance of differences in nonparametric indicators was determined using the Mann-Whitney *U* test. Differences in indicators at $P < 0.05$ were considered significant. The study was reviewed by our biostatistical expert.

RESULTS

Descriptive statistics

The age of the patients did not significantly differ between the two groups. The median age was 50 years (IQR: 39–67 years). Men predominated. The spread of the pathological process to the elbow joint was observed in 11 (61.1%) patients in the NPWT group and in 10 (55.6%) in the control group. The majority of patients in both groups were characterized by delays in seeking medical help. The associated treatment delay was not significantly different between groups ($P = 0.6$) and was 4 d (IQR: 3–7 d).

The average number of debridements *per patient* was 5 (IQR: 3–7), and there was no significant difference between the groups ($P = 0.94$). The average time to prepare wounds for surgical closure in the NPWT group was 11 ± 4 d, which was significantly less than that of the control group (29 ± 10 d) ($P = 0.00001$).

Laboratory investigations

During microbiological examination, in the vast majority of cases, *Staphylococcus aureus* (14; 38.9%) and *Streptococcus pyogenes* (22; 61.1%) were isolated, both in the form of a monoculture (24; 66.7%) and in association with each other (6; 16.7%). In 1 (2.7%) patient, the causative agent of NF was *Pasteurella multocida*. In 4 (11.1%) patients, it was not possible to identify the pathogen (Figure 1). Initial empirical antibiotic therapy did not differ between the two groups and included intravenous administration of broad-spectrum drugs, mainly penicillins and cephalosporins. Based on the results of a microbiological study of wound exudate, antibacterial therapy was adjusted considering the sensitivity of the bacterial flora. The overwhelming majority of patients with NF of the upper limb had pronounced leukocytosis [NPWT group: $21 \times 10^9/L$ (IQR: $18\text{--}27 \times 10^9/L$); control group: $22 \times 10^9/L$ (IQR: $17\text{--}27 \times 10^9$)] and an increase in the concentration of C-reactive protein [NPWT group: 103 mg/L (IQR: 49–231 mg/L); control group: 222 mg/L (IQR: 180–334)]. Procalcitonin levels were moderately increased in the two groups [NPWT group: 0.69 ng/mL (IQR: 0.5–1.1 ng/mL); control group: 3 ng/mL (IQR: 1–12 ng/mL)].

Plastic surgery

Surgical wound closure was performed in 15 (83.3%) patients in the NPWT group and in 8 (44.4%) in the control group. In the NPWT group, wounds were closed with local tissue in 12 (80%) patients and in 3 (20%) patients when a split-thickness skin graft was used. In the control group, full-thickness skin was used in 4 (50%) patients (Figure 2). In 2 patients, plastic surgery was performed with local tissues, and in 2 patients, plastic surgery was performed with a free vascularized fasciocutaneous flap from the groin area. Split-thickness grafting was performed in 4 (50%) patients (Supplementary Table 1).

Mortality

In-hospital mortality rate of the patients with NF of the upper extremities was 11.1% (4 patients), and NF was not associated with the method of wound treatment.

DISCUSSION

A significant delay in treatment in most patients led to the spread of the pathological process. Moreover, in more than one third of patients, inflammation involved the entire upper limb. The severity of the disease and systemic inflammatory response syndrome was indicated by a sharp increase in the levels of leukocytes in the blood and C-reactive protein in the plasma.

The vast majority of microflora species identified were associated with *Staphylococcus aureus* and *Streptococcus pyogenes*, which allowed these patients to be classified as NF type II[4]. The typical approach to surgical treatment of this disease has become multiple rounds of debridement, which creates conditions not only for eliminating the infection but also for the effective use of the NPWT method. The use of vacuum-assisted dressings has made it possible to significantly reduce the time required to prepare wounds for surgical closure[3]. As a result, there was a change in treatment tactics. In the NPWT group, one-stage inpatient treatment was used, which made it possible to eliminate the infection and close the wounds. After this, the patient was sent for rehabilitation. In the control group, a three-stage treatment option had to be used. At the first inpatient stage, the infection was eliminated. In the second stage, outpatient, local treatment of the wound was carried out until it was ready for surgical closure. In the third stage, stationary, skin grafting was performed (Figure 3). In addition to the difference in the timing of preparing wounds for plastic surgery, the difference in the range of surgical interventions also attracts attention. Early wound closure in the NPWT group allowed for more frequent use of local tissue, which provided better functional and cosmetic results. Delayed wound closure in the control group occurred against the background of a decrease in the plastic reserves of the surrounding skin and the scarring process[7]. The result has been an increase in the number of split skin grafts with poorer results (Figure 4, Figure 5 and Figure 6). The use of NPWT had no effect on the level of hospital mortality, which complies with data from other researchers[14].




Figure 1 Microorganisms cultured from patients with necrotizing fasciitis of the upper limb. MSSA: Methicillin-sensitive staphylococcus aureus.




Figure 2 Plastic surgery of postnecrectomy wounds. NPWT: Negative pressure wound therapy.




Figure 3 Treatment strategy for patients with necrotizing fasciitis of the upper limb. VAC: Vacuum-assisted closure.




Figure 4 The first case. A: Clinical presentation at admission; B: After first debridement; C: After negative pressure wound therapy; D: Long-term result of plastic surgery.

CONCLUSION

NF of the upper extremities is a serious disease, for which timely and complete debridement is important. NPWT is an effective method for the local treatment of postnecrectomy wounds and can significantly reduce the duration of their preparation for surgical closure. Reaching a better cosmetic result is extremely important for the aesthetic appearance of the upper limb.



Figure 5 The second case. A: Clinical presentation at admission; B: After debridement; C: After negative pressure wound therapy; D: After plastic surgery; E and F: Long-term outcomes after 6 mo.




Figure 6 The third case. A: Clinical presentation at admission; B: After first debridement; C: After third debridement; D: Negative pressure wound therapy; E: After plastic surgery; F: Result.

FOOTNOTES

Author contributions: Lipatov KV performed conceptualization, and manuscript writing, review, and editing; Asatryan A and

Melkonyan G performed methodology and writing of the original draft; Kazantsev AD performed visualization, and manuscript writing, review, and editing; Solov'eva EI and Gorbacheva IV performed investigation and writing of the original draft; Krivikhin DV and Cherkasov UE performed formal analysis and supervision.

Institutional review board statement: The study was reviewed and approved for publication by our Institutional Reviewer.

Informed consent statement: All study participants or their legal guardian provided informed written consent about personal and medical data collection prior to study enrolment.

Conflict-of-interest statement: All the authors have no conflict of interest related to the manuscript.

Data sharing statement: The original anonymous dataset is available on request from the corresponding author at lipatov_k_v@staff.sechenov.ru.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: Russia

ORCID number: Konstantin V Lipatov 0000-0002-9902-2650; Arthur Asatryan 0000-0002-8409-2605; George Melkonyan 0000-0001-7234-4185; Aleksandr D Kazantsev 0000-0003-1238-1990; Ekaterina I Solov'eva 0000-0003-4143-3593; Denis V Krivikhin 0009-0000-6899-4804; Irina V Gorbacheva 0000-0002-1060-1163; Urii E Cherkasov 0000-0003-0137-4517.

S-Editor: Liu H

L-Editor: Wang TQ

P-Editor: Zhao YQ

REFERENCES

- Chen LL, Fasolka B, Treacy C. Necrotizing fasciitis: A comprehensive review. *Nursing* 2020; **50**: 34-40 [PMID: 32826674 DOI: 10.1097/01.NURSE.0000694752.85118.62]
- Nawijn F, Verhiel SHWL, Lunn KN, Eberlin KR, Hietbrink F, Chen NC. Factors Associated with Mortality and Amputation Caused by Necrotizing Soft Tissue Infections of the Upper Extremity: A Retrospective Cohort Study. *World J Surg* 2020; **44**: 730-740 [PMID: 31664494 DOI: 10.1007/s00268-019-05256-9]
- Hakkarainen TW, Kopari NM, Pham TN, Evans HL. Necrotizing soft tissue infections: review and current concepts in treatment, systems of care, and outcomes. *Curr Probl Surg* 2014; **51**: 344-362 [PMID: 25069713 DOI: 10.1067/j.cpsurg.2014.06.001]
- Choueka J, De Tolla JE. Necrotizing Infections of the Hand and Wrist: Diagnosis and Treatment Options. *J Am Acad Orthop Surg* 2020; **28**: e55-e63 [PMID: 31403486 DOI: 10.5435/JAAOS-D-17-00716]
- Stevens DL, Bryant AE. Necrotizing Soft-Tissue Infections. *N Engl J Med* 2017; **377**: 2253-2265 [PMID: 29211672 DOI: 10.1056/NEJMrat1600673]
- Tessier JM, Sanders J, Sartelli M, Ulrych J, De Simone B, Grabowski J, Buckman S, Duane TM. Necrotizing Soft Tissue Infections: A Focused Review of Pathophysiology, Diagnosis, Operative Management, Antimicrobial Therapy, and Pediatrics. *Surg Infect (Larchmt)* 2020; **21**: 81-93 [PMID: 31584343 DOI: 10.1089/sur.2019.219]
- Melillo A, Addagatla K, Jarrett NJ. Necrotizing Soft Tissue Infections of the Upper Extremity. *Hand Clin* 2020; **36**: 339-344 [PMID: 32586460 DOI: 10.1016/j.hcl.2020.03.007]
- El-Menyar A, Asim M, Mudali IN, Mekkodathil A, Latifi R, Al-Thani H. The laboratory risk indicator for necrotizing fasciitis (LRINEC) scoring: the diagnostic and potential prognostic role. *Scand J Trauma Resusc Emerg Med* 2017; **25**: 28 [PMID: 28270166 DOI: 10.1186/s13049-017-0359-z]
- Ditsios K, Chitas K, Christidis P, Charatsis K, Katsimentzas T, Papadopoulos P. Necrotizing Fasciitis of the Upper Extremity - A Review. *Orthop Rev (Pavia)* 2022; **14**: 35320 [PMID: 36034724 DOI: 10.52965/001c.35320]
- Gawaziuk JP, Liu T, Sigurdson L, Buchel E, Hayakawa TEJ, Shiga S, Logsetty S. Free tissue transfer for necrotizing fasciitis reconstruction: A case series. *Burns* 2017; **43**: 1561-1566 [PMID: 28536044 DOI: 10.1016/j.burns.2017.04.007]
- Stevens DL, Bisno AL, Chambers HF, Dellinger EP, Goldstein EJ, Gorbach SL, Hirschmann JV, Kaplan SL, Montoya JG, Wade JC; Infectious Diseases Society of America. Practice guidelines for the diagnosis and management of skin and soft tissue infections: 2014 update by the Infectious Diseases Society of America. *Clin Infect Dis* 2014; **59**: e10-e52 [PMID: 24973422 DOI: 10.1093/cid/ciu444]
- Corona PS, Erimeiku F, Reverté-Vinaixa MM, Soldado F, Amat C, Carrera L. Necrotising fasciitis of the extremities: implementation of new management technologies. *Injury* 2016; **47** Suppl 3: S66-S71 [PMID: 27692110 DOI: 10.1016/S0020-1383(16)30609-X]
- Lee JY, Jung H, Kwon H, Jung SN. Extended negative pressure wound therapy-assisted dermatotraction for the closure of large open fasciotomy wounds in necrotizing fasciitis patients. *World J Emerg Surg* 2014; **9**: 29 [PMID: 24731449 DOI: 10.1186/1749-7922-9-29]
- Zhang R, Zhang Y, Hou L, Yan C. Vacuum-assisted closure versus conventional dressing in necrotizing fasciitis: a systematic review and meta-analysis. *J Orthop Surg Res* 2023; **18**: 85 [PMID: 36737764 DOI: 10.1186/s13018-023-03561-7]
- Yin Y, Zhang R, Li S, Guo J, Hou Z, Zhang Y. Negative-pressure therapy versus conventional therapy on split-thickness skin graft: A systematic review and meta-analysis. *Int J Surg* 2018; **50**: 43-48 [PMID: 29292216 DOI: 10.1016/j.ijssu.2017.12.020]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

