

World Journal of *Clinical Cases*

World J Clin Cases 2024 October 6; 12(28): 6132-6249



EDITORIAL

- 6132** Approach to cardiac masses: Thinking inside and outside the box
De Maria E
- 6137** Stroke: Evolution of newer treatment modalities for acute ischemic stroke
Nag DS, Swain A, Sahu S, Sen B, Vatsala, Parween S
- 6148** Risk of appendiceal neoplasm in patients with appendix disorders
Ferrara F, Peltrini R
- 6151** Multi-systemic melioidosis
Nagoba BS, Dhotre SV, Rayate AS, Mumbre SS, Dhotre PS
- 6155** Pituitary metastasis from lung adenocarcinoma
Wang XJ
- 6159** Hidden army within: Harnessing the microbiome to improve cancer treatment outcomes
Messaritakis I, Vougiouklakis G, Koulouridi A, Agouridis AP, Spernovasilis N

ORIGINAL ARTICLE**Case Control Study**

- 6165** Effects of traditional Chinese medicine on symptoms of patients with spleen and stomach deficiency-related tinnitus
Cui N, Zhao C, Xue JL, Zhu XW
- 6173** Efficacy of graphene nanocomposites for air disinfection in dental clinics: A randomized controlled study
Ju YQ, Yu XH, Wu J, Hu YH, Han XY, Fang D

Retrospective Study

- 6180** Self-expanding metal stent for relieving the stricture after endoscopic injection for esophageal varices
Zhang FL, Xu J, Jiang YH, Zhu YD, Shi Y, Li X, Wang H, Huang CJ, Zhou CH, Zhu Q, Chen JW
- 6187** Incidental renal cell carcinoma post bilateral nephrectomy in autosomal dominant polycystic kidney disease
Shin MH, Choi NK

Clinical Trials Study

- 6195** Effect of intraoperative injection of esketamine on postoperative analgesia and postoperative rehabilitation after cesarean section
Chen HZ, Gao Y, Li KK, An L, Yan J, Li H, Zhang J

Prospective Study

- 6204** Virtual reality for preoperative patient education: Impact on satisfaction, usability, and burnout from the perspective of new nurses
Kim J, Kim D, Oh SH, Kwon H

CASE REPORT

- 6217** Uveitis glaucoma hyphema syndrome as a result of glaucoma implant: A case report
Altwijri RJ, Alsirhy E
- 6222** Recurrent multisystem Langerhans cell histiocytosis involving the female genitalia: A case report
Yuan CY, Zhang ZR, Guo MF, Zhang N
- 6230** Toripalimab in combination with chemotherapy effectively suppresses local recurrence and metastatic sarcomatoid renal cell carcinoma: A case report
Gao MZ, Wang NF, Wang JY, Ma L, Yang YC

LETTER TO THE EDITOR

- 6237** Re-evaluating the necessity of routine laboratory monitoring during isotretinoin therapy for acne
Zhu YX, Wu L, Zhu SC, Wang XP, Zhang D, Tan YP, Ouyang XL, Li CM
- 6241** Conversion therapy for hepatocellular carcinoma to improve treatment strategies for intermediate and advanced stages
Giorgio A, De Luca M
- 6244** Old, the new, more or less and the conundrum for biomarkers in cardiovascular diseases?
Iyngkaran P
- 6247** Early pirfenidone treatment enhances lung function in idiopathic pulmonary fibrosis patients
Zhang JW

ABOUT COVER

Peer Reviewer of *World Journal of Clinical Cases*, Saptarshi Bhattacharya, FACE, MD, Doctor, Endocrinology, Indraprastha Apollo Hospitals, Sarita Vihar, New Delhi 110076, India. saptarshi515@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in PubMed, PubMed Central, *Reference Citation Analysis*, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for WJCC as 1.0; JIF without journal self cites: 0.9; 5-year JIF: 1.1; JIF Rank: 168/325 in medicine, general and internal; JIF Quartile: Q3; and 5-year JIF Quartile: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yu; Production Department Director: Xu Guo; Cover Editor: Jin-Li Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Bao-Gan Peng, Salim Surani

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

October 6, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Prospective Study

Virtual reality for preoperative patient education: Impact on satisfaction, usability, and burnout from the perspective of new nurses

Jiyoung Kim, Donghyun Kim, Sang-Ha Oh, Hyeokjae Kwon

Specialty type: Nursing

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade B

Novelty: Grade A

Creativity or Innovation: Grade A

Scientific Significance: Grade B

P-Reviewer: Varama A

Received: June 7, 2024

Revised: July 27, 2024

Accepted: July 31, 2024

Published online: October 6, 2024

Processing time: 66 Days and 18 Hours



Jiyoung Kim, College of Nursing, Woosuk University, Wanju-gun 55338, South Korea

Donghyun Kim, Department of Medical Science, Chungnam National University College of Medicine, Daejeon 35015, South Korea

Donghyun Kim, Sang-Ha Oh, Hyeokjae Kwon, Department of Plastic and Reconstructive Surgery, Chungnam National University Hospital, Daejeon 35015, South Korea

Sang-Ha Oh, Department of Plastic and Reconstructive Surgery, Chungnam National University College of Medicine, Daejeon 35015, South Korea

Co-corresponding authors: Sang-Ha Oh and Hyeokjae Kwon.

Corresponding author: Hyeokjae Kwon, MD, PhD, Surgeon, Department of Plastic and Reconstructive Surgery, Chungnam National University Hospital, 282 Munhwa-ro Jung-gu Daejeon, Daejeon 35015, South Korea. kwon.hyeokjae@cnuh.co.kr

Abstract

BACKGROUND

Traditional paper-based preoperative patient education is a struggle for new nurses and requires extensive training. In this situation, virtual reality technology can help the new nurses. Despite its potential benefits, there are studies on patient satisfaction but there is limited information on the usability of virtual reality (VR) technology for new nurses in giving preoperative education to patients.

AIM

To investigate the impact on satisfaction, usability, and burnout of a system using VR technology in preoperative patient education.

METHODS

The study involved 20 nurses from the plastic surgery ward and 80 patients admitted between April and May 2019. Each nurse taught four patients: Two using traditional verbal education and two using virtual reality. The System Usability Scale, After-Scenario Questionnaire, and Maslach Burnout Inventory (MBI) were employed to evaluate the impact of these education methods.

RESULTS

The VR education groups showed a statistically higher satisfaction than the traditional verbal education groups. Among the three subscales of the MBI, emotional exhaustion and personal accomplishment improved statistically significantly. VR was also better in terms of usability.

CONCLUSION

This study suggests VR enhances usability and reduces burnout in nurses, but further research is needed to assess its impact on depersonalization and objective measures like stress and heart rate.

Key Words: Virtual reality; Preoperative care; Patient education; Burnout; Usability; Satisfaction; Prospective studies

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Virtual reality technology proved to be significantly more satisfactory and usable than traditional paper based methods for healthcare providers (especially for nurses) working in plastic surgeries wards.

Citation: Kim J, Kim D, Oh SH, Kwon H. Virtual reality for preoperative patient education: Impact on satisfaction, usability, and burnout from the perspective of new nurses. *World J Clin Cases* 2024; 12(28): 6204-6216

URL: <https://www.wjgnet.com/2307-8960/full/v12/i28/6204.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v12.i28.6204>

INTRODUCTION

In recent years, virtual reality (VR) technology has evolved rapidly, opening new possibilities for many industries, including healthcare[1]. With the aid of VR technology, users can engage in immersive and realistic experiences in computer-generated environments and simulations. It has been applied to a variety of patient care, including psychotherapy[2], medical education[3], pain management[4], and rehabilitation[5]. VR can provide patients with a simpler approach for understanding complicated medical procedures in preoperative situations[6,7]. More dynamically and interestingly, it enables patients to experience the procedure and understand the possible risks and complications[8].

In addition, traditional paper-based preoperative patient education is stressful for new nurses, requires extensive training[9], and the patients receive varying levels of instruction as a result of nurses' various understandings and practices[10].

Receiving varying degrees of care from various nurses is a concern in terms of quality of nursing. Furthermore, the learning curve that new nurses must go through to overcome this can deplete their energy and produce excessive stress, leading to work dissatisfaction and maladjustment, poor mental and physical impacts, and burnout[11].

Burnout is a phenomenon characterized by emotional exhaustion, depersonalization, and a sense of poor accomplishment in life as a result of job-related stress, which can result in a worsening in the quality of care provided[12]. It is caused by prolonged exposure to job stresses such as high patient caseloads, long work hours, and challenging interactions with patients[13]. Nurse burnout is a serious problem that must be addressed for a variety of reasons. It endangers the emotional and physical health of nurses, causing symptoms such as depression, anxiety, sleep disturbances and an increased risk of chronic diseases[14-16]. Furthermore, it can impair patient care quality by decreasing nurses' motivation, attention, and ability to offer compassionate care[17]. This burnout ultimately puts patients at risk and reduces the efficiency of the healthcare system.

According to one study, 43% of nurses who reported high levels of burnout showed plans to quit compared to just 11% of non-burned-out nurses, indicating that managing burnout can help keep experienced nurses in the field for a longer period[18]. Another study showed that healthcare employees had a 32% greater risk of suicide than non-healthcare workers. Medical support staff had the highest annual suicide rate per 100000 among healthcare workers, followed by nurses at 16.0[19].

In this situation, virtual reality technology can help both the patient and the new nurse. Despite its potential benefits, there are studies on patient satisfaction[6,20,21] but there are limited studies on the satisfaction and usability of VR technology for preoperative education of new nurses and its impact on burnout of new nurses. We hypothesized that using VR technology would result in higher satisfaction and usability for new nurses compared to traditional paper-based methods, and that it would reduce burnout and promote wellness. Thus, this study aims to compare the satisfaction and usability, impact on burnout of new nurses of a system using VR technology with traditional paper-based methods in preoperative patient education from the perspective of new nurses.

MATERIALS AND METHODS

This study was approved by the Research Ethics Board of Chungnam National University Hospital (approval no. CNUH 2019-06-008-002) and informed consent was obtained from all participants. From April to December 2019, this study involved 20 nurses from the plastic surgery ward of Chungnam National University Hospital in Daejeon, South Korea. All nurses who participated in the study had been working at the advanced beginner level with less than two years of experience[22] (Table 1).

VR contents

The Department of Anesthesiology and Pain Medicine and the Department of Plastic Surgery collaborated to create a scenario of educational content using VR before and after surgery. Subsequently, VR contents for the full hospitalization process, pre-surgery education, procedures on the day of surgery, and discharge were produced based on the completed scenario. In particular, general education the day before and at the time for surgery, shifting from the ward bed to the stretcher car, arrival in front of the operating room *via* the elevator, wait and entry into the waiting room, entry into the operating room, laying on the operating bed for general anesthesia, shifting of the patient to the recovery room and hospital room after surgery, and discharge from the hospital after recovery was implemented using VR (Figure 1). An Insta360 ONE X (Insta 360, Seoul, South Korea) was used to record the video. Multiple persons can watch the presentation at once using a PICO G2 head mount device (PICO Interactive Korea Co., Ltd., Seoul, South Korea). The full video can be viewed on YouTube at <https://www.youtube.com/watch?v=thq3GikfPL4&t=11s>.

Patients and Survey tools

Patients who were admitted to our plastic surgery department between April 21, 2019, and May 31, 2019, and underwent their first operation under general anesthesia served as the study participants. Patients who could not benefit from VR education because of cognitive disability, visual impairment, claustrophobia, or those who refused to participate in the questionnaire were excluded. After excluding unsuitable patients, we used randomization software (Random Allocation Software, version 1.0; Isfahan Medical Sciences) to randomly assign the 40 patients in the control group to 20 ward nurses in groups of two (Figure 2).

Each patient was hospitalized the day before surgery, and the nurses who participated in the study ($n = 20$) provided traditional paper-based education to the two control group patients assigned to each participating nurse and completed satisfaction, usability, burnout surveys the following day. After the preoperative patient education of the control group and the satisfaction, usability, burnout surveys of the nurses who participated in the study, the preoperative education using VR and satisfaction, usability, burnout surveys were conducted in the same manner as for the control group ($n = 40$) (Figure 3). Patient characteristics are listed in Table 2. Age was tested for homogeneity between the two groups using an independent t-test, sex was tested using a χ^2 test, and the type of surgery was tested using Fisher's exact test. After patient education, and satisfaction, usability, burnout surveys were conducted.

To investigate satisfaction, the seven items in the questionnaire were scored on a 5-point Likert scale. Items 3 and 4 were designed as reverse questions, and considered in the following statistical analyses (Table 3).

To investigate usability of virtual reality system, we used two survey scales; The System Usability Scale and After-Scenario Questionnaire. The System Usability Scale (SUS) was developed in 1986 by John Brooke as a "Quick and Dirty" tool for measuring usability[23]. It consists of 10 items, each of which can be answered on a five-point scale from strongly agree to strongly disagree, and is a reliable tool used to evaluate the usability of various products and services (Table 4).

Due to its effectiveness and ease of use—just three questions—the After-Scenario Questionnaire (ASQ)[24], a usability survey created in 1995 by J.R. Lewis, is widely in use today (Table 5).

To measure burnout, we used the Maslach Burnout Inventory. The Maslach Burnout Inventory (MBI) is a widely used and rigorously validated psychometric instrument developed by Maslach *et al*[25] to assess and quantify burnout among healthcare professionals, including nurses. Comprising three distinct subscales—emotional exhaustion, depersonalization, and personal accomplishment—the MBI provides a comprehensive assessment of the multidimensional nature of burnout (Table 6).

Statistical analysis

A paired *t*-test was used to statistically compare the satisfaction before and after using VR by converting the Likert scale for each item in the satisfaction survey into a score.

After the survey of the nurses who performed the intervention for the control and experimental groups, the scores for each item of the SUS and the total score were calculated and compared using paired-*t* test. ASQ scores were also statistically compared in the same manner.

MBI scores were also calculated and compared statistically using a paired *t* test. Reverse-coded survey items were handled and statistically analyzed appropriately.

All analyses were conducted using Statistical Package for Social Science software (version 26.0, IBM).

RESULTS

Satisfaction

The average overall score in the satisfaction survey taken after giving the control group traditional verbal education was

Table 1 Demographics of participating nurses

Characteristics of participating nurses			
Sex	Male	0	(0/20)
	Female	20	(20/20)
Mean age, years		24.4	
Mean years of experience		2.3	

Table 2 Demographics of the patients

Characteristics of the patients		Control group (n = 40), n of mean ± SD		VR Group(n = 40), n of mean ± SD		t	P value
Sex	Male	25	(25/40)	27	(27/40)		0.815
	Female	15	(15/40)	13	(13/40)		
Mean age, years		42.75 ± 13.23		40.75 ± 15.60		0.62	0.538
Type of operation	Facial trauma	26	(26/40)	25	(25/40)		0.717
	Limb or trunk reconstruction	3	(3/40)	3	(3/40)		
	Corrective rhinoplasty	2	(2/40)	1	(1/40)		
	Benign tumor excision	7	(7/40)	10	(10/40)		
	Breast augmentation	0	(0/40)	1	(1/40)		
	<i>etc.</i> (Skin cancer, burn)	2	(2/40)	0	(0/40)		

VR: virtual reality.

Table 3 Questionnaire items

	Strongly disagree (1)	Disagree (2)	Neutral (3)	Agree (4)	Strongly disagree (5)
1 Does the amount of time needed for preoperative education make sense?					
2 Is a preoperative education making your job more difficult?					
3 Does a preoperative education make you more physically stressed?					
4 Does a preoperative education make you more mentally stressed?					
5 After explaining the procedure to the patient, do you feel satisfied?					
6 Do you believe that by giving the patient precise information, your preoperative education satisfied them?					
7 Do you employ effective communication skills or tools in your preoperative education?					

20.50 ± 1.76 points. In addition, the experimental group received preoperative patient education through virtual reality, and a satisfaction survey was conducted. The average overall score was 30.60 ± 2.37. With a paired t-test P value of less than 0.05, the experimental group showed a statistically higher satisfaction than the control group (Table 7). The experimental group reported significantly higher satisfaction than the control group across all seven survey items when comparing each item (Table 8). Bar plots with error bars diagrams are shown in Figure 4.

Usability


Based on the results of the SUS survey, the traditional paper-based method showed statistically significant results for the items "System too complex," "Need of post-support help," "Too much inconsistent," "Cumbersome to use," and "Need of pre-learn training." However, all 20 nurses in the study demonstrated statistically significant superior results for the VR-based patient education system for the items "It will be used frequently," "Easy to use," "Well integrated," "Learned very quickly," and "Confident to use." The results are shown in Table 9 and Figure 5. Although the total score was statistically significantly higher for the VR-enabled system compared to the traditional system (68.00 ± 5.42), this is only the average system usability scale found in other studies[26].

Table 4 System usability scale

System usability scale	Strongly disagree (1)	Disagree (2)	Neutral (3)	Agree (4)	Strongly agree (5)
I think that I would like to use this system frequently (SUS1)					
I found the system unnecessarily complex (SUS2)					
I thought the system was easy to use (SUS3)					
I think that I would need the support of a technical person to be able to use this system (SUS4)					
I found the various functions in this system were well integrated (SUS5)					
I thought there was too much inconsistency in this system (SUS6)					
I would imagine that most people would learn to use this system very quickly (SUS7)					
I found the system very cumbersome to use (SUS8)					
I felt very confident using the system (SUS9)					
I needed to learn a lot of things before I could get going with this system (SUS10)					

Table 5 After-Scenario Questionnaire

After-Scenario Questionnaire	Strongly disagree (1)	(2)	(3)	(4)	(5)	(6)	Strongly agree (7)
I am satisfied with the ease of completing the task in this scenario (ASQ1)							
I am satisfied with the amount of time it took to complete the task in this scenario (ASQ2)							
I am satisfied with the support information (on-line help, messages, documentation) when completing the task (ASQ3)							

**Figure 1 Overall design of the virtual reality educational contents.** VR: Virtual reality.


The ASQ survey also showed that the usability of the VR system was significantly higher than the traditional paper-based system. The results are shown in [Table 10](#) and [Figure 6](#).

Burnout

Among the three subscales of the MBI (emotional exhaustion, depersonalization, and personal accomplishment), emotional exhaustion and personal accomplishment improved statistically significantly from 2.53 ± 0.31 to 1.98 ± 0.17 and 2.80 ± 0.20 to 2.54 ± 0.44 , respectively. However, Depersonalization increased from 1.97 ± 0.30 to 2.01 ± 0.27 , which was not statistically significant. The results are shown in [Table 11](#) and [Figure 7](#).

Table 6 Maslach Burnout Inventory

Questions	Never	Rarely	Sometimes	Frequently	Always
I Emotional Exhaustion					
I feel emotionally drained from my work	1	2	3	4	5
I feel used up at the end of the workday	1	2	3	4	5
I feel fatigued when I get up in the morning and have to face another day on the job	1	2	3	4	5
Working with people all day is really a strain for me	1	2	3	4	5
I feel burned out from my work	1	2	3	4	5
I feel frustrated by my job	1	2	3	4	5
I feel I'm working too hard on my job	1	2	3	4	5
Working with people directly puts too much stress on me	1	2	3	4	5
I feel like I'm at the end of my rope	1	2	3	4	5
II Personal Accomplishment					
I can easily understand how my recipients feel about things	1	2	3	4	5
I deal very effectively with the problems of my recipients	1	2	3	4	5
I feel I'm positively influencing other people's lives through my work	1	2	3	4	5
I feel very energetic	1	2	3	4	5
I can easily create a relaxed atmosphere with my recipients	1	2	3	4	5
I feel exhilarated after working closely with my recipients	1	2	3	4	5
I have accomplished many worthwhile things in this job	1	2	3	4	5
In my work, I deal with emotional problems very calmly	1	2	3	4	5
III Depersonalization					
I feel I treat some recipients as if they were impersonal 'objects'	1	2	3	4	5
I've become more callous toward people since I took this job	1	2	3	4	5
I worry that this job is hardening me emotionally	1	2	3	4	5
I don't really care what happens to some recipients	1	2	3	4	5
I feel recipients blame me for some of their problems	1	2	3	4	5

**Figure 2 Diagram of patient allocation to each nurse.**

DISCUSSION

This study demonstrates the effectiveness of preoperative patient education using appropriate technology for new nurses. For new nurses, the use of VR technology proved to be superior to the traditional paper-based method in terms of satisfaction, usability and wellness. However, this does not mean that VR is the tool of choice for preoperative patient

Table 7 Comparison of satisfaction survey results of healthcare providers in VR and Control Group

	Item	Score					mean \pm SD	t	P value (n = 20)
		1	2	3	4	5			
After verbal education	1	9 (45.0)		9 (45.0)	1 (5.0)	1 (5.0)	20.50 \pm 1.76	14.44	< 0.001
	2	7 (35.0)		11 (55.0)	1 (5.0)	1 (5.0)			
	3	3 (15.0)		9 (45.0)	4 (20.0)	4 (20.0)			
	4	7 (35.0)		8 (40.0)	5 (25.0)				
	5	6 (30.0)		7 (35.0)	4 (20.0)	3 (15.0)			
	6	9 (45.0)		9 (45.0)	1 (5.0)	1 (5.0)			
	7	6 (30.0)		13 (65.0)	1 (5.0)				
After VR education	1			2 (10.0)	8 (40.0)	10 (50.0)	30.60 \pm 2.37		
	2			2 (10.0)	9 (45.0)	9 (45.0)			
	3			3 (15.0)	10 (50.0)	7 (35.0)			
	4			2 (10.0)	11 (55.0)	7 (35.0)			
	5			2 (10.0)	7 (35.0)	11 (55.0)			
	6			3 (15.0)	3 (15.0)	14 (70.0)			
	7			3 (15.0)	6 (30.0)	11 (55.0)			

VR: Virtual reality.

Table 8 Comparison of satisfaction results of each item

		Number	mean \pm SD	t	P value
Item 1	Verbal education	20	2.70 \pm 0.80	8.23	< 0.001
	VR education	20	4.40 \pm 0.68		
Item 2	Verbal education	20	2.80 \pm 0.77	6.31	< 0.001
	VR education	20	4.35 \pm 0.67		
Item 3	Verbal education	20	3.45 \pm 1.00	3.00	0.003
	VR education	20	4.20 \pm 0.70		
Item 4	Verbal education	20	2.90 \pm 0.79	5.81	< 0.001
	VR education	20	4.25 \pm 0.64		
Item 5	Verbal education	20	3.20 \pm 1.06	4.19	< 0.001
	VR education	20	4.45 \pm 0.69		
Item 6	Verbal education	20	2.70 \pm 0.80	7.59	< 0.001
	VR education	20	4.55 \pm 0.76		
Item 7	Verbal education	20	2.75 \pm 0.55	7.47	< 0.001
	VR education	20	4.40 \pm 0.75		

VR: Virtual reality.

education. Although the VR method scores higher on the System Usability Scale than the traditional paper-based method, it is only average at 68 points[26]. The System Usability Scale is not a diagnostic method, so a low System Usability Scale means that we need to further identify and improve the problem.

VR has been used in many healthcare fields. It has been used to train medical students[3], for preoperative planning and simulation by surgeons[27,28], and for rehabilitation therapy[29] and psychotherapy[30]. There are also studies showing the benefits of using VR to educate patients before surgery[7]. However, there is a lack of study on how beneficial the use of suitable technology is in terms of usability for healthcare professionals (especially for new nurses

Table 9 Comparison of traditional paper-based verbal method and virtual reality method using System Usability Scale

Variables	Verbal education, mean \pm SD	VR education, mean \pm SD	<i>t</i>	<i>P</i> value
SUS1	2.35 \pm 1.04	4.10 \pm 0.72	-7.68	< 0.001
SUS2	3.90 \pm 0.79	2.65 \pm 0.81	4.80	< 0.001
SUS3	2.25 \pm 0.72	4.10 \pm 0.79	-7.00	< 0.001
SUS4	4.00 \pm 0.79	2.75 \pm 0.79	4.63	< 0.001
SUS5	2.10 \pm 0.91	4.30 \pm 0.73	-7.03	< 0.001
SUS6	4.00 \pm 0.79	2.75 \pm 0.91	5.22	< 0.001
SUS7	2.05 \pm 0.89	4.30 \pm 0.66	-8.64	< 0.001
SUS8	4.10 \pm 0.72	2.90 \pm 0.85	4.33	< 0.001
SUS9	2.05 \pm 0.83	4.00 \pm 0.73	-7.06	< 0.001
SUS10	3.95 \pm 0.76	2.55 \pm 0.83	5.98	< 0.001
Score of SUS	27.13 \pm 8.82	68.00 \pm 5.42	-16.79	< 0.001

SUS: System Usability Scale; VR: Virtual reality.

Table 10 Comparison of traditional paper-based verbal method and virtual reality method using After-Scenario Questionnaire

Variables	Verbal education, mean \pm SD	VR education, mean \pm SD	<i>t</i>	<i>P</i> value
ASQ1	4.05 \pm 1.19	5.75 \pm 0.97	-5.23	< 0.001
ASQ2	2.75 \pm 1.12	5.55 \pm 1.00	-11.33	< 0.001
ASQ3	2.95 \pm 1.10	3.85 \pm 0.99	-4.72	< 0.001
Average of ASQ	3.25 \pm 0.59	5.05 \pm 0.61	-10.59	< 0.001

ASQ: After-Scenario Questionnaire; VR: Virtual reality.

Table 11 Comparison of traditional paper-based verbal method and virtual reality method using Maslach Burnout Inventory

Variables	Verbal education, mean \pm SD	VR education, mean \pm SD	<i>t</i>	<i>P</i> value
Emotional exhaustion	2.53 \pm 0.31	1.98 \pm 0.17	6.77	< 0.001
Personal accomplishment	2.80 \pm 0.20	2.54 \pm 0.44	2.28	0.034
Depersonalization	1.97 \pm 0.30	2.01 \pm 0.27	-0.54	0.599

VR: Virtual reality.

who are new to the job).

The problem with traditional paper-based education is that it must be repeated for each patient and cannot be performed concurrently when there are several patients. Numerous patients can be managed simultaneously if preoperative patient education in virtual reality is performed using a head-mounted display, as the present study. Additionally, the quality of education, which varies according to the healthcare professionals' abilities, has the benefit of remaining consistent. From the perspective of healthcare providers, patients may receive suitable and easy-to-understand information more readily than through conventional methods, and there may be a positive impact on lowering physical and mental stress.

Because it is simple to use and has high usability for new nurses who are challenging at work, virtual reality technology will assist improve the quality of nursing care. Since preoperative patient education using virtual reality technology is beneficial for patients and usable for medical staff (especially new nurses), it should be actively considered for acceptance and adoption.

The MBI was used in the study to compare traditional methods of patient education with virtual reality technology. The findings provided important insights into the impact of these two methods of instruction on the burnout levels of nurses, as evaluated by the three MBI subscales: Emotional exhaustion, depersonalization, and personal accomplishment. The first important finding was a statistically significant decrease in emotional exhaustion levels. Nurses who provided



Figure 3 Two patients in the experimental group being educated in virtual reality under a nurse's supervision.




Figure 4 Bar plots with error bars of the satisfaction score of each item virtual reality. VR: Virtual reality.

patient education *via* virtual reality technology exhibited a decrease in emotional exhaustion, dropping from 2.53 ± 0.31 to 1.98 ± 0.17 . This is consistent with other studies that shown the use of VR technology reduces emotional weariness among ENT residents[31]. A decrease in emotional exhaustion is critical because it implies a possible improvement in emotional well-being and resilience in the face of the emotionally demanding nature of nursing. Similarly, the personal accomplishment subscale improved statistically significantly. Scores improved from 2.80 ± 0.20 to 2.54 ± 0.44 , indicating an improvement in nurses' sense of accomplishment and efficacy. This advancement suggests that virtual reality methods of education contributed to the nursing staff's sense of professional growth and achievement. Such beneficial changes can be critical in increasing work happiness and lowering burnout, eventually improving not only nurses but also the quality of patient care. The results of the depersonalization subscale, on the other hand, were intriguing. The increase in depersonalization scores from 1.97 ± 0.30 to 2.01 ± 0.27 was not statistically significant. Because of the lack of statistical significance, virtual reality-based education appears to have had little effect on reducing depersonalization. Depersonalization is defined as an unfeeling and impersonal attitude to those who receive one's service, care, treatment, or education. Because there is little face-to-face interaction with the patient, the use of VR is not regarded to be a major improvement over the traditional method of nurses interacting and explaining to patients in person. It is necessary to investigate more why depersonalization was mostly unaffected and whether further interventions or modifications to methods of education are required to address this component of burnout.

The limitations of this study are obvious. These are as follows:

Patients with visual impairments or claustrophobia may find it challenging to understand the information provided in virtual reality. This potentially leads to biased satisfaction and usability ratings that do not accurately reflect the general patient population.

While the System Usability Scale and the After-Scenario Questionnaire are time-tested, valid, and reliable tools, they are not diagnostic tools in and of themselves, necessitating additional inquiry into the actual causes for low scores owing




Figure 5 Bar plot with error bar showing comparison of traditional paper-based method and VR method using System Usability Scale. VR: Virtual reality.




Figure 6 Bar plot with error bar showing comparison of traditional paper-based method and VR method using After-Scenario Questionnaire. VR: Virtual reality.

to a lack of context in the results.

The traditional paper-based method may be more intuitive to use than the VR one for experienced nurses as compared to younger, more familiar with technology new nurses. This difference could skew the results.

It is expensive to acquire equipment and provide VR content appropriate for each hospital environment. This financial barrier could limit the widespread adoption of VR, affecting the generalizability of the study findings to other settings that may not have the resources to implement such technology.

Even healthcare providers who have previously explained this need time to get trained on how to use the equipment and contents. This could introduce variability in how effectively VR is implemented across different users, potentially impacting the consistency and reliability of the study's findings.

If the VR content needs to be changed owing to a change in the situation in the hospital, additional costs for reproducing the contents may be incurred. This limitation may affect the long-term feasibility and sustainability of VR as a patient education tool, influencing the study's implications for future practice.

The usability survey of a small number of healthcare providers ($n = 20$) was assessed using a subjective questionnaire; however, it would be preferable if actual stress reduction could be demonstrated through more objective indications.




Figure 7 Bar plot with error bar showing comparison of traditional paper-based verbal method and virtual reality method using Maslach Burnout Inventory. MBI: Maslach Burnout Inventory; VR: Virtual reality.

CONCLUSION

This study provides a valuable reference for future research regarding the possible advantages and drawbacks of employing VR technology in a preoperative environment. In addition, the study provides valuable insights into the impact of patient education using VR technology on nurse burnout. Despite above limitations, the study demonstrates that VR technology is significantly more usable than traditional paper-based methods for healthcare providers (especially for new nurses) working in plastic surgery wards. In addition, it demonstrates the usefulness of virtual reality-based patient education in lowering emotional exhaustion and increasing personal accomplishment of nurses. However, the persistence of depersonalization in lack of a statistically significant changes implies that a more nuanced evaluation of educational content and methods to address this particular aspect of burnout among nurses is required. As this study has the limitation of being a subjective survey, it is crucial to assess qualitative changes to confirm the efficacy to the quality of nursing care, such as the change in the number of questions or the amount of time required for a preoperative patient education when the healthcare provider provides preoperative patient education using VR. Additionally, further research is required to confirm the impact of objective vital signs of healthcare providers, such as stress signs or heart rate variability, when the healthcare provider is offering the preoperative patient education.

FOOTNOTES

Author contributions: Kwon H and Oh SH contributed to study conception and design; Kim J and Kim D contributed to data collection; Kim J contributed to data analysis and interpretation; All authors contributed to drafting of the article; Kwon H and Oh SH contributed to critical revision of the article. Kwon H and Oh SH conceptualized and designed the research; Kim J and Kim D screened patients and acquired clinical data; Kim J contributed to data analysis; Kim J, Kwon H and Oh SH wrote the paper. All the authors have read and approved the final manuscript. Both Kwon H and Oh SH have played important and indispensable roles in the experimental design, data interpretation and manuscript preparation as the co-corresponding authors. Oh SH applied for and obtained the funds for this research project. Kwon H and Oh SH conceptualized, designed, and supervised the whole process of the project. They searched the literature, revised and submitted the early version of the manuscript. Kwon H was instrumental and responsible for data re-analysis and re-interpretation, figure plotting, comprehensive literature search, preparation and submission of the current version of the manuscript. This collaboration between Kwon H and Oh SH is crucial for the publication of this manuscript and other manuscripts still in preparation.

Supported by Research Fund of Chungnam National University, Chungnam National University, the Ministry of Trade, Industry, and Energy, Korea, under the “Regional industry-based organization support program”, No. P0001940; the Korea Institute for Advancement of Technology, and a grant of the Korea Health Technology R&D Project through the Korea Health Industry Development Institute, funded by the Ministry of Health & Welfare, Republic of Korea, No. HI20C2088.

Institutional review board statement: This study was approved by the Research Ethics Board of Chungnam National University Hospital (approval no. CNUH 2019-06-008-002).

Clinical trial registration statement: Trial registration CRIS, KCT0007489. Registered June 30, 2022. <http://cris.nih.go.kr/cris/>.

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: No potential conflict of interest relevant to this article was reported.

Data sharing statement: No additional data are available.

CONSORT 2010 statement: The authors have read the CONSORT 2010 statement, and the manuscript was prepared and revised according to the CONSORT 2010 statement.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: South Korea

ORCID number: Jiyoung Kim 0000-0002-2021-5290; Donghyun Kim 0000-0001-8296-9920; Sang-Ha Oh 0000-0003-3734-5005; Hyeokjae Kwon 0000-0002-1418-3448.

S-Editor: Liu JH

L-Editor: A

P-Editor: Zhang XD

REFERENCES

- Li L, Yu F, Shi D, Shi J, Tian Z, Yang J, Wang X, Jiang Q. Application of virtual reality technology in clinical medicine. *Am J Transl Res* 2017; **9**: 3867-3880 [PMID: 28979666]
- de Vries FS, van Dongen RTM, Bertens D. Pain education and pain management skills in virtual reality in the treatment of chronic low back pain: A multiple baseline single-case experimental design. *Behav Res Ther* 2023; **162**: 104257 [PMID: 36731183 DOI: 10.1016/j.brat.2023.104257]
- Pottle J. Virtual reality and the transformation of medical education. *Future Healthc J* 2019; **6**: 181-185 [PMID: 31660522 DOI: 10.7861/fhj.2019-0036]
- Ahmadpour N, Randall H, Choksi H, Gao A, Vaughan C, Poronnik P. Virtual Reality interventions for acute and chronic pain management. *Int J Biochem Cell Biol* 2019; **114**: 105568 [PMID: 31306747 DOI: 10.1016/j.biocel.2019.105568]
- Howard MC. A meta-analysis and systematic literature review of virtual reality rehabilitation programs. *CHB* 2017; **70**: 317-327 [DOI: 10.1016/j.chb.2017.01.013]
- Bekelis K, Calnan D, Simmons N, MacKenzie TA, Kakoulides G. Effect of an Immersive Preoperative Virtual Reality Experience on Patient Reported Outcomes: A Randomized Controlled Trial. *Ann Surg* 2017; **265**: 1068-1073 [PMID: 27906757 DOI: 10.1097/SLA.0000000000002094]
- Kwon H, Lee J, Park YS, Oh SH, Kim J. Effects of preoperative education using virtual reality on preoperative anxiety and information desire: a randomized clinical trial. *J Clin Monit Comput* 2023; **37**: 1401-1407 [PMID: 36933168 DOI: 10.1007/s10877-023-00988-5]
- Shepherd T, Trinder M, Theophilus M. Does virtual reality in the perioperative setting for patient education improve understanding? A scoping review. *Surg Pract Sci* 2022; 100101 [DOI: 10.1016/j.sipas.2022.100101]
- Bazi KN. Effective Training for Novice Nurses Transition to the Perioperative Role: An Integrative Literature Review. 2019
- Fitzpatrick E, Hyde A. Nurse-related factors in the delivery of preoperative patient education. *J Clin Nurs* 2006; **15**: 671-677 [PMID: 16684162 DOI: 10.1111/j.1365-2702.2006.01327.x]
- Wu TY, Fox DP, Stokes C, Adam C. Work-related stress and intention to quit in newly graduated nurses. *Nurse Educ Today* 2012; **32**: 669-674 [PMID: 21983395 DOI: 10.1016/j.nedt.2011.09.002]
- Kim MH, Mazenga AC, Simon K, Yu X, Ahmed S, Nyasulu P, Kazembe PN, Ngoma S, Abrams EJ. Burnout and self-reported suboptimal patient care amongst health care workers providing HIV care in Malawi. *PLoS One* 2018; **13**: e0192983 [PMID: 29466443 DOI: 10.1371/journal.pone.0192983]
- Chandawarkar A, Chaparro JD. Burnout in clinicians. *Curr Probl Pediatr Adolesc Health Care* 2021; **51**: 101104 [PMID: 34789423 DOI: 10.1016/j.cppeds.2021.101104]
- Bondagji D, Fakeerh M, Alwafi H, Khan AA. The Effects of Long Working Hours on Mental Health Among Resident Physicians in Saudi Arabia. *Psychol Res Behav Manag* 2022; **15**: 1545-1557 [PMID: 35757011 DOI: 10.2147/PRBM.S370642]
- Zarei S, Fooladvand K. Mediating effect of sleep disturbance and rumination on work-related burnout of nurses treating patients with coronavirus disease. *BMC Psychol* 2022; **10**: 197 [PMID: 35962404 DOI: 10.1186/s40359-022-00905-6]
- von Känel R, Princip M, Holzgang SA, Fuchs WJ, van Nuffel M, Pazhenkottil AP, Spiller TR. Relationship between job burnout and somatic diseases: a network analysis. *Sci Rep* 2020; **10**: 18438 [PMID: 33116176 DOI: 10.1038/s41598-020-75611-7]
- Hamaideh SH. Burnout, social support, and job satisfaction among Jordanian mental health nurses. *Issues Ment Health Nurs* 2011; **32**: 234-242 [PMID: 21355758 DOI: 10.3109/01612840.2010.546494]
- Aiken LH, Clarke SP, Sloane DM, Sochalski J, Silber JH. Hospital nurse staffing and patient mortality, nurse burnout, and job dissatisfaction. *JAMA* 2002; **288**: 1987-1993 [PMID: 12387650 DOI: 10.1001/jama.288.16.1987]
- Olsson M, Cosgrove CM, Wall MM, Blanco C. Suicide Risks of Health Care Workers in the US. *JAMA* 2023; **330**: 1161-1166 [PMID: 37750880 DOI: 10.1001/jama.2023.15787]
- Gold JJ, Annick ET, Lane AS, Ho K, Marty RT, Espinoza JC. "Doc McStuffs: Doctor for a Day" Virtual Reality (DocVR) for Pediatric Preoperative Anxiety and Satisfaction: Pediatric Medical Technology Feasibility Study. *J Med Internet Res* 2021; **23**: e25504 [PMID: 33730687 DOI: 10.2196/25504]
- Llorens R, Colomer-Font C, Alcañiz M, Noé-Sebastián E. BioTrak virtual reality system: Effectiveness and satisfaction analysis for balance rehabilitation in patients with brain injury. *Neurología* 2013; **28**: 268-275 [DOI: 10.1016/j.nrleng.2012.04.016]
- Benner P. From novice to expert. *Menlo Park* 1984; **84**: 10-1097 [DOI: 10.1097/00000446-198412000-00027]

- 23 **Brooke J.** SUS: A “quick and dirty” usability Scale. 1st Edition, Usability evaluation in industry. 1996; 189: 189-194
- 24 **Lewis JR.** IBM computer usability satisfaction questionnaires: Psychometric evaluation and instructions for use. *Int J Hum-Comput Int* 1995; 7: 57-78 [DOI: [10.1080/10447319509526110](https://doi.org/10.1080/10447319509526110)]
- 25 **Maslach C, Jackson SE, Leiter MP.** Maslach burnout inventory: Scarecrow Education; 1997
- 26 **Melnick ER, Dyrbye LN, Sinsky CA, Trockel M, West CP, Nedelec L, Tutty MA, Shanafelt T.** The Association Between Perceived Electronic Health Record Usability and Professional Burnout Among US Physicians. *Mayo Clin Proc* 2020; **95**: 476-487 [PMID: [31735343](https://pubmed.ncbi.nlm.nih.gov/31735343/) DOI: [10.1016/j.mayocp.2019.09.024](https://doi.org/10.1016/j.mayocp.2019.09.024)]
- 27 **Ulbrich M, Van den Bosch V, Bönsch A, Gruber LJ, Ooms M, Melchior C, Motmaen I, Wilpert C, Rashad A, Kuhlen TW, Hölzle F, Puladi B.** Advantages of a Training Course for Surgical Planning in Virtual Reality for Oral and Maxillofacial Surgery: Crossover Study. *JMIR Serious Games* 2023; **11**: e40541 [PMID: [36656632](https://pubmed.ncbi.nlm.nih.gov/36656632/) DOI: [10.2196/40541](https://doi.org/10.2196/40541)]
- 28 **Freidin D, Singolda R, Tejman-Yarden S, Parmat I, Liran A, Ofir H, Saukhat O, Haik J, Barnea Y, Tessone A.** Using Virtual Reality for Deep Inferior Epigastric Perforator Flap Preoperative Planning. *Plast Reconstr Surg Glob Open* 2023; **11**: e4773 [PMID: [36660058](https://pubmed.ncbi.nlm.nih.gov/36660058/) DOI: [10.1097/GOX.0000000000004773](https://doi.org/10.1097/GOX.0000000000004773)]
- 29 **Phelan I, Carrion-Plaza A, Furness PJ, Dimitri P.** Home-based immersive virtual reality physical rehabilitation in paediatric patients for upper limb motor impairment: a feasibility study. *Virtual Real* 2023; **27**: 1-16 [PMID: [36686613](https://pubmed.ncbi.nlm.nih.gov/36686613/) DOI: [10.1007/s10055-023-00747-6](https://doi.org/10.1007/s10055-023-00747-6)]
- 30 **Riches S, Jeyarajaguru P, Taylor L, Fialho C, Little J, Ahmed L, O'Brien A, van Driel C, Veling W, Valmaggia L.** Virtual reality relaxation for people with mental health conditions: a systematic review. *Soc Psychiatry Psychiatr Epidemiol* 2023; **58**: 989-1007 [PMID: [36658261](https://pubmed.ncbi.nlm.nih.gov/36658261/) DOI: [10.1007/s00127-022-02417-5](https://doi.org/10.1007/s00127-022-02417-5)]
- 31 **Weitzman RE, Wong K, Worrall DM, Park C, McKee S, Tufts RE, Teng MS, Illoreta AM.** Incorporating Virtual Reality to Improve Otolaryngology Resident Wellness: One Institution's Experience. *Laryngoscope* 2021; **131**: 1972-1976 [PMID: [33764531](https://pubmed.ncbi.nlm.nih.gov/33764531/) DOI: [10.1002/lary.29529](https://doi.org/10.1002/lary.29529)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

