World Journal of *Gastrointestinal Endoscopy*

World J Gastrointest Endosc 2024 October 16; 16(10): 545-580





Published by Baishideng Publishing Group Inc

World Journal of Gastrointestinal Endoscopy

Contents

Monthly Volume 16 Number 10 October 16, 2024

ORIGINAL ARTICLE

Retrospective Study

Endoscopic "calabash" ligation and resection for small gastric mesenchymal tumors 545 Lin XM, Peng YM, Zeng HT, Yang JX, Xu ZL

Prospective Study

557 Gastric fundoplication with endoscopic technique: A novel approach for gastroesophageal reflux disease treatment

Gadour E, Hoff AC

META-ANALYSIS

Safety and efficacy of peroral endoscopic myotomy for treating achalasia in pediatric and geriatric 566 patients: A meta-analysis

Pu XX, Huang S, Zhong CY, Wang X, Fu SF, Lv YQ, Zou K, Lü MH, Peng Y, Tang XW



Contents

Monthly Volume 16 Number 10 October 16, 2024

ABOUT COVER

Editorial Board Member of World Journal of Gastrointestinal Endoscopy, Takashi Murakami, MD, PhD, Associate Professor, Department of Gastroenterology, Juntendo University School of Medicine, Tokyo 113-8421, Japan. t-murakm@juntendo.ac.jp

AIMS AND SCOPE

The primary aim of World Journal of Gastrointestinal Endoscopy (WJGE, World J Gastrointest Endosc) is to provide scholars and readers from various fields of gastrointestinal endoscopy with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGE mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal endoscopy and covering a wide range of topics including capsule endoscopy, colonoscopy, double-balloon enteroscopy, duodenoscopy, endoscopic retrograde cholangiopancreatography, endosonography, esophagoscopy, gastrointestinal endoscopy, gastroscopy, laparoscopy, natural orifice endoscopic surgery, proctoscopy, and sigmoidoscopy.

INDEXING/ABSTRACTING

The WJGE is now abstracted and indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for WJGE as 1.4; JIF without journal self cites: 1.4; 5-year JIF: 1.7; JIF Rank: 111/143 in gastroenterology and hepatology; JIF Quartile: Q4; and 5-year JIF Quartile: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yi-Xuan Cai; Production Department Director: Xu Guo; Cover Editor: Jia-Ping Yan.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Gastrointestinal Endoscopy	https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 1948-5190 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
October 15, 2009	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Monthly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Bing Hu, JooYoung Cho	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/1948-5190/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
October 16, 2024	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com
PUBLISHING PARTNER	PUBLISHING PARTNER'S OFFICIAL WEBSITE
Digestive Endoscopy Center of West China Hospital, SCU	http://www.cd120.com/index.html

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



E WU

World Journal of Gastrointestinal Endoscopy

Submit a Manuscript: https://www.f6publishing.com

World J Gastrointest Endosc 2024 October 16; 16(10): 566-580

DOI: 10.4253/wjge.v16.i10.566

ISSN 1948-5190 (online)

META-ANALYSIS

Safety and efficacy of peroral endoscopic myotomy for treating achalasia in pediatric and geriatric patients: A meta-analysis

Xin-Xin Pu, Shu Huang, Chun-Yu Zhong, Xia Wang, Su-Fen Fu, Ying-Qin Lv, Kang Zou, Mu-Han Lü, Yan Peng, Xiao-Wei Tang

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification Scientific Quality: Grade B, Grade D, Grade D, Grade D Novelty: Grade B, Grade C, Grade D, Grade D Creativity or Innovation: Grade B, Grade C, Grade D, Grade D Scientific Significance: Grade A, Grade C, Grade C, Grade D

P-Reviewer: Jankovic J; Tawheed A; Wang Y

Received: May 24, 2024 Revised: August 17, 2024 Accepted: September 14, 2024 Published online: October 16, 2024 Processing time: 144 Days and 1.5 Hours



Xin-Xin Pu, Xia Wang, Su-Fen Fu, Department of Gastroenterology, Dechang People's Hospital of Sichuan Province, Liangshan 615500, Sichuan Province, China

Xin-Xin Pu, Ying-Qin Lv, Kang Zou, Mu-Han Lü, Yan Peng, Xiao-Wei Tang, Department of Gastroenterology, The Affiliated Hospital of Southwest Medical University, Luzhou 646000, Sichuan Province, China

Shu Huang, Department of Gastroenterology, Lianshui People's Hospital of Kangda College Affiliated to Nanjing Medical University, Huaian 223499, Jiangsu Province, China

Chun-Yu Zhong, Department of Ultrasound, The Affiliated Hospital of Southwest Medical University, Luzhou 646000, Sichuan Province, China

Co-first authors: Xin-Xin Pu and Shu Huang.

Corresponding author: Xiao-Wei Tang, MD, PhD, Associate Professor, Postdoc, Department of Gastroenterology, The Affiliated Hospital of Southwest Medical University, No. 25 Taiping Street, Jiangyang Region, Luzhou 646000, Sichuan Province, China. solitude5834@hotmail.com

Abstract

BACKGROUND

As a less invasive technique, peroral endoscopic myotomy (POEM) has recently been widely accepted for treating achalasia with an excellent safety profile, durability, and efficacy in adults. In pediatric and geriatric patients, the treatment is more difficult.

AIM

To discuss the clinical outcomes of POEM in pediatric and geriatric patients with achalasia.

METHODS

We conducted a comprehensive search of PubMed, Embase and Cochrane Library databases from inception to July 2024. The primary outcomes were technical and clinical success. Secondary outcomes of interest included adverse events and gastroesophageal reflux disease (GERD). The pooled event rates were calculated by comprehensive meta-analysis software.

Raisbideng® WJGE | https://www.wjgnet.com

RESULTS

A total of 32 studies with 547 pediatric patients and 810 geriatric patients were included in this study. The pooled event rates of technical success, clinical success, GERD and adverse events of POEM for treating achalasia in pediatric patients were 97.1% [95% confidence interval (CI): 95.0%-98.3%; $l^2 = 0\%$; P < 0.000], 93.2% (95%CI: 90.5%-95.2%; $l^2 = 0\%$; P < 0.000), 22.3% (95%CI: 18.4%-26.7%; $l^2 = 43.874\%$; P < 0.000) and 20.4% (95%CI: 16.6%-24.8%; $l^2 = 67.217\%$; P < 0.000), respectively. Furthermore, in geriatric patients, the pooled event rates were 97.7% (95%CI: 95.8%-98.7%; $l^2 = 15.200\%$; P < 0.000), 93.2% (95%CI: 90.3%-95.2%; $l^2 = 0\%$; P < 0.000), 23.9% (95%CI: 19.4%-29.1%; $l^2 = 75.697\%$; P < 0.000) and 10.8% (95%CI: 8.3%-14.0%; $l^2 = 62.938\%$; P < 0.000], respectively.

CONCLUSION

Our findings demonstrated that POEM was an effective and safe technique for pediatric and geriatric patients with achalasia.

Key Words: Achalasia; Peroral endoscopic myotomy; Pediatric; Geriatric; Meta-analysis

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Peroral endoscopic myotomy (POEM), a minimally invasive procedure, has gained substantial acceptance as a treatment for achalasia due to its excellent safety profile, durability, and efficacy in adults. We conducted this systematic review and meta-analysis to summarize the durability, safety, and efficacy of POEM for treating achalasia in pediatric and geriatric patients. Our results indicated that POEM was an effective and safe technique for pediatric and geriatric patients with achalasia.

Citation: Pu XX, Huang S, Zhong CY, Wang X, Fu SF, Lv YQ, Zou K, Lü MH, Peng Y, Tang XW. Safety and efficacy of peroral endoscopic myotomy for treating achalasia in pediatric and geriatric patients: A meta-analysis. *World J Gastrointest Endosc* 2024; 16(10): 566-580

URL: https://www.wjgnet.com/1948-5190/full/v16/i10/566.htm **DOI:** https://dx.doi.org/10.4253/wjge.v16.i10.566

INTRODUCTION

Achalasia is a moderately uncommon esophageal smooth muscle motility disorder characterized by the absence or spastic contractions of the esophageal body and the loss of deglutition-induced relaxation of the lower esophageal sphincter (LES)[1]. Typically, dysphagia is experienced by the patient. Regurgitation of undigested food, heartburn, respiratory symptoms, chest pain, and weight loss are other clinical symptoms of achalasia[2]. The estimated annual incidence rate of achalasia is between 0.03 and 1.63 *per* 100000 people, and the annual prevalence of achalasia is reported to range from 1.8 to 12.6 *per* 100000 people[3]. Achalasia can occur at all ages and equally in men and women, with no racial predilection [3], but it is an exceedingly rare illness in the pediatric population, with an estimated prevalence between 0.02 and 0.31 *per* 100000 children, approximately 10 times lower than that in adults[4-6]. The prevalence increases with age, with a peak in the seventh decade of life[7].

Botulinum toxin injection, laparoscopic Heller myotomy (LHM) and pneumatic dilation (PD) are available therapeutic approaches for achalasia[8-11]. In the past decade, a novel technique called peroral endoscopic myotomy (POEM) was created to combine an endoscopic approach with the principles of natural orifice transluminal endoscopic surgery to perform myotomy for achalasia. The American College of Gastroenterology clinical guidelines indicate that POEM and LHM result in comparable symptomatic improvements in patients with achalasia[3]. A systematic review and meta-analysis compared the results between 1958 patients after POEM and 5834 patients after LHM, and reported that the predicted probability of improvement in dysphagia was 93.5% for POEM and 91.0% for LHM at 12 months, and 92.7% for POEM and 90.0% for LHM at 24 months[12].

Briefly, the POEM procedure requires submucosal injection with indigo carmine dye, mucosal incision, submucosal dissection, myotomy and closure of the mucosal incision[3]. The POEM procedure in special populations is similar to that in general patients, with only minor modifications. However, the length of the pediatric esophagus is short, and the esophageal wall in children is relatively thin, which undoubtedly increases the difficulty of the operation[13]. In addition, geriatric patients tend to have more comorbidities, and thus have a higher risk during surgery and anesthesia[14,15].

Therefore, we aimed to conduct this systematic review and meta-analysis to summarize the durability, safety and efficacy of POEM in pediatric and geriatric patients with achalasia.

Saisbideng® WJGE | https://www.wjgnet.com

MATERIALS AND METHODS

This systematic review and meta-analysis strictly conformed to the preferred reporting items for systematic reviews and meta-analyses statement^[16]. Our study did not require ethical approval or written consent.

Search strategy

A systematic literature search was conducted via the PubMed, Embase and Cochrane Library databases from inception to July 2024, with studies limited to those written in the English language. The key words included "peroral endoscopic myotomy", "per-oral endoscopic myotomy", "POEM", "achalasia" and all possible combinations (Supplementary Table 1). To ensure a thorough search of the literature, the words "pediatric" and "geriatric" were not used. In addition to searching the literature using keywords, the inclusion of articles following a meticulous review of references was pursued to identify those that satisfied the specified criteria for inclusion.

Study selection

Two authors screened the titles and abstracts of all articles separately, in accordance with the exclusion and inclusion criteria. Next, the full texts of relevant articles were reviewed after screening. Any disagreements between reviewers in the search process were resolved by discussion with a third reviewer. The inclusion criteria were as follows: (1) Population: Patients diagnosed with achalasia and aged ≤ 18 years (pediatric) or ≥ 60 years (geriatric); (2) Treatment: POEM; and (3) Outcomes: Technical and clinical success, adverse events and gastroesophageal reflux disease (GERD). Exclusion criteria were as follows: (1) Case reports with < 5 patients, reviews, and animal experiments; (2) Studies that did not provide enough data; and (3) Studies not published in the English language.

Data extraction and definition

Data regarding the characteristics of the selected studies (first author, year of publication, country, study type, study interval, number of patients, gender and age), pre- and postoperative data of the patients (duration of symptoms, myotomy length, operation time, length of hospital stay and follow-up time), and clinical outcomes (technical success and clinical success rates, adverse events and GERD rates) were independently extracted by two authors using a prepared standardized form. The definition of technical success was completion of the entire POEM procedure. Clinical success was regarded as an Eckardt score \leq 3 during the follow-up period after POEM. The severity of adverse events was graded on the basis of the American Society for Gastrointestinal Endoscopy (ASGE) lexicon[17]. GERD included symptomatic reflux and reflux esophagitis.

Quality assessment of the studies

Two reviewers independently conducted a quality assessment using the National Institutes of Health (NIH) quality assessment tool, which is applicable for pre-post studies with no control group[18].

Statistical analysis

Statistical analysis was performed using comprehensive meta-analysis software version 3.0 (Biostat, Englewood, NJ, United States). The outcomes are presented as pooled event rates and 95% confidence interval (CI), and the significance degree P was set at < 0.05. Heterogeneity among the studies was assessed using the l^2 statistic. Significant heterogeneity was considered, if the l^2 value was \geq 50%, and a random-effect model was used. A funnel plot was used to evaluate publication bias.

RESULTS

Eligible studies

A total of 4620 articles was identified in the initial search, and 3206 studies remained after the removal of duplicates. On the basis of the inclusion and exclusion criteria, 41 articles were eligible after the titles and abstracts were reviewed. Nine studies were excluded as the cohorts included fewer than five patients or due to overlapping publications. Ultimately, 32 articles were included in this review (pediatric patients, n = 20[19-38]; geriatric patients, n = 12[13,39-49]) (Figure 1).

According to the NIH quality assessment tool, 13 studies were of good quality, eight studies showed fair quality, and one study exhibited poor quality (Supplementary Table 2 and Supplementary Table 3). The reasons why the study was rated as poor quality were as follows: The eligibility criteria and study population were not clearly described; Only five pediatric patients who underwent POEM were enrolled in this study; The definition of outcomes was not clearly described; Blinding or masking means were used for the outcome evaluations; P values were not reported. Due to the lack of information in articles published in conference abstracts, no quality assessment was conducted.

Role of POEM for achalasia in pediatric patients

Baseline characteristics: Table 1 shows the baseline characteristics of the original studies of POEM for treating achalasia in pediatric patients [19-38]. Twenty articles involving 547 pediatric patients investigated the efficacy and safety of POEM, including 5 prospective studies and 15 retrospective studies. Among these articles, seven studies were conducted in China, four in the United States, two in Japan, two in Italy, two in India, one in France, one in Israel and one in Chile. These studies were performed between 2007 and 2021. The age of the patients in these studies ranged from 0.9 to 18 years,



Table 1 Baseline characteristics of the studies on peroral endoscopic myotomy for achalasia in pediatric patients												
Ref.	Country	Study type	Study interval	Patients (<i>n</i>)	Gender (M:F)	Age (yr)	Duration of symptoms (month)	Myotomy length (cm)	Operation time (min)	Hospital stay (day)	Follow-up (month)	Method of GERD diagnosis
Li et al[<mark>19</mark>], 2015	China	Prospective	October 2011 to March 2014	9	4:5	14.1 (10-17)	26.4 (6-60)	8.3	56.7	-	16.3 (3-30)	Esophageal manometry, barium esophagram and EGD
Chen <i>et al</i> [20], 2015	China	Prospective	August 2010 to July 2012	27	11:16	13.8 (6-17)	20.4 (6-36)	9.6 (7-11)	39.4 (21-90)	3.2 (1-7)	24.6 (15-38)	EGD
Caldaro <i>et al</i> [21], 2015	Italy	Retrospective	2009 to 2014	9	3:6	12.2 ± 3.8	-	11 ± 2	62 ± 12.7	4.1 (2-7)	12.7 (5-28)	pH-monitoring and EGD
Tang et al[<mark>22</mark>], 2015	China	Retrospective	July 2012 to August 2014	5	3:2	15	12 (3-15)	8 (6-11)	50 (40-90)	7 (5-13)	18 (12-23)	EGD, mano-metry
Tan <i>et al</i> [23], 2016	China	Retrospective	January 2007 to June 2015	12	6:6	13.7 ± 2.6	23.6 ± 16.8	-	-	-	36	EGD, esophageal manometry
Stavropoulos <i>et al</i> [24], 2017	United States	Retrospective ¹	2013 to 2016	10	7:3	14.7 (10-17)	21 (3-84)	11.4 ± 5.98	55 (33-111)	1.2 (1-2)	15 (1-30)	-
Zangen <i>et al</i> [25], 2017	Israel	Retrospective ¹	-	5	2:3	15.4 (10- 18)	-	11.2 (10-14)	62 (43-73)	-	6	HRM
Kethman <i>et al</i> [26], 2018	United States	Prospective	2014 to 2016	10	8:2	13.4 ± 3.3	-	7 (4-9)	142 (60-259)	-	-	-
Miao <i>et al</i> [27], 2018	China	Prospective	October 2014 to October 2016	21	9:12	5.5 (0.9-18)	18 (3.6-30)	9 (6-11)	40 (30-55)	9 (7-12)	13.2 (3-24)	99mTc DTPA scintigraphic examin- ations, gastroscopy and esophageal manometry
Korrapati <i>et al</i> [<mark>28</mark>], 2018	India	Retrospective ¹	-	15	10:5	15 (3-18)	21.9 (6-54)	8 (6-11)	85.3 ± 31	4.4 ± 2.5	19.8 (1.5-51)	EGD
Nishimoto <i>et al</i> [29], 2018	Japan	Retrospective ¹	May 2015 to November 2017	13	-	15 (10-18)	-	-	-	-	18.3 (0-30)	-
Mangiola <i>et al</i> [<mark>30]</mark> , 2018	Italy	Retrospective ¹	January 2012 to June 2017	26	12:14	10.9 (2-17)	18.2 ± 14.9	10 ± 2.6	56.2 ± 12.6	3.7 ± 1.7	30.2 ± 15.4	Manometry, 24 h pH-monitoring, and EGD
Yamashita <i>et al</i> [<mark>31</mark>], 2018	Japan	Retrospective ¹	September 2011 to June 2017	7	-	15.0 (9-18)	30.7 (1-84)	14 (7-24)	-	-	39.6 (18-54)	-
Choné <i>et al</i> [<mark>32</mark>], 2019	France	Retrospective	January 2012 to August 2018	117	69:48	14.2 (3.7)	21.3 (21.2)	8.3 (3-21)	72.5 (16-240)	3.9 (1-14)	18 (3.3-53.7)	-
Nabi <i>et al</i> [33] , 2019	India	Retrospective	September 2013 to Jan 2018	44	22:21	14.58 ± 3.41	24 (2-96)	10.09 (5-15)	65.46 (18-240)	3 (2-4)	18 (1-53.1)	Symptoms, EGD and 24-h pH- impedance
Liu et al[<mark>34</mark>], 2019	China	Retrospective	August 2010 to	130	82:48	-	12 (0-13)	7.2 ± 1.4	30 (15-255)	3 (1-21)	40 (4-88)	Barium swallow, EGD and HRM

Pu XX et al. POEM in pediatric and geriatric patients

			August 2017									
Saez et al[<mark>35</mark>], 2020	Chile	Retrospective	March 2017 to November 2019	5	4:1	11 (5-15)	-	9.8 (9-11)	70 (50-120)	2 (1-3)	20.5 (4 -37)	EGD and HRM
Wood <i>et al</i> [<mark>36</mark>], 2020	United States	Prospective	2014 to 2019.	21	14:7	13 (2-17)	-	7±1.1	92 ± 52	1 ± 0.5	12	-
Peng <i>et al</i> [37], 2022	China	Retrospective	October 2011 to November 2016	24	14:10	14.42 ± 2.65	14.5 (3-84)	9 (5-10)	58.67 ± 19.10	6.42 ± 2.15	85.75 ± 25.91	Gerd Q score and EGD
Petrosyan <i>et al</i> [38], 2022	United States	Retrospective	July 2015 to September 2021	37	23:14	11.6 ± 4.5	-	6.5 ± 0.93	138.1 ± 62.2	2.4 ± 0.9	22.6 ± 20	-

¹Published conference abstracts.

M: Male; F: Female; POEM: Peroral endoscopic myotomy; pH: Potential of hydrogen; EGD: Esophagogastroduodenoscopy; HRM: High resolution esophageal manometry; 99mTc DTPA: Diethylenetriaminepentaacetic acid.

and 55.4% of the patients were male. The duration of symptoms ranged from 12 to 26.4 months. The mean myotomy length and operation time was mentioned in 18 and 17 studies, which ranged from 6.5 to 14 cm and 30 to 142 minutes, respectively. Furthermore, the length of hospital stay and follow-up time ranged from 1 to 9 days and from 6 to 85.75 months, respectively.

Clinical outcomes: In total, 20 studies (Table 2) reported the technical success and clinical success rates of POEM for achalasia in pediatric patients; The pooled rates were 97.1% (95%CI: 95.0%-98.3%; $l^2 = 0\%$; P < 0.000) and 93.2% (95%CI: 90.5%-95.2%, $l^2 = 0\%$; P < 0.000], respectively (Figure 2A and B). On the other hand, the pooled rates of GERD and adverse events were 22.3% (95%CI: 18.4%-26.7%; $l^2 = 43.874\%$; P < 0.000) and 20.4% (95%CI: 16.6%-24.8%; $l^2 = 67.217\%$; P < 0.000), respectively (Figure 2C and D). It must be noted that the definition of adverse events in each study was inconsistent, with some authors defining adverse events as those requiring intervention or major adverse events[33,34], whereas others defined gas-related complications without clinical symptoms as adverse events[19,23,27].

Role of POEM for achalasia in geriatric patients

Baseline characteristics: There were eight original studies and four conference abstracts on the efficacy and safety of POEM in geriatric patients, and the results are shown in Table 3[13,39-49]. Among these articles, four studies were conducted in Japan, three in the United States, three in China, one in Italy, and one in the Netherlands. All studies were retrospective cohort studies, and were conducted from September 2008 to May 2021. A total of 810 geriatric patients aged 67.9 to 84 years (aged \geq 80 years in one study) were included in this study, and 50% of the patients were male. The duration of symptoms, reported in nine studies, ranged from 4.4 to 30 years. Myotomy length was mentioned in 10 studies, which ranged from 10 to 14.5 cm. In addition, the operation duration of POEM, reported in 8 studies, ranged from 46.87 to 138.3 minutes. Moreover, the duration of hospital stay and the follow-up time ranged from 1 to 9.8 years and from 2 to 41 months, respectively.

Clinical outcomes: Overall, the 12 studies (Table 4) had pooled rates of technical success, clinical success, GERD and adverse events of 97.7% (95%CI: 95.8%-98.7%; *I*² = 15.200%; *P* < 0.000), 93.2% (95%CI: 90.3%-95.2%; *I*² = 0%, *P* < 0.000), 23.9% (95%CI: 19.4%-29.1%; *I*² = 75.697%; *P* < 0.000), and 10.8% (95%CI: 8.3%-14.0%; *I*² = 62.938%; *P* < 0.000), respectively (Figure 3).

Table 2 Clinical outcomes of the studies on peroral endoscopic myotomy for achalasia in pediatric patients, <i>n</i> (%)											
Ref.	Technical success	Clinical success	GERD	Adverse events							
Li et al[19], 2015	9/9 (100)	9/9 (100)	1/9 (11.1)	2/9 (11.1)							
Chen <i>et al</i> [20], 2015	26/27 (96.3)	26/26 (100)	5/26 (19.2)	9/26 (34.6) (Cumulative adverse events on CT scan: 53)							
Caldaro <i>et al</i> [21], 2015	9/9 (100)	9/9 (100)	1/9 (11.1)	1/9 (11.1)							
Tang <i>et al</i> [22], 2015	5/5 (100)	4/4 (100)	0 (0)	0 (0)							
Tan <i>et al</i> [23], 2016	12/12 (100)	12/12 (100)	2/12 (16.7)	3/12 (8.3)							
Stavropoulos et al[24], 2017	10/10 (100)	10/10 (100)	2/5 (40)	4/10 (40)							
Zangen <i>et al</i> [25], 2017	5/5 (100)	5/5 (100)	0 (0)	0 (0)							
Kethman <i>et al</i> [26], 2018	10/10 (100)	8/10 (80)	0 (0)	3/10 (30)							
Miao <i>et al</i> [27], 2018	21/21 (100)	21/21 (100)	6/21 (28.6)	12/21 (57.1)							
Korrapati <i>et al</i> [28], 2018	15/15 (100)	15/15 (100)	0 (0)	2/15 (13.3)							
Nishimoto et al[29], 2018	13/13 (100)	12/13 (92.3)	-	2/13 (15.4)							
Mangiola <i>et al</i> [30], 2018	25/26 (96.2)	26/26 (100)	4/17 (23.5)	6/26 (23.1)							
Yamashita <i>et al</i> [<mark>31</mark>], 2018	7/7 (100)	7/7 (100)	-	0 (0)							
Choné <i>et al</i> [32], 2019	116/117 (99.1)	106/117 (90.6)	17/117 (21.4)	7/117 (8.5)							
Nabi <i>et al</i> [33], 2019	43/44 (97.7)	40/44 (90.9)	11/20 (55)	11/43 (25.6)							
Liu et al[<mark>34</mark>], 2019	129/130 (99.2)	108/113 (95.6)	30/111 (27.0)	5/130 (3.8)							
Saez et al[35], 2020	5/5 (100)	5/5 (100)	2/5 (40)	0 (0)							
Wood <i>et al</i> [36], 2020	21/21 (100)	21/21 (100)	-	6/21 (28.6)							
Peng et al[37], 2022	24/24 (100)	23/24 (95.8)	5/21 (23.8)	0 (0)							
Petrosyan <i>et al</i> [38], 2022	37/37 (100)	37/37 (100)	3/37 (8.1)	9/37 (24.3)							

GERD: Gastroesophageal reflux disease; CT: Computed tomography.

Publication Bias: Most of the funnel plots were relatively symmetric, suggesting that publication bias was not significant (Figure 4, Figure 5A and B). However, funnel plots regarding GERD and adverse events in geriatric patients displayed substantial asymmetry (Figure 5C and D).

DISCUSSION

Achalasia is a relatively uncommon disorder of the esophageal smooth muscle, and the annual incidence and prevalence increases with age, but it can affect all ages[1]. Due to the short length and weak wall of the esophagus in pediatric patients, and esophagus modifications contributing to a corkscrew esophagus, increased comorbidities and poor treatment tolerance in geriatric patients [43,49-52], the difficulty of surgery and the incidence of postoperative adverse events in these populations are increased.

Currently, the purpose of all treatment methods for achalasia is to reduce the hypertonicity of the LES to attain the goals of reducing symptoms, improving esophageal emptying, and avoiding further dilation of the esophagus[3]. Pharmacologic, botulinum toxin injection, PD, POEM, and LHM are currently available therapeutic approaches. Specifically, pharmacologic therapy is the least effective treatment for achalasia, resulting in a short-term reduction in LES pressure in 13%-65% of patients and symptom relief in 0%-87% of patients[8]. The effect of botulinum toxin is limited as it is not long-lasting and requires repeated therapy[9]. PD is an effective choice for patients with achalasia, and reports suggest that 50%-93% of patients might achieve symptom relief[1]. However, a recent meta-analysis demonstrated that the long-term efficacy of PD was inferior to that of POEM[10]. LHM is generally regarded as the gold standard because it can provide long-lasting symptom relief; however, for geriatric patients and patients with multiple comorbidities, it is not appropriate as it is an invasive procedure and can result in operative complications[11-13].

The minimally invasive technique POEM was first described by Inoue et al[15]. This technique is used for treating achalasia in adults, and is becoming increasingly available in pediatric and geriatric patients. The clinical guidelines established by the ASGE in the Preservation and Incorporation of Valuable Endoscopic Innovations (PIVI) paper[3] showed that POEM for achalasia treatment is considered a viable therapeutic modality and should fulfil the following criteria: (1) \ge 80% efficacy at 12 months after the procedure (Eckardt score \le 3 with a dysphagia component of \le 2); and (2)

Baishidena® WJGE | https://www.wjgnet.com

Table 3 Baseline characteristics of the studies on p	eroral endoscopic myotom	y for achalasia in geriatric patients
--	--------------------------	---------------------------------------

Ref.	Country	Study type	Study interval	Patients (<i>n</i>)	Gender (M:F)	Age (yr)	Duration of symptoms (month)	Myotomy length (cm)	Operation time (min)	Hospital stay (day)	Follow-up (month)	Method of GERD diagnosis
Wang et al <mark>[39]</mark> , 2016	China	Retrospective	January 2010 to December 2015	21	12:9	67.9 ± 4.3	13.9 ± 11.7	-	-	-	21.8	EGD, esophageal manometry and barium esophagram
Chen <i>et al</i> [<mark>13</mark>], 2018	United States	Retrospective	January 2010 to January 2016	76	40:36	84 ± 3.2	24.0 (17.8-30.3)	10.0 ± 4.0	103.7 ± 47.9	3	8.5 (IQR: 2.2- 18.2)	-
Landi <i>et al</i> [40], 2018	Italy	Retrospective ¹	May 2011 to April 2017	88	39:49	72.2 ± 4.7	-	12.3 ± 3	-	-	24	-
Liu <i>et al</i> [<mark>41</mark>], 2019	China	Retrospective	August 2010 to December 2017	139	65:74	70.22 ± 5.68	8 (IQR: 2-20)	10.57 ± 1.81	50 (IQR: 36-76)	3 (IQR: 2-4)	41 (IQR: 26- 60)	Barium swallow, EGD and HRM
Klair <i>et al</i> [<mark>42</mark>], 2019	United States	Retrospective ¹	December 2014 to October 2018	62	36:26	72.3 ± 5.7	-	13.8	-	1.9	-	-
Abe <i>et al</i> [<mark>43</mark>], 2020	Japan	Retrospective	April 2015 to March 2019	28	12:16	≥80	5.5 (0.25-59)	14.5 (4-26)	60.5 (36-124)	9.8 (4-51)	-	EGD and HRM
Sanaka <i>et al</i> [44], 2020	United States	Retrospective	April 2014 to May 2019	55	31:24	74 (70-79)	30 (12-60)	10 (IQR: 8-10)	90 (IQR: 7.5- 110)	1 (IQR: 1-2)	2.4 (IQR: 2.2- 2.7)	HRM, timed barium esophagram and 24-h esophageal pH study
Angeli Abad et al[45], 2020	Netherlands	Retrospective ¹	September 2008 to June 2019	66	28:38	83 (80-92)	10.5 (0.2-62.4)	-	-	7 ± 4.1	12	-
Okada <i>et al</i> [<mark>46</mark>], 2021	Japan	Retrospective	September 2011 to March 2020	100	40:60	74.2 (65-93)	10.3 ± 13.6	12.5 (3-25)	138.3 (50-460)	-	36	-
Nakamura <i>et al</i> [<mark>47</mark>], 2021	Japan	Retrospective	August 2014 to May 2021	11	7:4	81 (75-87)	5 (2-40)	13 (8-19)	109 (62-144)	-	36	EGD, esophagography and HRM
Ujiie <i>et al</i> [<mark>48</mark>], 2021	Japan	Retrospective	January 2015 to December 2019	18	12:6	78 (75-86)	4.4 (0.05-50.2)	10 (5-16)	104 (45-165)	-	2	IRP, HRM and EGD
Zhao <i>et al</i> [<mark>49</mark>], 2022	China	Retrospective ¹	November 2010 to September 2019	146	-	-	-	7.09 ± 2.49	46.87 ± 19.29	-	-	GerdQ score

¹Published conference abstracts.

M: Male; F: Female; EGD: Esophagogastroduodenoscopy; HRM: High resolution esophageal manometry; IRP: Integrated relaxation pressure; IQR: Interquartile range; pH: Potential of hydrogen.

Serious adverse event rate $\leq 6\%$ and mortality rate $\leq 0.1\%$ within 30 days after the procedure.

Previous investigations have demonstrated that POEM can provide short-term benefits in pediatric patients, with a median follow-up period of approximately 13.2 to 40 months. Given that children have a longer life expectancy, the long-term therapeutic outcomes for this patient group are especially significant[37]. The largest series of POEM for children with achalasia was conducted in China by Liu *et al*[34] published in 2019. The authors retrospectively evaluated a total of 130 pediatric patients. The technical success rate for POEM in these patients was 99.2%, and the clinical success rates at 1,

Table 4 Results of the studies on peroral endoscopic myotomy for achalasia in geriatric patients, <i>n</i> (%)										
Ref.	Technical success	Clinical success	GERD	Adverse events						
Wang et al[39], 2016	21/21 (100)	20/21 (95.2)	2/21 (9.5)	1/21 (4.8)						
Chen <i>et al</i> [13], 2018	71/76 (93.4)	59/65 (90.8)	13/76 (16.1)	11/76 (14.5)						
Landi <i>et al</i> [40], 2018	88/88 (100)	84/88 (95.4)	-	-						
Liu et al[41], 2019	138/139 (99.3)	79/85 (92.9)	20/85 (23.53)	4/139 (2.88)						
Klair <i>et al</i> [42], 2019	62/62 (100)	55/62 (88.7)	-	5/62 (8.1)						
Abe <i>et al</i> [43], 2020	28/28 (100)	17/17 (100)	3/23 (13.0)	8/28 (28.6)						
Sanaka et al[44], 2020	55/55 (100)	36/38 (94.7)	6/51(11.8)	3/55 (5.5)						
Angeli Abad et al[45], 2020	66/66 (100)	19/20 (95.0)	-	5/66 (7.6)						
Okada et al[46], 2021	100/100 (100)	92/92 (100)	-	11/100 (11)						
Nakamura <i>et al</i> [47], 2021	11/11 (100)	11/11 (100)	1/11 (9)	3/11 (27.3)						
Ujiie et al[48], 2021	18/18 (100)	18/18 (100)	1/18 (5.6)	0 (0)						
Zhao et al[49], 2022	146/146 (100)	(96.33)	(15.60)	-						

GERD: Gastroesophageal reflux disease.



Figure 1 Flow diagram showing study selection. POEM: Peroral endoscopic myotomy.

3, and 5 years were 98.2%, 96.5%, and 95.6%, respectively. In our study, the pooled technical and clinical success rates for pediatric patients were 97.1% (95%CI: 95.0%-98.3%; P = 0%; P < 0.000) and 93.2% (95%CI: 90.5%-95.2%; P = 0%, P < 0.000), and the follow-up time ranged from 6 to 85.75 months. These findings build on the outcomes of earlier research and offer further support that POEM is a highly effective therapeutic approach for childhood achalasia.

Saishideng® WJGE | https://www.wjgnet.com

A

Study name Statistics for each study							Event	rate and	95%CI	
	Event rate	Lower limit	Upper limit	Z-value	P-value					
Li et al, 2015	0.950	0.525	0.997	2.029	0.042	1	1			- I
Chen et al, 2015	0.963	0.779	0.995	3.197	0.001					
Caldaro et al, 2015	0.950	0.525	0.997	2.029	0.042					
Tang et al, 2015	0.917	0.378	0.995	1.623	0.105			_ ·	-	
Tan et al, 2016	0.962	0.597	0.998	2.232	0.026					
Stavropoulos et al, 2017	0.955	0.552	0.997	2.103	0.035				-	
Zangen et al, 2017	0.917	0.378	0.995	1.623	0.105			- 1 -	-	
Kethman et al, 2018	0.955	0.552	0.997	2.103	0.035				-	
Miao et al, 2018	0.977	0.723	0.999	2.629	0.009					
Korrapati et al, 2018	0.969	0.650	0.998	2.390	0.017					
Nishimoto et al, 2018	0.964	0.616	0.998	2.289	0.022					
Mangiola et al, 2018	0.962	0.772	0.995	3.156	0.002				-	
Yamashita et al, 2018	0.938	0.461	0.996	1.854	0.064				-	
Choné et al, 2019	0.991	0.942	0.999	4.733	0.000					
Nabi et al, 2019	0.977	0.856	0.997	3.718	0.000					
Liu et al, 2019	0.992	0.947	0.999	4.841	0.000					
Saez et al, 2020	0.917	0.378	0.995	1.623	0.105			- I -	-	
Wood et al, 2020	0.977	0.723	0.999	2.629	0.009					
Peng et al, 2022	0.980	0.749	0.999	2.724	0.006				- 4	
Petrosyan et al, 2022	0.987	0.822	0.999	3.033	0.002					
-	0.971	0.950	0.983	12.169	0.000					
						-2.00	-1.00	0.00	1.00	2.00

В

Clinical success

Study name		Statisti	cs for ea	ch study			Event rate and 95%CI				
	Event rate	Lower limit	Upper limit	Z-value	P-value						
Li et al, 2015	0.950	0.525	0.997	2.029	0.042	1				1	
Chen et al, 2015	0.981	0.764	0.999	2.781	0.005						
Caldaro et al, 2015	0.950	0.525	0.997	2.029	0.042				-		
Tang et al, 2015	0.900	0.326	0.994	1.474	0.140			<u> </u>	-		
Tan et al, 2016	0.962	0.597	0.998	2.232	0.026						
Stavropoulos et al, 2017	0.955	0.552	0.997	2.103	0.035				-		
Zangen et al, 2017	0.917	0.378	0.995	1.623	0.105			- 1 -	-		
Kethman et al, 2018	0.800	0.459	0.950	1.754	0.080						
Miao et al, 2018	0.977	0.723	0.999	2.629	0.009				-		
Korrapati et al, 2018	0.969	0.650	0.998	2.390	0.017						
Nishimoto et al, 2018	0.923	0.609	0.989	2.387	0.017						
Mangiola et al, 2018	0.981	0.764	0.999	2.781	0.005						
Yamashita et al, 2018	0.938	0.461	0.996	1.854	0.064				-		
Choné et al, 2019	0.906	0.838	0.947	7.152	0.000						
Nabi et al, 2019	0.909	0.782	0.965	4.391	0.000						
Liu et al, 2019	0.956	0.898	0.981	6.717	0.000						
Saez et al, 2020	0.917	0.378	0.995	1.623	0.105				-		
Wood et al, 2020	0.977	0.723	0.999	2.629	0.009						
Peng et al, 2022	0.958	0.756	0.994	3.069	0.002						
Petrosyan et al, 2022	0.987	0.822	0.999	3.033	0.002						
	0.932	0.905	0.952	13.996	0.000						
						-2.00	-1.00	0.00	1.00	2.00	

C GERD

GERD											
Study name		Statisti	cs for eac	h study			Event rate and 95%CI				
	Event rate	Lower limit	Upper limit	Z-value	P-value						
Li et al, 2015	0.111	0.015	0.500	-1.961	0.050		1		·		
Chen et al, 2015	0.192	0.082	0.387	-2.884	0.004			- H-			
Caldaro et al, 2015	0.111	0.015	0.500	-1.961	0.050			-			
Tang et al, 2015	0.083	0.005	0.622	-1.623	0.105			_ _	-		
Tan et al, 2016	0.167	0.042	0.477	-2.078	0.038			_ ₩			
Stavropoulos et al, 2017	0.400	0.100	0.800	-0.444	0.657				_		
Zangen et al, 2017	0.083	0.005	0.622	-1.623	0.105				-		
Kethman et al, 2018	0.045	0.003	0.448	-2.103	0.035			- -			
Miao et al, 2018	0.286	0.134	0.508	-1.897	0.058			- I 			
Korrapati et al, 2018	0.031	0.002	0.350	-2.390	0.017			_ ⊨			
Mangiola et al, 2018	0.235	0.091	0.486	-2.061	0.039						
Choné et al, 2019	0.145	0.092	0.221	-6.754	0.000						
Nabi et al, 2019	0.550	0.336	0.747	0.446	0.655			⊣	-		
Liu et al, 2019	0.270	0.196	0.360	-4.647	0.000						
Saez et al, 2020	0.400	0.100	0.800	-0.444	0.657				- 1		
Peng et al, 2022	0.238	0.103	0.460	-2.270	0.023			_ 			
Petrosyan et al, 2022	0.081	0.026	0.223	-4.031	0.000			•			
	0.223	0.184	0.267	-10.212	0.000			•			
						-2.00	-1.00	0.00	1.00	2.00	



	-	
	-	ł
	- 4	

Adverse events

Study name		Statisti	cs for ea	ch study			Event rate and 95%CI					
	Event rate	Lower limit	Upper limit	Z-value	P-value							
Li et al, 2015	0.222	0.056	0.579	-1.562	0.118	1	1	- ≡	· 1	- I		
Chen et al, 2015	0.346	0.191	0.543	-1.543	0.123			- H-	-			
Caldaro et al, 2015	0.111	0.015	0.500	-1.961	0.050							
Tang et al, 2015	0.083	0.005	0.622	-1.623	0.105				-			
Tan et al, 2016	0.250	0.083	0.552	-1.648	0.099				.			
Stavropoulos et al, 2017	0.400	0.158	0.703	-0.628	0.530			_ _	-			
Zangen et al, 2017	0.083	0.005	0.622	-1.623	0.105			-	-			
Kethman et al, 2018	0.300	0.100	0.624	-1.228	0.220				-			
Miao et al, 2018	0.571	0.360	0.760	0.652	0.514			_ _ _	-			
Korrapati et al, 2018	0.133	0.034	0.405	-2.464	0.014							
Nishimoto et al, 2018	0.154	0.039	0.451	-2.218	0.027							
Mangiola et al, 2018	0.231	0.108	0.428	-2.587	0.010			- H-				
Yamashita et al, 2018	0.063	0.004	0.539	-1.854	0.064			_ 				
Choné et al, 2019	0.060	0.029	0.120	-7.067	0.000							
Nabi et al, 2019	0.256	0.148	0.405	-3.055	0.002			- I -				
Liu et al, 2019	0.038	0.016	0.089	-7.058	0.000							
Saez et al, 2020	0.083	0.005	0.622	-1.623	0.105				-			
Wood et al, 2020	0.286	0.134	0.508	-1.897	0.058			- -				
Peng et al, 2022	0.020	0.001	0.251	-2.724	0.006			_ i ⊷ _				
Petrosyan et al, 2022	0.243	0.132	0.405	-2.962	0.003							
	0.204	0.166	0.248	-10.661	0.000			•				
						-2.00	-1.00	0.00	1.00	2.00		



Most cases of achalasia occur in individuals between the ages of 30 and 50 years, although it is not uncommon for older adults to be diagnosed with the condition. The risk of aspiration pneumonia, a potentially lethal complication of regurgitation in achalasia, is particularly high in the elderly, highlighting the significance of exploring effective clinical interventions for this patient population[46]. To date, many studies have reported the results of POEM in elderly patients [39-49]. Zhong *et al*[52] published a meta-analysis that included seven studies involving 469 geriatric patients. They reported that the pooled technical success rate was 98.1%, and the pooled clinical success rate was 92.5%. We conducted a more comprehensive study and found that the pooled technical and clinical success rates for geriatric patients were 97.7% (95%CI: 95.8%-98.7%; $I^2 = 15.200\%$; P < 0.000) and 93.2% (95%CI: 90.3%-95.2%; $I^2 = 0\%$, P < 0.000), respectively.

POEM can potentially enhance the quality of life in pediatric patients as they mature. Nevertheless, it is important to be aware of the potential adverse events following POEM, which may include GERD, pneumoperitoneum, pneumothorax, pneumonitis, mucosal injury, subcutaneous emphysema and mediastinal emphysema[53]. Our study revealed that the pooled adverse event rate following POEM in pediatric achalasia was 20.4% (95%CI: 16.6%-24.8%; $I^2 = 67.217\%$; P < 0.000), and the pooled GERD rate was 22.3% (95%CI: 18.4%-26.7%; $I^2 = 43.874\%$; P < 0.000). Chen *et al*[20] reported cumulative adverse events of 53 on the basis of computed tomography scans, which were related to gas. They revealed that this high incidence was due to air for insufflation in 20 patients (76.9%). They also demonstrated that air insufflation had a higher rate of postoperative gas-related adverse events than carbon dioxide insufflation [84.6% (11/13) *vs* 16.7% (1/6)][22]. In addition, Lee *et al*[54] included 12 studies involving 146 pediatric patients who underwent POEM. The authors found that at least 93% of the children experienced an improvement in achalasia symptoms after POEM, with a limited number of patients reporting minor adverse effects that could be controlled conservatively. Recently, Zhong *et al*[55] published an updated meta-analysis involving a total of 11 studies with 389 children. The pooled major adverse event rate was reported to be 12.8%, while the pooled GERD rate was 17.8%. Two meta-analyses suggested that POEM was effective and safe for treating achalasia in pediatric patients[54,55].

In geriatric patients, the pooled adverse events and GERD rates following POEM for achalasia were 10.8% (95%CI: 8.3%-14.0%; $l^2 = 62.938\%$; P < 0.000) and 23.9% (95%CI: 19.4%-29.1%; $l^2 = 75.697\%$; P < 0.000], respectively. Furthermore, Zhong *et al*[55] published a meta-analysis including 7 studies involving 469 geriatric patients, and the major adverse event rate was 9.0%, and the clinical reflux rate was 17.4%. In total, almost all these studies indicated that POEM is a safe and effective treatment for geriatric patients with esophageal achalasia[56].

There were several limitations in our analyses. First, the results of the NIH quality assessment revealed that one of our included studies was of poor quality, which may have reduced the evidence quality in our article. In addition, our study lacked several characteristic indicators, including achalasia type, American Society of Anesthesiology physical status classification system score, type of myotomy, pre- and postoperative Eckardt score, and LES pressure; thus, the analysis was not comprehensive. In addition, the study examining GERD and adverse events in geriatric patients exhibited publication bias; therefore, future investigations should include subgroup analyses to determine the underlying reasons for this. Furthermore, as the symptoms of GERD are very similar to those of achalasia, possible confusion could have occurred during interpretation of the results. A large proportion of cases with long-term follow-up could not be objectively evaluated for GERD, and the possibility of selection bias could not be completely ruled out. Both may have led to a higher pooled rate of GERD. Lastly, almost all of the studies were small sample, nonrandomized and observational studies, which may have been subject to a range of biases.

A

Technical	success
	0400000

Study name		Statisti	rs for eac	h study	Event rate and 95%CI						
Study hame	Event rate	Lower	Upper limit	<i>Z</i> -value	P-value		Event		55 /001		
Wang et al, 2016	0.977	0.723	0.999	2.629	0.009						
Chen et al, 2018	0.934	0.851	0.972	5.734	0.000						
Landi et al, 2018	0.994	0.917	1.000	3.650	0.000						
Liu et al, 2019	0.993	0.951	0.999	4.909	0.000				•		
Klair et al, 2019	0.992	0.885	1.000	3.401	0.001				- ÷ -		
Abe et al, 2020	0.983	0.777	0.999	2.834	0.005						
Sanaka et al, 2020	0.991	0.873	0.999	3.315	0.001						
Abad et al, 2020	0.993	0.892	1.000	3.445	0.001				- • ·		
Okada et al, 2021	0.995	0.926	1.000	3.741	0.000				- •		
Nakamura et al, 2021	0.958	0.575	0.997	2.170	0.030						
Ujiie et al, 2021	0.974	0.690	0.998	2.519	0.012						
Zhao et al, 2022	0.997	0.948	1.000	4.010	0.000						
	0.977	0.958	0.987	12.159	0.000						
						-2.00	-1.00	0.00	1.00	2.00	

В

Clinical success												
Study name		Statisti	cs for ea	ch study		Event rate and 95%CI						
	Event rate	Lower limit	Upper limit	Z-value	P-value							
Wang et al, 2016	0.952	0.729	0.993	2.924	0.003	- I	1			1		
Chen et al, 2018	0.908	0.809	0.958	5.334	0.000							
Landi et al, 2018	0.955	0.885	0.983	5.949	0.000							
Liu et al, 2019	0.929	0.852	0.968	6.087	0.000							
Klair et al, 2019	0.887	0.782	0.945	5.137	0.000				-			
Abe et al, 2020	0.972	0.678	0.998	2.479	0.013							
Sanaka et al, 2020	0.947	0.813	0.987	3.979	0.000				-			
Abad et al, 2020	0.950	0.718	0.993	2.870	0.004				-			
Okada et al, 2021	0.995	0.920	1.000	3.681	0.000				•			
Nakamura et al, 2021	0.958	0.575	0.997	2.170	0.030							
Ujiie et al, 2021	0.974	0.690	0.998	2.519	0.012							
	0.932	0.903	0.952	13.515	0.000							
						-2.00	-1.00	0.00	1.00	2.00		

C GERD

Study name		Statisti	cs for eac	ch study	Event rate and 95%CI					
	Event rate	Lower limit	Upper limit	Z-value	P-value					
Wang et al, 2016	0.095	0.024	0.311	-3.028	0.002		1	-		- I
Chen et al, 2018	0.171	0.102	0.273	-5.181	0.000					
Liu et al, 2019	0.235	0.157	0.337	-4.609	0.000			- E		
Abe et al, 2020	0.130	0.043	0.335	-3.064	0.002					
Sanaka et al, 2020	0.118	0.054	0.238	-4.636	0.000			-		
Abad et al, 2020	0.455	0.339	0.575	-0.738	0.461				•	
Nakamura et al, 2021	0.091	0.013	0.439	-2.195	0.028			- II-		
Ujiie et al, 2021	0.056	0.008	0.307	-2.753	0.006			_ ⊨		
	0.239	0.194	0.291	-8.502	0.000			+		
						-2.00	-1.00	0.00	1.00	2.00

D

Adverse events												
Study name		Statisti	cs for ea	ch study		Event rate and 95%CI						
	Event rate	Lower limit	Upper limit	Z-value	P-value							
Wang et al, 2016	0.048	0.007	0.271	-2.924	0.003	1	1	- H -				
Chen et al, 2018	0.145	0.082	0.243	-5.449	0.000			-				
Liu et al, 2019	0.029	0.011	0.074	-6.936	0.000							
Klair et al, 2019	0.081	0.034	0.180	-5.218	0.000							
Abe et al, 2020	0.286	0.150	0.476	-2.190	0.028			- I 				
Sanaka et al, 2020	0.055	0.018	0.156	-4.804	0.000			- F				
Abad et al, 2020	0.076	0.032	0.169	-5.377	0.000							
Okada et al, 2021	0.110	0.062	0.188	-6.542	0.000							
Nakamura et al, 2021	0.273	0.090	0.586	-1.449	0.147			- -	-			
Ujiie et al, 2021	0.026	0.002	0.310	-2.519	0.012			_ ∔				
	0.108	0.083	0.140	-14.047	0.000			•				
						-2.00	-1.00	0.00	1.00	2.00		

Figure 3 Forest plot of peroral endoscopic myotomy for achalasia in geriatric patients. A: Technical success; B: Clinical success; C: Gastroesophageal reflux disease; D: Adverse events. GERD: Gastroesophageal reflux disease.



Baisbideng® WJGE | https://www.wjgnet.com



Figure 4 Assessment of publication bias in pediatric patients. A: Technical success; B: Clinical success; C: Gastroesophageal reflux disease; D: Adverse events.



Figure 5 Assessment of publication bias in geriatric patients. A: Technical success; B: Clinical success; C: Gastroesophageal reflux disease; D: Adverse events.

WJGE | https://www.wjgnet.com

Paishideng⁴

CONCLUSION

Currently, on the basis of the available published evidence, POEM has been shown to be an effective and safe therapy for achalasia in both pediatric and geriatric patients. Nevertheless, there is a need for additional high-quality randomized controlled trials to establish the optimal treatment approach for achalasia within these specific populations.

FOOTNOTES

Author contributions: Tang XW and Pu XX contributed to study conception and design; Pu XX, Huang S and Zhong CY contributed to drafting of the manuscript; Wang X, Fu SF, Lv YQ, Zou K, Lü MH contributed to acquisition of data and critical revision; Tang XW and Peng Y contributed to revision of the manuscript, and final approval of the manuscript; All authors have read and approved the final manuscript.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country of origin: China

ORCID number: Shu Huang 0000-0001-7284-469X; Chun-Yu Zhong 0000-0002-6934-3795; Mu-Han Lü 0000-0002-0527-3695; Yan Peng 0000-0002-4007-0850; Xiao-Wei Tang 0000-0001-6064-0526.

S-Editor: Fan M L-Editor: Webster IR P-Editor: Zhao YQ

REFERENCES

- Savarino E, Bhatia S, Roman S, Sifrim D, Tack J, Thompson SK, Gyawali CP. Achalasia. Nat Rev Dis Primers 2022; 8: 28 [PMID: 35513420 1 DOI: 10.1038/s41572-022-00356-8]
- 2 Zaninotto G, Bennett C, Boeckxstaens G, Costantini M, Ferguson MK, Pandolfino JE, Patti MG, Ribeiro U Jr, Richter J, Swanstrom L, Tack J, Triadafilopoulos G, Markar SR, Salvador R, Faccio L, Andreollo NA, Cecconello I, Costamagna G, da Rocha JRM, Hungness ES, Fisichella PM, Fuchs KH, Gockel I, Gurski R, Gyawali CP, Herbella FAM, Holloway RH, Hongo M, Jobe BA, Kahrilas PJ, Katzka DA, Dua KS, Liu D, Moonen A, Nasi A, Pasricha PJ, Penagini R, Perretta S, Sallum RAA, Sarnelli G, Savarino E, Schlottmann F, Sifrim D, Soper N, Tatum RP, Vaezi MF, van Herwaarden-Lindeboom M, Vanuytsel T, Vela MF, Watson DI, Zerbib F, Gittens S, Pontillo C, Vermigli S, Inama D, Low DE. The 2018 ISDE achalasia guidelines. Dis Esophagus 2018; 31 [PMID: 30169645 DOI: 10.1093/dote/doy071]
- Vaezi MF, Pandolfino JE, Yadlapati RH, Greer KB, Kavitt RT. ACG Clinical Guidelines: Diagnosis and Management of Achalasia. Am J 3 Gastroenterol 2020; 115: 1393-1411 [PMID: 32773454 DOI: 10.14309/ajg.000000000000731]
- 4 Marlais M, Fishman JR, Fell JM, Haddad MJ, Rawat DJ. UK incidence of achalasia: an 11-year national epidemiological study. Arch Dis Child 2011; 96: 192-194 [PMID: 20515971 DOI: 10.1136/adc.2009.171975]
- Mayberry JF, Mayell MJ. Epidemiological study of achalasia in children. Gut 1988; 29: 90-93 [PMID: 3343019 DOI: 10.1136/gut.29.1.90] 5
- Sato H, Yokomichi H, Takahashi K, Tominaga K, Mizusawa T, Kimura N, Kawata Y, Terai S. Epidemiological analysis of achalasia in Japan 6 using a large-scale claims database. J Gastroenterol 2019; 54: 621-627 [PMID: 30607612 DOI: 10.1007/s00535-018-01544-8]
- 7 Li CJ, Tan YY, Wang XH, Liu DL. Peroral endoscopic myotomy for achalasia in patients aged \geq 65 years. World J Gastroenterol 2015; 21: 9175-9181 [PMID: 26290645 DOI: 10.3748/wjg.v21.i30.9175]
- Vaezi MF, Richter JE. Current therapies for achalasia: comparison and efficacy. J Clin Gastroenterol 1998; 27: 21-35 [PMID: 9706766 DOI: 8 10.1097/00004836-199807000-00006
- 9 Campos GM, Vittinghoff E, Rabl C, Takata M, Gadenstätter M, Lin F, Ciovica R. Endoscopic and surgical treatments for achalasia: a systematic review and meta-analysis. Ann Surg 2009; 249: 45-57 [PMID: 19106675 DOI: 10.1097/SLA.0b013e31818e43ab]
- Zhong C, Tan S, Huang S, Lü M, Peng Y, Fu X, Tang X. Peroral endoscopic myotomy versus pneumatic dilation for achalasia: a systematic 10 review and meta-analysis. Eur J Gastroenterol Hepatol 2020; 32: 1413-1421 [PMID: 32516175 DOI: 10.1097/MEG.000000000001800]
- Evensen H, Kristensen V, Larssen L, Sandstad O, Hauge T, Medhus AW. Outcome of peroral endoscopic myotomy (POEM) in treatment-11 naive patients. A systematic review. Scand J Gastroenterol 2019; 54: 1-7 [PMID: 30650313 DOI: 10.1080/00365521.2018.1549271]
- Schlottmann F, Luckett DJ, Fine J, Shaheen NJ, Patti MG. Laparoscopic Heller Myotomy Versus Peroral Endoscopic Myotomy (POEM) for 12 Achalasia: A Systematic Review and Meta-analysis. Ann Surg 2018; 267: 451-460 [PMID: 28549006 DOI: 10.1097/SLA.00000000002311]
- Chen YI, Inoue H, Ujiki M, Draganov PV, Colavita P, Mion F, Romanelli J, Chiu P, Balassone V, Patel L, Abbas A, Yang D, Dunst C, Pioche 13 M, Roman S, Rivory J, Ponchon T, Desilets D, Maselli R, Onimaru M, Nakamura J, Hata Y, Hajiyeva G, Ismail A, Ngamruengphong S, Bukhari M, Chavez YH, Kumbhari V, Repici A, Khashab MA. An international multicenter study evaluating the clinical efficacy and safety of



per-oral endoscopic myotomy in octogenarians. Gastrointest Endosc 2018; 87: 956-961 [PMID: 28235595 DOI: 10.1016/j.gie.2017.02.007]

- Finlayson EV, Birkmeyer JD. Operative mortality with elective surgery in older adults. Eff Clin Pract 2001; 4: 172-177 [PMID: 11525104] 14
- 15 Inoue H, Minami H, Kobayashi Y, Sato Y, Kaga M, Suzuki M, Satodate H, Odaka N, Itoh H, Kudo S. Peroral endoscopic myotomy (POEM) for esophageal achalasia. Endoscopy 2010; 42: 265-271 [PMID: 20354937 DOI: 10.1055/s-0029-1244080]
- Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the 16 PRISMA statement. PLoS Med 2009; 6: e1000097 [PMID: 19621072 DOI: 10.1371/journal.pmed.1000097]
- 17 Cotton PB, Eisen GM, Aabakken L, Baron TH, Hutter MM, Jacobson BC, Mergener K, Nemcek A Jr, Petersen BT, Petrini JL, Pike IM, Rabeneck L, Romagnuolo J, Vargo JJ. A lexicon for endoscopic adverse events: report of an ASGE workshop. Gastrointest Endosc 2010; 71: 446-454 [PMID: 20189503 DOI: 10.1016/j.gie.2009.10.027]
- NIH. Quality Assessment Tool for Before-After Studies with No Control Group. [cited 10 September 2024]. Available from: http://www. 18 nhlbi.nih.gov/health-pro/guidelines/in-develop/cardiovascular-risk-reduction/
- Li C, Tan Y, Wang X, Liu D. Peroral endoscopic myotomy for treatment of achalasia in children and adolescents. J Pediatr Surg 2015; 50: 19 201-205 [PMID: 25598123 DOI: 10.1016/j.jpedsurg.2014.10.017]
- Chen WF, Li QL, Zhou PH, Yao LQ, Xu MD, Zhang YQ, Zhong YS, Ma LL, Qin WZ, Hu JW, Cai MY, He MJ, Cui Z. Long-term outcomes 20 of peroral endoscopic myotomy for achalasia in pediatric patients: a prospective, single-center study. Gastrointest Endosc 2015; 81: 91-100 [PMID: 25088923 DOI: 10.1016/j.gie.2014.06.035]
- 21 Caldaro T, Familiari P, Romeo EF, Gigante G, Marchese M, Contini AC, Federici di Abriola G, Cucchiara S, De Angelis P, Torroni F, Dall'Oglio L, Costamagna G. Treatment of esophageal achalasia in children: Today and tomorrow. J Pediatr Surg 2015; 50: 726-730 [PMID: 25783358 DOI: 10.1016/j.jpedsurg.2015.02.047]
- Tang X, Gong W, Deng Z, Zhou J, Ren Y, Zhang Q, Chen Z, Jiang B. Usefulness of peroral endoscopic myotomy for treating achalasia in 22 children: experience from a single center. Pediatr Surg Int 2015; 31: 633-638 [PMID: 25957132 DOI: 10.1007/s00383-015-3717-9]
- 23 Tan Y, Zhu H, Li C, Chu Y, Huo J, Liu D. Comparison of peroral endoscopic myotomy and endoscopic balloon dilation for primary treatment of pediatric achalasia. J Pediatr Surg 2016; 51: 1613-1618 [PMID: 27339081 DOI: 10.1016/j.jpedsurg.2016.06.008]
- Stavropoulos SN, Sosulski AB, Modayil RJ, Gurram KC, Brathwaite CE, Coren CV, Boinpally H, Grendell JH. Sa2075 Use of Peroral 24 Endoscopic Myotomy (POEM) in Pediatric Patients as a Primary or Rescue Therapy for Achalasia. Gastrointest Endosc 2017; 85: AB285-AB286 [DOI: 10.1016/j.gie.2017.03.646]
- Abstract Supplement for NeuroGASTRO 2017 Congress, 24-26 August 2017, Cork, Ireland. 25 Neurogastroenterol Motil 2017; 29 Suppl 2: 3-140 [PMID: 28804938 DOI: 10.1111/nmo.13180]
- Kethman WC, Thorson CM, Sinclair TJ, Berquist WE, Chao SD, Wall JK. Initial experience with peroral endoscopic myotomy for treatment 26 of achalasia in children. J Pediatr Surg 2018; 53: 1532-1536 [PMID: 28827050 DOI: 10.1016/j.jpedsurg.2017.07.023]
- Miao S, Wu J, Lu J, Wang Y, Tang Z, Zhou Y, Huang Z, Ying H, Zhou P. Peroral Endoscopic Myotomy in Children With Achalasia: A 27 Relatively Long-term Single-center Study. J Pediatr Gastroenterol Nutr 2018; 66: 257-262 [PMID: 28691974 DOI: 10.1097/MPG.000000000001675]
- 28 Korrapati SK, Nemade P, Mahadik M, Biradar V, Pujari R, Bapaye J, Bapaye A. APDW 2018 E-poster Presentations. J Gastroen Hepatol 2018; 33: 54-221 [DOI: 10.1111/jgh.14482]
- Nishimoto M, Shimamura Y, Abad MRAZ, Tokunaga R, Mochizuki Y, Nishikawa Y, Ueno A, Fukuda H, Sumi K, Ikeda H, Onimaru M, 29 Goda K, Ito H, Inoue H. APDW 2018 E-poster Exhibitions – Pediatric. J Gastroen Hepatol 2018; 33: 546-548 [DOI: 10.1111/jgh.14490]
- Mangiola F, Familiari P, Landi R, Cali A, D'aversa F, Bove V, Boskoski I, Tringali A, Perri V, Costamagna G. Tul149 Peroral Endoscopic 30 Myotomy for The Treatment of Achalasia In Children: Experience of A Single Center With Long Term Follow-Up. Gastrointest Endosc 2018; 87: AB542 [DOI: 10.1016/j.gie.2018.04.2179]
- Ebrahim M, Schiodt FV, Willemoe GL, Dahl EE, Christensen E, Rashid S, Holm O, Zander M, Heitmann M, Hansen MB. UEG Week 2018 31 Poster Presentations. UEG J 2018; 6 [DOI: 10.1177/2050640618792819]
- 32 Choné A, Familiari P, von Rahden B, Desai P, Inoue H, Shimamura Y, Eleftheriadis N, Yamashita K, Khashab MA, Shiwaku H, Seewald S, Draganov PV, Alvarez LBM, Chaussade S, Tantau M, Abraham M, Marks J, Arevalo G, Albéniz E, Mion F, Roman S, Rivory J, Dubois R, Lachaux A, Benech N, Subtil F, Ponchon T, Barret M, Pioche M. Multicenter Evaluation of Clinical Efficacy and Safety of Per-oral Endoscopic Myotomy in Children. J Pediatr Gastroenterol Nutr 2019; 69: 523-527 [PMID: 31259787 DOI: 10.1097/MPG.00000000002432
- Nabi Z, Ramchandani M, Chavan R, Darisetty S, Kalapala R, Shava U, Tandan M, Kotla R, Reddy DN. Outcome of peroral endoscopic 33 myotomy in children with achalasia. Surg Endosc 2019; 33: 3656-3664 [PMID: 30671667 DOI: 10.1007/s00464-018-06654-1]
- Liu Z, Wang Y, Fang Y, Huang Y, Yang H, Ren X, Xu M, Chen S, Chen W, Zhong Y, Zhang Y, Qin W, Hu J, Cai M, Yao L, Li Q, Zhou P. 34 Short-term safety and efficacy of peroral endoscopic myotomy for the treatment of achalasia in children. J Gastroenterol 2020; 55: 159-168 [PMID: 31679066 DOI: 10.1007/s00535-019-01607-4]
- Saez J, Mejia R, Pattillo JC, Vuletin F, Monrroy H, Jaime F, Sharp A. Per oral endoscopic myotomy (POEM) in pediatric patients with 35 esophageal achalasia: First Latin-American experience. J Pediatr Surg 2021; 56: 706-710 [PMID: 32646661 DOI: 10.1016/i.jpedsurg.2020.06.007
- Wood LS, Chandler JM, Portelli KE, Taylor JS, Kethman WC, Wall JK. Treating children with achalasia using per-oral endoscopic myotomy 36 (POEM): Twenty-one cases in review. J Pediatr Surg 2020; 55: 1006-1012 [PMID: 32197825 DOI: 10.1016/j.jpedsurg.2020.02.028]
- Peng D, Tan Y, Li C, Lv L, Zhu H, Liang C, Li R, Liu D. Peroral Endoscopic Myotomy for Pediatric Achalasia: A Retrospective Analysis of 37 21 Cases With a Minimum Follow-Up of 5 Years. Front Pediatr 2022; 10: 845103 [PMID: 35444970 DOI: 10.3389/fped.2022.845103]
- Petrosyan M, Mostammand S, Shah AA, Darbari A, Kane TD. Per Oral Endoscopic Myotomy (POEM) for pediatric achalasia: Institutional 38 experience and outcomes. J Pediatr Surg 2022; 57: 728-735 [PMID: 35361482 DOI: 10.1016/j.jpedsurg.2022.02.017]
- Wang X, Tan Y, Lv L, Zhu H, Chu Y, Li C, Liu D. Peroral endoscopic myotomy versus pneumatic dilation for achalasia in patients aged ≥ 65 39 years. Rev Esp Enferm Dig 2016; 108: 637-641 [PMID: 27649684 DOI: 10.17235/reed.2016.4488/2016]
- 40 Landi R, Familiari P, Mangiola F, Calì A, D'aversa F, Bove V, Boskoski I, Tringali A, Perri V, Costamagna G. Tu1154 Peroral Endoscopic Myotomy for The Treatment of Esophageal Motility Disorders: Baseline Characteristics And Procedural Outcomes In Young And Elderly Patients. Gastrointest Endosc 2018; 87: AB544-AB545 [DOI: 10.1016/j.gie.2018.04.2184]
- Liu XY, Cheng J, Chen WF, Xu MD, Liu ZQ, Wang Y, Chen SY, Zhong YS, Zhang YQ, Qin WZ, Hu JW, Cai MY, Yao LQ, Li QL, Zhou 41 PH. Effect of peroral endoscopic myotomy in geriatric patients: a propensity score matching study. Surg Endosc 2020; 34: 2911-2917 [PMID:



31451922 DOI: 10.1007/s00464-019-07070-9]

- Klair JS, Hasan Y, Masadeh M, Gerke H, Murali A, Parekh K, Keech J, Nau P, El Abiad R. 379 Safety and Efficacy of Peroral Endoscopic 42 Myotomy (POEM) in Patients ≥ 65 Years at a Single Academic Tertiary Care Center. Am J Gastroenterol 2019; 114: S221-S221 [DOI: 10.14309/01.ajg.0000591048.43221.3c]
- Abe H, Tanaka S, Kawara F, Toyonaga T, Ariyoshi R, Sakaguchi H, Sako T, Ikezawa N, Kodama Y. Comparison of the safety and efficacy of 43 peroral endoscopic myotomy between octogenarians and non-octogenarians. Dig Endosc 2021; 33: 110-117 [PMID: 32274834 DOI: 10.1111/den.13686]
- Sanaka MR, Chadalavada P, Alomari M, Tang A, Parikh M, Garg R, Gupta N, Thota P, Gabbard S, Murthy S, Raja S. Peroral endoscopic 44 myotomy is a safe and effective treatment modality for geriatric patients with achalasia. Esophagus 2020; 17: 484-491 [PMID: 32394115 DOI: 10.1007/s10388-020-00746-5
- Angeli Abad MR, Inoue H, Fujiyoshi Y, Nishikawa Y, Sakaguchi T, Shimamura Y, Sumi K, Kimura R, Izawa S, Ikeda H, Onimaru M. 45 Sa1268 Clinical Outcomes of Peroral Endoscopic Myotomy In Octo- and Nonagenarian Achalasia Patients: A Large Single-Center Experience. Gastrointest Endosc 2020; 91: AB141 [DOI: 10.1016/j.gie.2020.03.1064]
- Okada H, Shiwaku H, Ohmiya T, Shiwaku A, Hasegawa S. Efficacy and safety of peroral endoscopic myotomy in 100 older patients. 46 Esophagus 2022; 19: 324-331 [PMID: 34626277 DOI: 10.1007/s10388-021-00881-7]
- Nakamura J, Hikichi T, Hashimoto M, Takasumi M, Kato T, Kobashi R, Yanagita T, Suzuki R, Sugimoto M, Sato Y, Irie H, Takagi T, 47 Kobayakawa M, Ohira H. Efficacy and Safety of Peroral Endoscopic Myotomy for Esophageal Achalasia and Achalasia-Related Diseases in Patients Aged 75 Years and Over. Healthcare (Basel) 2021; 9 [PMID: 34946392 DOI: 10.3390/healthcare9121668]
- Ujiie N, Sato C, Taniyama Y, Koseki K, Takaya K, Okamoto H, Fukutomi T, Unno M, Kamei T. Characteristics of esophageal achalasia in 48 geriatric patients over 75 years of age and outcomes after peroral endoscopic myotomy. Geriatr Gerontol Int 2021; 21: 788-793 [PMID: 34250704 DOI: 10.1111/ggi.14235]
- Zhao X, Chai N, Wu Q, Du R, Ye L, Li X, Linghu E. E-Poster Area: Blue Ribbon Posters. Dig Endosc 2022; 34: 64-213 [DOI: 49 10.1111/den.14292]
- Bartenstein P, Schober O. [In vivo imaging of benzodiazepine receptors of the central type]. Wien Klin Wochenschr 1991; 103: 647-648 50 [PMID: 1663299 DOI: 10.1097/01.sla.0000217524.75529.2d]
- Snyder CW, Burton RC, Brown LE, Kakade MS, Finan KR, Hawn MT. Multiple preoperative endoscopic interventions are associated with 51 worse outcomes after laparoscopic Heller myotomy for achalasia. J Gastrointest Surg 2009; 13: 2095-2103 [PMID: 19789928 DOI: 10.1007/s11605-009-1049-6]
- Zhong C, Huang S, Xia H, Tan S, Lü M, Peng Y, Tang X. Role of Peroral Endoscopic Myotomy in Geriatric Patients with Achalasia: A 52 Systematic Review and Meta-Analysis. Dig Dis 2022; 40: 106-114 [PMID: 33752208 DOI: 10.1159/000516024]
- Luvsandagva B, Adyasuren B, Bagachoimbol B, Luuzanbadam G, Bai T, Jalbuu N, Duger D, Hou X. Efficacy and safety of peroral 53 endoscopic myotomy for pediatric achalasia: A nationwide study. Medicine (Baltimore) 2024; 103: e38970 [PMID: 39121306 DOI: 10.1097/MD.00000000038970]
- Lee Y, Brar K, Doumouras AG, Hong D. Peroral endoscopic myotomy (POEM) for the treatment of pediatric achalasia: a systematic review 54 and meta-analysis. Surg Endosc 2019; 33: 1710-1720 [PMID: 30767141 DOI: 10.1007/s00464-019-06701-5]
- Zhong C, Tan S, Huang S, Peng Y, Lü M, Tang X. Clinical outcomes of peroral endoscopic myotomy for achalasia in children: a systematic 55 review and meta-analysis. Dis Esophagus 2021; 34 [PMID: 33316041 DOI: 10.1093/dote/doaa112]
- Peng D, Tan Y, Chen X, Lv L, Zhu H, Li R, Liu D. Prognostic factors analysis of symptoms recurrence after peroral endoscopic myotomy for 56 the treatment of achalasia in older adults. J Gastrointest Surg 2024; 28: 1349-1351 [PMID: 38796138 DOI: 10.1016/j.gassur.2024.05.023]



WJGE | https://www.wjgnet.com



Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

