







Supplementary Table 1


Retrieval strategy

Search: (((((((("Therapy, Neoadjuvant") OR ("Neoadjuvant Treatment")) OR ("Neoadjuvant Chemotherapy")) OR ("Chemotherapy, Neoadjuvant")) OR ("Neoadjuvant Chemotherapies")) OR ("Neoadjuvant Therapy"[Mesh])) AND (((((((((((((((("Colorectal Neoplasm") OR ("Neoplasm, Colorectal")) OR ("Neoplasms, Colorectal")) OR ("Colorectal Tumors")) OR ("Colorectal Tumor")) OR ("Tumor, Colorectal")) OR ("Tumors, Colorectal")) OR ("Colorectal Cancer")) OR ("Cancer, Colorectal")) OR ("Cancers, Colorectal")) OR ("Colorectal Cancers")) OR ("Colorectal Carcinoma")) OR ("Carcinoma, Colorectal")) OR ("Carcinomas, Colorectal")) OR ("Colorectal Carcinomas")) OR ("Colorectal Neoplasms"[Mesh])))) AND (((("colorectal liver metastases") OR ("colonic liver metastases")) OR ("rectal liver metastases"))









Supplementary Figure 1 Sensitivity analyses results. A: Sensitivity analysis of duration of surgery; B: Sensitivity analysis of the number of liver metastases; C: Sensitivity analyses of duration of the length of hospital stay; D: Sensitivity analysis of the size of the largest metastasis; E: Sensitivity analysis of blood loss during surgery; F: Sensitivity analysis of synchronous metastases; G: Sensitivity analysis of perioperative complications.