

Name of Journal: *World Journal of Gastrointestinal Surgery*

ESPS Manuscript NO: 21146

Manuscript Type: Review

Risk Factors and Implications of Anastomotic <sup>2</sup>Complications After Surgery for Crohn's Disease

Kristen T. Crowell, Evangelos Messaris

### Abstract

Anastomotic complications <sup>5</sup>occur more frequently in patients with Crohn's disease leading to postoperative <sup>1</sup>intra-abdominal septic complications (IASC). Patients with IASC often require re-operation or drainage to control the sepsis and have an increased frequency of disease <sup>9</sup>recurrence. The aim of this article was to examine the factors

### Match Overview

Match Number	Source	Words	Percentage
1	CrossCheck	77 words	2%
	Shental, Omri, Hagit Tulchinsky, Ron Greenberg, Josep h M. Klausner, and Shmuel Avital. "Positive Histological I		
2	CrossCheck	41 words	1%
	Takayuki Yamamoto. "Risk factors for intra-abdominal ... eptis after surgery in Crohn's disease", Diseases of the		
3	Internet	28 words	1%
	crawled on 04-Nov-2014 onlinelibrary.wiley.com		
4	CrossCheck	27 words	1%
	for the European Crohn's and Colitis Organisation (ECC O). "The second European evidence-based consensus		
5	CrossCheck	25 words	1%
	Aberra, Faten N, and Gary R Lichtenstein. "Methods to ... oid infections in Patients with Inflammatory Bowel Disea		
6	CrossCheck	23 words	1%
	Paparo, Francesco, Matteo Revelli, Cristina Puppo, Lore nzo Bacigalupo, Isabella Garello, Alessandro Gartaschi,		





Risk Factors and Implications of Anastomotic Complications After Surgery for



学术搜索

找到约 15,300 条结果 (用时0.09秒)

我的著作引用情况

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

包括专利

包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

### Risk factors for intra-abdominal sepsis after surgery in Crohn's disease

T Yamamoto, RN Allan, MRB Keighley - *Diseases of the colon & rectum*, 2000 - Springer

... risk factors, 16 per-cent (33/209 operations) in patients with only one risk factor, and 5 ... and fistula at the time of laparotomy were independent significant risk factors for postoperative ... Our results indicate that multiple anastomoses can be used without additional risks in Crohn's ...

被引用次数: 212 相关文章 所有 8 个版本 引用 保存

researchgate.net 中的 [PDF]

### Risks of intestinal anastomoses in Crohn's disease.

S Post, M Betzler, B von Ditfurth, G SCHÜRMANN... - ... of surgery, 1991 - ncbi.nlm.nih.gov

Risks of Intestinal Anastomoses in Crohn's Disease ... in noninflammatory diseases.6 In Crohn's disease intestinal resections frequently are connected with factors known or ... more than one anastomosis during the same procedure (maximum, 9) without increased risk (Table 3 ...

被引用次数: 153 相关文章 所有 6 个版本 引用 保存

nih.gov 中的 [PDF]

### Risk factors for intra-abdominal septic complications after a first ileocecal resection for Crohn's disease: a multivariate analysis in 161 consecutive patients

A Alves, Y Paris, Y Bouhnik, M Pocard, E Vicaut... - *Diseases of the colon & ...*, 2007 - Springer

... S, Betzler M, von Ditfurth B, Schümann G, Küppers P, Herfarth C. Risks of intestinal ... Risk factors for intra-abdominal sepsis after surgery in Crohn's disease. ... Factors affecting the incidence of postoperative septic complications and recurrence after stricturoplasty for jejunoileal ...

被引用次数: 140 相关文章 所有 8 个版本 引用 保存

### Early postoperative complications are not increased in patients with Crohn's disease treated perioperatively with infliximab or immunosuppressive therapy

JF Colombel, EV Loftus, WJ Tremaine... - *The American journal of ...*, 2004 - nature.com

... Antitumor necrosis factor therapy for inflammatory bowel disease: A review of agents.

nature.com 中的 [HTML]







Scholar

About 14,700 results (0.14 sec)

My Citations

Articles

Case law

My library

Any time

Since 2015

Since 2014

Since 2011

Custom range...

Sort by relevance

Sort by date

 include patents

 include citations

 Create alert

### Risk factors for intra-abdominal sepsis after surgery in Crohn's disease

T Yamamoto, RN Allan, MRB Keighley - *Diseases of the colon & rectum*, 2000 - Springer

... **risk factors**, 16 per- cent (33/209 operations) in patients with only one **risk factor**, and 5 ... and fistula at the time of laparot- omy were independent significant **risk factors** for postoperative ... Our results indicate that multiple anas- tomoses can be used without additional **risks** in **Crohn's** ...

Cited by 208 Related articles All 8 versions Cite Save

[PDF] from lww.com

Get It! @ Penn State

### Risks of intestinal anastomoses in Crohn's disease.

S Post, M Betzler, B von Ditfurth, G SCHÜRMAN... - ... of *surgery*, 1991 - ncbi.nlm.nih.gov

**Risks of Intestinal Anastomoses in Crohn's Disease** ... in noninflammatory **diseases**.6 In **Crohn's disease** intestinal resections frequently are connected with **factors** known or ... more than one **anastomosis** during the same **procedure** (maximum, 9) without increased **risk** (Table 3 ...

Cited by 151 Related articles All 6 versions Cite Save

[PDF] from nih.gov

Get It! @ Penn State

### Risk factors for intra-abdominal septic complications after a first ileocecal resection for Crohn's disease: a multivariate analysis in 161 consecutive patients

A Alves, Y Panis, Y Bouhnik, M Pocard, E Vicaut... - *Diseases of the colon & ...*, 2007 - Springer

... S, Betzler M, von Ditfurth B, Schümann G, Küppers P, Herfarth C. **Risks of intestinal ... Risk factors** for intra-abdominal sepsis **after surgery** in Crohn\_s **disease**. ... **Factors** affecting the incidence of postoperative septic **complications** and recurrence **after** stricturoplasty for jejunoileal ...

Cited by 135 Related articles All 8 versions Cite Save

[PDF] from lww.com

Get It! @ Penn State

### Early postoperative complications are not increased in patients with Crohn's disease treated perioperatively with infliximab or immunosuppressive therapy

JF Colombel, EV Loftus, WJ Tremaine... - *The American journal of ...*, 2004 - nature.com

... Antitumor necrosis **factor** therapy for inflammatory bowel **disease**: A review of agents, pharmacology, clinical results, and safety. ... **Risk factors** for intra-abdominal sepsis **after surgery** in **Crohn's disease**. ... **Risks of intestinal anastomoses in Crohn's disease**. ...

Cited by 249 Related articles All 6 versions Cite Save

[HTML] from nature.com

Get It! @ Penn State

### [PDF] Factors affecting recurrence after surgery for Crohn's disease

T Yamamoto - *World Journal of Gastroenterology*, 2005 - researchgate.net

... The endoscopic and radiological rate of recurrence was also significantly decreased with relative **risks** of 0.654 (90%CI, 0.47 ... Thus, the length of resected bowel does not seem to be a definite predictive **factor** of post ... **Risk factors** for postoperative recurrence of **Crohn's disease**. ...

Cited by 197 Related articles All 7 versions Cite Save More

[PDF] from researchgate.net

Get It! @ Penn State

### Risk factors for anastomotic leakage after surgery for colorectal cancer: results of prospective surveillance

T Konishi, T Watanabe, J Kishimoto... - ... *College of Surgeons*, 2006 - Elsevier

... suggested that **anastomosis** below the peritoneal reflection is an important **risk factor** for

[HTML] from journalacs.org



[网页](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 244,000 条结果（用时 0.73 秒）

## Google 学术: Risk factors and implications of anastomotic complications after surgery for Crohn's disease

... intra-abdominal sepsis after surgery in Crohn's disease - Yamamoto - 被引用次数: 213

Risks of intestinal anastomoses in Crohn's disease. - Post - 被引用次数: 153

Risk factors for intra-abdominal septic complications ... - Alves - 被引用次数: 141

### Risk factors for intra-abdominal sepsis after surgery in ...

[www.ncbi.nlm.nih.gov/pubmed/10950014](http://www.ncbi.nlm.nih.gov/pubmed/10950014) ▾ 翻译此页

作者: T Yamamoto - 2000 - 被引用次数: 213 - 相关文章

Risk factors for intra-abdominal sepsis after surgery in Crohn's disease. ... RESULTS:  
Intra-abdominal septic complications, defined as anastomotic leak, ...

### Risk factors for postoperative intra-abdominal septic ...

[www.ncbi.nlm.nih.gov/pubmed/22874602](http://www.ncbi.nlm.nih.gov/pubmed/22874602) ▾ 翻译此页

作者: A Kanazawa - 2012 - 被引用次数: 29 - 相关文章

Risk factors for postoperative intra-abdominal septic complications after bowel resection in patients with Crohn's disease. ... Adult; Aged; Anastomosis, Surgical/adverse effects; Anastomotic Leak/etiology; Crohn Disease/complications; Crohn ...

### Influence of risk factors on the safety of ileocolic ...

[www.ncbi.nlm.nih.gov/pubmed/22513434](http://www.ncbi.nlm.nih.gov/pubmed/22513434) ▾ 翻译此页

作者: A Tzivanakis - 2012 - 被引用次数: 36 - 相关文章

Influence of risk factors on the safety of ileocolic anastomosis in Crohn's disease surgery. ... with their associated morbidity, are the most feared risks of surgery.