

World Journal of *Diabetes*

World J Diabetes 2024 July 15; 15(7): 1384-1653



EDITORIAL

- 1384 Remission of type 2 diabetes mellitus
Nakhleh A, Halfin E, Shehadeh N
- 1390 Diabetes remission and nonalcoholic fatty pancreas disease
Wu WJ
- 1394 Management of gestational diabetes mellitus *via* nutritional interventions: The relevance of gastric emptying
Huang WK, Jalleh RJ, Rayner CK, Wu TZ
- 1398 MicroRNA-630: A promising avenue for alleviating inflammation in diabetic kidney disease
Donate-Correa J, González-Luis A, Díaz-Vera J, Hernandez-Fernaud JR
- 1404 Adiposity in Chinese people with type 1 diabetes
Wu NW, Lyu XF, An ZM, Li SY
- 1409 Diabetes and tuberculosis: An emerging dual threat to healthcare
Shetty S, Pappachan JM, Fernandez CJ

REVIEW

- 1417 Patient-centered care in diabetes care-concepts, relationships and practice
Chen TT, Su WC, Liu MI
- 1430 Insulin resistance as the molecular link between diabetes and Alzheimer's disease
Abdalla MMI

MINIREVIEWS

- 1448 Obstructive sleep apnea: Overlooked comorbidity in patients with diabetes
Tenda ED, Henrina J, Cha JH, Triono MR, Putri EA, Aristy DJ, Tahapary DL
- 1461 Update on evidence-based clinical application of sodium-glucose cotransporter inhibitors: Insight to uncommon cardiovascular disease scenarios in diabetes
Tao SB, Lu X, Ye ZW, Tong NW

ORIGINAL ARTICLE**Retrospective Cohort Study**

- 1477 Association between glucose levels of children with type 1 diabetes and parental economic status in mobile health application

Zhang WH, Wang CF, Wang H, Tang J, Zhang HQ, Zhu JY, Zheng XY, Luo SH, Ding Y

Retrospective Study

- 1489 Association between glucose-lowering drugs and circulating insulin antibodies induced by insulin therapy in patients with type 2 diabetes

Zhang P, Jiang Q, Ding B, Yan RN, Hu Y, Ma JH

- 1499 Clinical efficacy of endovascular revascularization combined with vacuum-assisted closure for the treatment of diabetic foot

Lei FR, Shen XF, Zhang C, Li XQ, Zhuang H, Sang HF

- 1509 Magnetic resonance imaging combined with serum endolipin and galactaglobin-3 to diagnose cerebral infarction in the elderly with diabetes mellitus

Zhang YH, Liang D

- 1518 Dapagliflozin in heart failure and type 2 diabetes: Efficacy, cardiac and renal effects, safety

Yu PL, Yu Y, Li S, Mu BC, Nan MH, Pang M

Observational Study

- 1531 Cut-off value of glycated hemoglobin A1c for detecting diabetic retinopathy in the Chinese population

Wen Y, Wang Q

- 1537 Glymphatic function and its influencing factors in different glucose metabolism states

Tian B, Zhao C, Liang JL, Zhang HT, Xu YF, Zheng HL, Zhou J, Gong JN, Lu ST, Zeng ZS

Clinical and Translational Research

- 1551 Does type 1 diabetes serve as a protective factor against inflammatory bowel disease: A Mendelian randomization study

Tong KK, Yu YF, Yang XY, Wu JY, Yu R, Tan CC

- 1562 Network pharmacology and molecular dynamics study of the effect of the *Astragalus-Coptis* drug pair on diabetic kidney disease

Zhang MY, Zheng SQ

Basic Study

- 1589 Interactions between myoblasts and macrophages under high glucose milieu result in inflammatory response and impaired insulin sensitivity

Luo W, Zhou Y, Wang LY, Ai L

SYSTEMATIC REVIEWS

- 1603** Natural product-based treatment potential for type 2 diabetes mellitus and cardiovascular disease
Shrivastav D, Kumbhakar SK, Srivastava S, Singh DD

META-ANALYSIS

- 1615** Evaluation of teplizumab's efficacy and safety in treatment of type 1 diabetes mellitus: A systematic review and meta-analysis
Ma XL, Ge D, Hu XJ

SCIENTOMETRICS

- 1627** Global trends in publications regarding macrophages-related diabetic foot ulcers in the last two decades
Wen JP, Ou SJ, Liu JB, Zhang W, Qu YD, Li JX, Xia CL, Yang Y, Qi Y, Xu CP

LETTER TO THE EDITOR

- 1645** Atrial fibrillation and prediabetes: A liaison that merits attention!
Batta A, Hatwal J
- 1648** Serum tumor markers: Can they clinically implicate in type 2 diabetes mellitus?
Reddy KS, Pandiaraj IP, Gaur A, Varatharajan S
- 1651** Bidirectional link between periodontitis and systemic inflammation in diabetic retinopathy
Nishant P, Sinha S, Sinha RK, Morya AK

ABOUT COVER

Peer Review of *World Journal of Diabetes*, Erkan Gokce, MD, Professor, Department of Radiology, Tokat Gaziosmanpasa University, School of Medicine, Tokat 60100, Türkiye. drerkangokce@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Diabetes (WJD, World J Diabetes)* is to provide scholars and readers from various fields of diabetes with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJD mainly publishes articles reporting research results and findings obtained in the field of diabetes and covering a wide range of topics including risk factors for diabetes, diabetes complications, experimental diabetes mellitus, type 1 diabetes mellitus, type 2 diabetes mellitus, gestational diabetes, diabetic angiopathies, diabetic cardiomyopathies, diabetic coma, diabetic ketoacidosis, diabetic nephropathies, diabetic neuropathies, Donohue syndrome, fetal macrosomia, and prediabetic state.

INDEXING/ABSTRACTING

The *WJD* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for *WJD* as 4.2; JIF without journal self cites: 4.1; 5-year JIF: 4.2; JIF Rank: 40/186 in endocrinology and metabolism; JIF Quartile: Q1; and 5-year JIF Quartile: Q2.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Yu-Xi Chen*; Production Department Director: *Xu Guo*; Cover Editor: *Jia-Ru Fan*.

NAME OF JOURNAL

World Journal of Diabetes

ISSN

ISSN 1948-9358 (online)

LAUNCH DATE

June 15, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Lu Cai, Md. Shahidul Islam, Michael Horowitz

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9358/editorialboard.htm>

PUBLICATION DATE

July 15, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Remission of type 2 diabetes mellitus

Afif Nakhleh, Elya Halfin, Naim Shehadeh

Specialty type: Endocrinology and metabolism

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade B, Grade D

Novelty: Grade B, Grade C

Creativity or Innovation: Grade B, Grade C

Scientific Significance: Grade B, Grade C

P-Reviewer: Srinivasan AR, India

Received: December 30, 2023

Revised: April 13, 2024

Accepted: April 28, 2024

Published online: July 15, 2024

Processing time: 190 Days and 12.9 Hours



Afif Nakhleh, Naim Shehadeh, Institute of Endocrinology, Diabetes and Metabolism, Rambam Health Care Campus, Haifa 3109601, Israel

Afif Nakhleh, Elya Halfin, Naim Shehadeh, Diabetes and Endocrinology Clinic, Maccabi Healthcare Services, Haifa 3299001, Israel

Afif Nakhleh, Naim Shehadeh, The Azrieli Faculty of Medicine, Bar-Ilan University, Safed 1311502, Israel

Corresponding author: Afif Nakhleh, MD, Doctor, Institute of Endocrinology, Diabetes and Metabolism, Rambam Health Care Campus, 8 Haaliya Hashniya Street, Haifa 3109601, Israel. anakhleh@gmail.com

Abstract

The surge in type 2 diabetes mellitus (T2DM) is tightly linked to obesity, leading to ectopic fat accumulation in internal organs. Weight management has become a cornerstone of T2DM treatment, with evidence suggesting that significant weight loss can induce remission. Remission, defined as sustained hemoglobin (HbA1c) below 6.5% for at least 3 months without medication, can be achieved through various approaches, including lifestyle, medical, and surgical interventions. Metabolic bariatric surgery offers significant remission rates, particularly for patients with severe obesity. Intensive lifestyle modifications, including low-calorie diets and exercise, have also demonstrated significant potential. Medications like incretin-based agents show robust results in improving beta-cell function, achieving glycemic control, and promoting weight loss. While complete remission without medication may not be attainable for everyone, especially those with severe insulin resistance or deficiency, early and aggressive glycemic control remains a crucial strategy. Maintaining HbA1c below 6.5% from the time of diagnosis reduces the risk of long-term complications and mortality. Moreover, considering a broader definition of remission, encompassing individuals with sustained control on medication, could offer a more comprehensive and inclusive approach to managing this chronic disease.

Key Words: Type 2 diabetes mellitus; Remission; Lifestyle intervention; Metabolic bariatric surgery; Glucose-lowering medications; Weight management medications

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The rise in type 2 diabetes mellitus (T2DM) parallels the obesity epidemic. Notably, significant weight loss can induce remission, with hemoglobin levels below 6.5% for three months without medication. Herein we present the recent data on a variety of approaches, including lifestyle interventions, glucose-lowering and weight management medications and metabolic surgery that hold promise for inducing T2DM remission.

Citation: Nakhleh A, Halfin E, Shehadeh N. Remission of type 2 diabetes mellitus. *World J Diabetes* 2024; 15(7): 1384-1389

URL: <https://www.wjgnet.com/1948-9358/full/v15/i7/1384.htm>

DOI: <https://dx.doi.org/10.4239/wjd.v15.i7.1384>

INTRODUCTION

The surge in type 2 diabetes mellitus (T2DM) is undeniably linked to the alarming rise in obesity. While mammalian physiology utilizes subcutaneous adipose tissue as a natural reservoir for energy storage, obesity disrupts this equilibrium. It triggers ectopic fat deposition, where adipose tissue infiltrates internal organs such as the liver, pancreas, heart, kidneys, and muscles[1]. Therefore, in most cases of newly diagnosed T2DM, weight management has moved to the forefront of treatment approach, as growing evidence suggests that weight loss interventions can lead to T2DM remission[2].

The recent consensus statement defines T2DM remission as a sustained return of glycated hemoglobin (HbA1c) levels to below 6.5% (or 48 mmol/mol) for at least 3 months, without the use of glucose-lowering medications[3]. This improvement can occur spontaneously or through intervention, but crucially, it must be achieved and maintained without medications. In situations where HbA1c is not a reliable indicator of long-term blood sugar control, alternative criteria can be employed. These include a fasting plasma glucose level below 126 mg/dL (7.0 mmol/L) or an estimated HbA1c value of less than 6.5%, calculated using continuous glucose monitoring data. This consensus definition will help standardize future studies. However, the current definition does not include individuals who achieve remission (HbA1c < 6.5%) while maintaining glucose-lowering medications.

Several therapeutic modalities can be employed to achieve T2DM remission, including intensive lifestyle modifications, pharmacological agents, and surgical interventions targeting glycemic control and weight management.

INTENSIVE LIFESTYLE INTERVENTIONS FOR T2DM REMISSION

Studies employing intensive lifestyle modification programs have demonstrated that substantial weight loss in patients with T2DM is successful in reversing hyperglycemia and lowering risk factors.

The DiRECT trial, which assessed a very low-calorie diet using meal replacements, demonstrated remission rates of 46% at 1 year and 36% at 2 years versus 4.0% and 3.4% in control groups, respectively[4]. Another similar trial, DIADEM-I, achieved even a higher remission rate of 61% at 1 year by using short-term, very low-calorie diets and facilitating a significant weight loss of 12 kg[5].

The recently published DiRECT-Aus found that over half (56%) of newly diagnosed patients with T2DM achieved remission after 1 year of intensive lifestyle intervention using low-energy total diet replacement[6]. Remission rates were similar to previous studies in other parts of the world, suggesting the effectiveness of this approach across ethnicities and cultures. DiRECT-Aus suggests prioritizing intensive lifestyle interventions, especially within 6 years of diagnosis, to maximize remission chances[6].

While the LOOK AHEAD trial's intensive diet and exercise program achieved remission in only 11.5% of participants after 1 year and 7.3% after 4 years[7], a follow-up study revealed a crucial link between weight loss and long-term health benefits[8].

This study found that participants who maintained remission for a longer period had a greater decrease in systolic blood pressure and a larger increase in high-density lipoprotein cholesterol. However, there was only a small reduction in low-density lipoprotein cholesterol[9].

The LOOK AHEAD study found that the greater the weight loss, the lower the risk of developing cardiovascular disease over 10 years. Specifically, a 10% weight loss was associated with a 21% reduction in cardiovascular disease risk [8].

We can speculate that the low rates of diabetes remission in the LOOK AHEAD trial as compared to the DiRECT trial may be related to the longer T2DM duration of participants in the LOOK AHEAD trial (mean of 5 years) versus 3 years in the DiRECT trial.

A recent post-hoc analysis of the DiRECT study offers compelling evidence that weight loss-induced remission can lead to a remarkable reversal of abnormal pancreas structure in individuals with T2D. After 2 years of remission, the study found a significant increase in pancreas volume and a decrease in its shape irregularity[10]. This remarkable finding suggests that gradual morphological and functional recovery of the entire pancreas is possible during remission, highlighting the potential for T2DM reversal.

Other approaches, including time-restricted eating[11] and even high-protein diets[12], have successfully induced remission.

Emerging evidence suggests that incorporating exercise into diabetes management plans can effectively enhance beta cell function in a dose-dependent manner, which is crucial for better glycemic control and induction of diabetes remission. For adults with newly diagnosed T2DM, adding regular exercise to a weight-loss program significantly improved their beta cell function over a 16-wk period. The most significant benefits were seen in those who exercised six times per week[13].

Lean individuals [body mass index (BMI) < 25 kg/m²] with T2DM can also benefit from an intensive physical activity and dietary program to achieve diabetes remission. A recent study in China showed that with a program combining a low-carbohydrate diet, exercise, and diabetes management support, approximately 60% of lean individuals with T2DM achieved diabetes remission (HbA1c < 6.5%) after 6 months without the need for any anti-diabetic medications[14].

Overall, these trials demonstrate that well-supported lifestyle changes can achieve T2DM remission in a substantial proportion of individuals, with benefits extending beyond remission to weight loss and likely improved overall health. However, the current evidence underscores the importance of early intervention, coinciding with the putative timeframe of preserved beta-cell function.

MEDICAL INTERVENTIONS-GLUCOSE-LOWERING AND WEIGHT MANAGEMENT MEDICATIONS FOR T2DM REMISSION

T2DM is characterized by a progressive decline in beta-cell function, which is already reduced to approximately 50% of normal function at diagnosis[15]. Short-term intensive insulin therapy has been investigated for potential to reverse this decline and achieve diabetes remission. This approach involves actively titrating insulin for 2 wk to attain beta-cell rest and glycemic control. This can improve beta-cell function and enhance insulin sensitivity. Studies have shown that intensive insulin therapy can lead to drug-free T2DM remission in 42% of participants after 2 years, with higher success rates observed in those who initiate treatment earlier[16]. However, potential drawbacks like the complexity of treatment and risk of hypoglycemia limit its use[17].

Studies suggest that initial insulin therapy followed by non-insulin antidiabetic drugs (NIAD) can preserve beta-cell function as compared to NIADs alone, potentially leading to better long-term glycemic control. Studies like REMIT and REMIT-dapa exemplify this approach[18,19]. In REMIT, the intervention group received a combination of metformin, acarbose, and insulin glargine[18]. REMIT-dapa used a similar approach with insulin glargine, metformin, and dapagliflozin[19]. Both studies demonstrated the potential of intensive combined therapy with NIADs and insulin. Up to 25% of patients in REMIT and 14% in REMIT-dapa achieved remission after 1 year, with the likelihood of success increasing with a longer treatment duration[18,19].

Incretin-based medications like glucagon-like peptide 1 receptor agonists (GLP-1 RAs), GLP-1/glucose-dependent insulinotropic polypeptide (GIP) dual agonists, and GLP-1/GIP/glucagon tri-agonists can improve glycemic control, preserve beta-cell function, and promote weight loss in patients with T2DM[2,20]. This suggests they may offer less-invasive diabetes remission potential as compared to more aggressive interventions like metabolic bariatric surgery (MBS).

Semaglutide significantly reduced HbA1c to ≤ 6.5% in 60% to 68% of T2DM patients with overweight or obesity over 68 wk[21].

GLP-1/GIP dual agonists are even more effective for glycemic control and weight loss. Furthermore, they are associated with greater beta-cell function and improvements in insulin sensitivity as compared to GLP-1 RAs alone. In the SURPASS-4 trial, 66% to 81% of T2DM patients on tirzepatide achieved an HbA1c ≤ 6.5% after 1 year[22]. In SURMOUNT-1, a remarkable 95.3% of participants with prediabetes and obesity in the tirzepatide groups achieved normoglycemia by week 72[23]. In a recent study of overweight or obese T2DM patients, the GLP-1/GIP/glucagon tri-agonist retatrutide successfully lowered HbA1c levels, with up to 82% of participants reaching ≤ 6.5% after 24 wk[24].

Cases of T2DM “remission” achieved through incretin-based medications do not technically meet the proposed definition of remission and only persist while patients are keeping their weight loss achievement mostly by actively taking the medication. However, achieving medication-free remission may not be feasible for all T2DM patients. This is due to the diverse nature of T2DM, including factors like severe insulin resistance (particularly in non-obese individuals) or marked insulin deficiency, as identified in the recent adult-onset diabetes subclassification by Ahlqvist *et al*[25].

Although complete remission without glucose-lowering medication may not be achievable for all patients, we firmly advocate for early and proactive glycemic control as a crucial aspect of long-term diabetes management. Maintaining good control early on can prevent future complications and improve outcomes. The legacy effect supports this approach. A large cohort study with a 13-year follow-up period emphasized the importance of achieving good glycemic control (HbA1c < 6.5%) within the first year of diagnosis. The study clearly demonstrates that the longer HbA1c levels remain below 6.5%, the greater the protection against both microvascular and macrovascular complications, and the lower the mortality risk. Conversely, sustained exposure to higher HbA1c levels significantly increases these risks. Therefore, taking prompt action to achieve early glycemic control can significantly enhance long-term outcomes and prevent debilitating complications.

SURGICAL INTERVENTIONS FOR T2DM REMISSION

MBS can reverse T2DM in many patients, especially those with severe obesity (BMI \geq 35). Weight reduction can result in better glycemic control, reduced medication use, improved quality of life, and improved cardiovascular outcomes[26]. Among MBS procedures, malabsorptive techniques like Roux-en-Y gastric bypass (RYGB), biliopancreatic diversion (BPD) and the newly used mini-gastric bypass surgery achieve higher remission rates compared to purely restrictive methods like gastric banding. It's worth noting, however, that the utilization of BPD has waned in recent years[27]. Sleeve gastrectomy (SG) is a restrictive procedure that offers promising results; its mechanism primarily reduces stomach volume, ultimately falling behind the efficacy of malabsorptive approaches that alter the digestive pathway.

For severe obesity (BMI \geq 35) with T2DM, RYGB offers a 67% remission rate at 3 years that is double that of SG (33%) [28]. At 5 years of follow-up, RYGB demonstrates superior and sustained glycemic control as compared to SG, with a 25% lower rate of T2DM relapse[29]. However, remission rates typically decrease over time, with relapse rates reaching around 50% after 5 years for both RYGB and SG[28]. The likelihood of T2DM remission after MBS surgery increases with greater weight loss, younger age, shorter duration of diabetes, reasonable pre-operative glycemic control, fewer diabetes medications, and not using insulin[30]. Careful patient selection, risk/benefit assessment, and lifelong monitoring are crucial for successful MBS implementation. More research is needed on long-term outcomes and expanding access to MBS for eligible patients. Studies exploring the potential for remission of T2DM with MBS lack a standardized definition of "remission", making comparisons and conclusions challenging. Furthermore, these studies often relied on comparator groups focused on glycemic control rather than weight loss. While MBS has demonstrated promising results in achieving sustained weight loss and potentially inducing T2DM remission, its broader availability remains limited.

CONCLUSION

Current evidence suggests that T2DM can be effectively managed and potentially put into remission through various lifestyle, medical, and surgical interventions. In these interventions, weight loss remains the primary driver of remission. Significant weight loss (greater than 10%) and early intervention after T2DM diagnosis enhance the likelihood of achieving long-term remission. Taking into consideration the approval of new effective and safe medications for weight reduction, we propose a broader definition of T2DM remission, including individuals who achieve sustained glycemic control using glucose-lowering and weight management medications. For newly diagnosed T2DM patients, achieving an HbA1c below 6.5% should be the primary objective. Therefore, an aggressive, comprehensive approach is recommended, with patients fully informed about all available remission options, including potential medication combinations, right from the beginning of their diagnosis.

FOOTNOTES

Author contributions: Nakhleh A drafted the manuscript; Nakhleh A, Halfin E, and Shehadeh N edited and approved the final version of the manuscript.

Conflict-of-interest statement: The authors declare that there are no conflicts of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: Israel

ORCID number: Afif Nakhleh 0000-0003-1206-1516; Elya Halfin 0009-0000-3747-1971; Naim Shehadeh 0000-0003-3158-6254.

S-Editor: Chen YL

L-Editor: A

P-Editor: Zheng XM

REFERENCES

- 1 Chan TT, Tse YK, Lui RN, Wong GL, Chim AM, Kong AP, Woo J, Yeung DK, Abrigo JM, Chu WC, Wong VW, Tang RS. Fatty Pancreas Is Independently Associated With Subsequent Diabetes Mellitus Development: A 10-Year Prospective Cohort Study. *Clin Gastroenterol Hepatol* 2022; **20**: 2014-2022.e4 [PMID: 34571257 DOI: 10.1016/j.cgh.2021.09.027]
- 2 Lingvay I, Sumithran P, Cohen RV, le Roux CW. Obesity management as a primary treatment goal for type 2 diabetes: time to reframe the conversation. *Lancet* 2022; **399**: 394-405 [PMID: 34600604 DOI: 10.1016/S0140-6736(21)01919-X]
- 3 Riddle MC, Cefalu WT, Evans PH, Gerstein HC, Nauck MA, Oh WK, Rothberg AE, le Roux CW, Rubino F, Schauer P, Taylor R, Twenefour D. Consensus Report: Definition and Interpretation of Remission in Type 2 Diabetes. *J Clin Endocrinol Metab* 2022; **107**: 1-9 [PMID: 35111111 DOI: 10.1210/clinem.2022-01111]

- 34459898 DOI: [10.1210/clinem/dgab585](https://doi.org/10.1210/clinem/dgab585)]
- 4 **Lean ME**, Leslie WS, Barnes AC, Brosnahan N, Thom G, McCombie L, Peters C, Zhyzhneuskaya S, Al-Mrabeh A, Hollingsworth KG, Rodrigues AM, Rehackova L, Adamson AJ, Sniehotta FF, Mathers JC, Ross HM, McIlvenna Y, Stefanetti R, Trenell M, Welsh P, Kean S, Ford I, McConnachie A, Sattar N, Taylor R. Primary care-led weight management for remission of type 2 diabetes (DiRECT): an open-label, cluster-randomised trial. *Lancet* 2018; **391**: 541-551 [PMID: [29221645](https://pubmed.ncbi.nlm.nih.gov/29221645/) DOI: [10.1016/S0140-6736\(17\)33102-1](https://doi.org/10.1016/S0140-6736(17)33102-1)]
 - 5 **Taheri S**, Zaghoul H, Chagoury O, Elhadad S, Ahmed SH, El Khatib N, Amona RA, El Nahas K, Suleiman N, Alnaama A, Al-Hamaq A, Charlson M, Wells MT, Al-Abdulla S, Abou-Samra AB. Effect of intensive lifestyle intervention on bodyweight and glycaemia in early type 2 diabetes (DIADEM-I): an open-label, parallel-group, randomised controlled trial. *Lancet Diabetes Endocrinol* 2020; **8**: 477-489 [PMID: [32445735](https://pubmed.ncbi.nlm.nih.gov/32445735/) DOI: [10.1016/S2213-8587\(20\)30117-0](https://doi.org/10.1016/S2213-8587(20)30117-0)]
 - 6 **Hocking SL**, Markovic TP, Lee CMY, Picone TJ, Gudorf KE, Colagiuri S. Intensive Lifestyle Intervention for Remission of Early Type 2 Diabetes in Primary Care in Australia: DiRECT-Aus. *Diabetes Care* 2024; **47**: 66-70 [PMID: [37840461](https://pubmed.ncbi.nlm.nih.gov/37840461/) DOI: [10.2337/dc23-0781](https://doi.org/10.2337/dc23-0781)]
 - 7 **Gregg EW**, Chen H, Wagenknecht LE, Clark JM, Delahanty LM, Bantle J, Pownall HJ, Johnson KC, Safford MM, Kitabchi AE, Pi-Sunyer FX, Wing RR, Bertoni AG; Look AHEAD Research Group. Association of an intensive lifestyle intervention with remission of type 2 diabetes. *JAMA* 2012; **308**: 2489-2496 [PMID: [23288372](https://pubmed.ncbi.nlm.nih.gov/23288372/) DOI: [10.1001/jama.2012.67929](https://doi.org/10.1001/jama.2012.67929)]
 - 8 **Look AHEAD Research Group**, Gregg EW, Jakicic JM, Blackburn G, Bloomquist P, Bray GA, Clark JM, Coday M, Curtis JM, Egan C, Evans M, Foreyt J, Foster G, Hazuda HP, Hill JO, Horton ES, Hubbard VS, Jeffery RW, Johnson KC, Kitabchi AE, Knowler WC, Kriska A, Lang W, Lewis CE, Montez MG, Nathan DM, Neiberg RH, Patricio J, Peters A, Pi-Sunyer X, Pownall H, Redmon B, Regensteiner J, Rejeski J, Ribisl PM, Safford M, Stewart K, Trencle D, Wadden TA, Wing RR, Yanovski SZ. Association of the magnitude of weight loss and changes in physical fitness with long-term cardiovascular disease outcomes in overweight or obese people with type 2 diabetes: a post-hoc analysis of the Look AHEAD randomised clinical trial. *Lancet Diabetes Endocrinol* 2016; **4**: 913-921 [PMID: [27595918](https://pubmed.ncbi.nlm.nih.gov/27595918/) DOI: [10.1016/S2213-8587\(16\)30162-0](https://doi.org/10.1016/S2213-8587(16)30162-0)]
 - 9 **Gregg EW**, Chen H, Bancks MP, Manalac R, Maruthur N, Munshi M, Wing R; Look AHEAD Research Group. Impact of remission from type 2 diabetes on long-term health outcomes: findings from the Look AHEAD study. *Diabetologia* 2024; **67**: 459-469 [PMID: [38233592](https://pubmed.ncbi.nlm.nih.gov/38233592/) DOI: [10.1007/s00125-023-06048-6](https://doi.org/10.1007/s00125-023-06048-6)]
 - 10 **Al-Mrabeh A**, Hollingsworth KG, Shaw JAM, McConnachie A, Sattar N, Lean MEJ, Taylor R. 2-year remission of type 2 diabetes and pancreas morphology: a post-hoc analysis of the DiRECT open-label, cluster-randomised trial. *Lancet Diabetes Endocrinol* 2020; **8**: 939-948 [PMID: [33031736](https://pubmed.ncbi.nlm.nih.gov/33031736/) DOI: [10.1016/S2213-8587\(20\)30303-X](https://doi.org/10.1016/S2213-8587(20)30303-X)]
 - 11 **Yang X**, Zhou J, Shao H, Huang B, Kang X, Wu R, Bian F, Hu M, Liu D. Effect of an Intermittent Calorie-restricted Diet on Type 2 Diabetes Remission: A Randomized Controlled Trial. *J Clin Endocrinol Metab* 2023; **108**: 1415-1424 [PMID: [36515429](https://pubmed.ncbi.nlm.nih.gov/36515429/) DOI: [10.1210/clinem/dgac661](https://doi.org/10.1210/clinem/dgac661)]
 - 12 **Clina JG**, Sayer RD, Pan Z, Cohen CW, McDermott MT, Catenacci VA, Wyatt HR, Hill JO. High- and normal-protein diets improve body composition and glucose control in adults with type 2 diabetes: a randomized trial. *Obesity (Silver Spring)* 2023; **31**: 2021-2030 [PMID: [37475689](https://pubmed.ncbi.nlm.nih.gov/37475689/) DOI: [10.1002/oby.23815](https://doi.org/10.1002/oby.23815)]
 - 13 **Legaard GE**, Lyngbæk MPP, Almdal TP, Karstoft K, Bennetsen SL, Feineis CS, Nielsen NS, Durrer CG, Liebetrau B, Nystrup U, Østergaard M, Thomsen K, Trinh B, Solomon TPJ, Van Hall G, Brønd JC, Holst JJ, Hartmann B, Christensen R, Pedersen BK, Ried-Larsen M. Effects of different doses of exercise and diet-induced weight loss on beta-cell function in type 2 diabetes (DOSE-EX): a randomized clinical trial. *Nat Metab* 2023; **5**: 880-895 [PMID: [37127822](https://pubmed.ncbi.nlm.nih.gov/37127822/) DOI: [10.1038/s42255-023-00799-7](https://doi.org/10.1038/s42255-023-00799-7)]
 - 14 **Zou W**, Luo K, Hu Z, Zhang X, Feng C, Ye D, Liu S, Zhang Q, Tan R, Fu C. Diabetes remission after a lifestyle-medicine intervention on type 2 diabetes in lean and obese Chinese subjects: a prospective study. *Ann Palliat Med* 2022; **11**: 1462-1472 [PMID: [35523754](https://pubmed.ncbi.nlm.nih.gov/35523754/) DOI: [10.21037/apm-22-369](https://doi.org/10.21037/apm-22-369)]
 - 15 **Saisho Y**. β -cell dysfunction: Its critical role in prevention and management of type 2 diabetes. *World J Diabetes* 2015; **6**: 109-124 [PMID: [25685282](https://pubmed.ncbi.nlm.nih.gov/25685282/) DOI: [10.4239/wjd.v6.i1.109](https://doi.org/10.4239/wjd.v6.i1.109)]
 - 16 **Kramer CK**, Zinman B, Retnakaran R. Short-term intensive insulin therapy in type 2 diabetes mellitus: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol* 2013; **1**: 28-34 [PMID: [24622264](https://pubmed.ncbi.nlm.nih.gov/24622264/) DOI: [10.1016/S2213-8587\(13\)70006-8](https://doi.org/10.1016/S2213-8587(13)70006-8)]
 - 17 **Nakhleh A**, Shehadeh N. Hypoglycemia in diabetes: An update on pathophysiology, treatment, and prevention. *World J Diabetes* 2021; **12**: 2036-2049 [PMID: [35047118](https://pubmed.ncbi.nlm.nih.gov/35047118/) DOI: [10.4239/wjd.v12.i12.2036](https://doi.org/10.4239/wjd.v12.i12.2036)]
 - 18 **McInnes N**, Smith A, Otto R, Vandermey J, Punthakee Z, Sherifali D, Balasubramanian K, Hall S, Gerstein HC. Piloting a Remission Strategy in Type 2 Diabetes: Results of a Randomized Controlled Trial. *J Clin Endocrinol Metab* 2017; **102**: 1596-1605 [PMID: [28324049](https://pubmed.ncbi.nlm.nih.gov/28324049/) DOI: [10.1210/jc.2016-3373](https://doi.org/10.1210/jc.2016-3373)]
 - 19 **McInnes N**, Hall S, Sultan F, Aronson R, Hramiak I, Harris S, Sigal RJ, Woo V, Liu YY, Gerstein HC. Remission of Type 2 Diabetes Following a Short-term Intervention With Insulin Glargine, Metformin, and Dapagliflozin. *J Clin Endocrinol Metab* 2020; **105** [PMID: [32403130](https://pubmed.ncbi.nlm.nih.gov/32403130/) DOI: [10.1210/clinem/dgaa248](https://doi.org/10.1210/clinem/dgaa248)]
 - 20 **Nakhleh A**, Goldenberg-Furmanov M, Goldstein R, Shohat M, Shehadeh N. A beneficial role of GLP-1 receptor agonist therapy in ABCC8-MODY (MODY 12). *J Diabetes Complications* 2023; **37**: 108566 [PMID: [37536118](https://pubmed.ncbi.nlm.nih.gov/37536118/) DOI: [10.1016/j.jdiacomp.2023.108566](https://doi.org/10.1016/j.jdiacomp.2023.108566)]
 - 21 **Davies M**, Færch L, Jeppesen OK, Pakseresht A, Pedersen SD, Perreault L, Rosenstock J, Shimomura I, Viljoen A, Wadden TA, Lingvay I; STEP 2 Study Group. Semaglutide 2.4 mg once a week in adults with overweight or obesity, and type 2 diabetes (STEP 2): a randomised, double-blind, double-dummy, placebo-controlled, phase 3 trial. *Lancet* 2021; **397**: 971-984 [PMID: [33667417](https://pubmed.ncbi.nlm.nih.gov/33667417/) DOI: [10.1016/S0140-6736\(21\)00213-0](https://doi.org/10.1016/S0140-6736(21)00213-0)]
 - 22 **Del Prato S**, Kahn SE, Pavo I, Weerakkody GJ, Yang Z, Doupis J, Aizenberg D, Wynne AG, Riesmeyer JS, Heine RJ, Wiese RJ; SURPASS-4 Investigators. Tirzepatide versus insulin glargine in type 2 diabetes and increased cardiovascular risk (SURPASS-4): a randomised, open-label, parallel-group, multicentre, phase 3 trial. *Lancet* 2021; **398**: 1811-1824 [PMID: [34672967](https://pubmed.ncbi.nlm.nih.gov/34672967/) DOI: [10.1016/S0140-6736\(21\)02188-7](https://doi.org/10.1016/S0140-6736(21)02188-7)]
 - 23 **Jastreboff AM**, Aronne LJ, Ahmad NN, Wharton S, Connery L, Alves B, Kiyosue A, Zhang S, Liu B, Bunck MC, Stefanski A; SURMOUNT-1 Investigators. Tirzepatide Once Weekly for the Treatment of Obesity. *N Engl J Med* 2022; **387**: 205-216 [PMID: [35658024](https://pubmed.ncbi.nlm.nih.gov/35658024/) DOI: [10.1056/NEJMoa2206038](https://doi.org/10.1056/NEJMoa2206038)]
 - 24 **Rosenstock J**, Frias J, Jastreboff AM, Du Y, Lou J, Gurbuz S, Thomas MK, Hartman ML, Haupt A, Milicevic Z, Coskun T. Retatrutide, a GIP, GLP-1 and glucagon receptor agonist, for people with type 2 diabetes: a randomised, double-blind, placebo and active-controlled, parallel-group, phase 2 trial conducted in the USA. *Lancet* 2023; **402**: 529-544 [PMID: [37385280](https://pubmed.ncbi.nlm.nih.gov/37385280/) DOI: [10.1016/S0140-6736\(23\)01053-X](https://doi.org/10.1016/S0140-6736(23)01053-X)]
 - 25 **Ahqvist E**, Storm P, Käräjämäki A, Martinell M, Dorkhan M, Carlsson A, Vikman P, Prasad RB, Aly DM, Almgren P, Wessman Y, Shaat N, Spégel P, Mulder H, Lindholm E, Melander O, Hansson O, Malmqvist U, Lernmark Å, Lahti K, Forsén T, Tuomi T, Rosengren AH, Groop L. Novel subgroups of adult-onset diabetes and their association with outcomes: a data-driven cluster analysis of six variables. *Lancet Diabetes*

- Endocrinol* 2018; **6**: 361-369 [PMID: 29503172 DOI: 10.1016/S2213-8587(18)30051-2]
- 26 **Chumakova-Orin M**, Vanetta C, Moris DP, Guerron AD. Diabetes remission after bariatric surgery. *World J Diabetes* 2021; **12**: 1093-1101 [PMID: 34326957 DOI: 10.4239/wjd.v12.i7.1093]
- 27 **Jin ZL**, Liu W. Progress in treatment of type 2 diabetes by bariatric surgery. *World J Diabetes* 2021; **12**: 1187-1199 [PMID: 34512886 DOI: 10.4239/wjd.v12.i8.1187]
- 28 **McTigue KM**, Wellman R, Nauman E, Anau J, Coley RY, Odor A, Tice J, Coleman KJ, Courcoulas A, Pardee RE, Toh S, Janning CD, Williams N, Cook A, Sturtevant JL, Horgan C, Arterburn D; PCORnet Bariatric Study Collaborative. Comparing the 5-Year Diabetes Outcomes of Sleeve Gastrectomy and Gastric Bypass: The National Patient-Centered Clinical Research Network (PCORNet) Bariatric Study. *JAMA Surg* 2020; **155**: e200087 [PMID: 32129809 DOI: 10.1001/jamasurg.2020.0087]
- 29 **Schauer PR**, Bhatt DL, Kirwan JP, Wolski K, Aminian A, Brethauer SA, Navaneethan SD, Singh RP, Pothier CE, Nissen SE, Kashyap SR; STAMPEDE Investigators. Bariatric Surgery versus Intensive Medical Therapy for Diabetes - 5-Year Outcomes. *N Engl J Med* 2017; **376**: 641-651 [PMID: 28199805 DOI: 10.1056/NEJMoa1600869]
- 30 **Akkus G**, Tetiker T. Which predictors could effect on remission of type 2 diabetes mellitus after the metabolic surgery: A general perspective of current studies? *World J Diabetes* 2021; **12**: 1312-1324 [PMID: 34512896 DOI: 10.4239/wjd.v12.i8.1312]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

