Supplementary Table 1 Recording items of the eight scale items of SF-36 (step one)

Item number	Change	original	To recorded value
	response category ¹		
1, 2, 20, 22, 34, 36	1 →		100
	$2 \rightarrow$		75
	$3 \rightarrow$		50
	$4 \rightarrow$		25
	$5 \rightarrow$		0
3, 4, 5, 6, 7, 8, 9, 10, 11, 12	1 →		0
	$2 \rightarrow$		50
	$3 \rightarrow$		100
13, 14, 15, 16, 17, 18, 19	$1 \rightarrow$		0
	$2 \rightarrow$		100
21, 23, 26, 27, 30	$1 \rightarrow$		100
	$2 \rightarrow$		80
	$3 \rightarrow$		60
	$4 \rightarrow$		40
	$5 \rightarrow$		20
	$6 \rightarrow$		0
24, 25, 28, 29, 31	$1 \rightarrow$		0
	$2 \rightarrow$		20
	$3 \rightarrow$		40
	$4 \rightarrow$		60
	5 →		80
	6 →		100
32, 33, 35	1 →		0

$2 \rightarrow$	25
$3 \rightarrow$	50
$4 \rightarrow$	75
$5 \rightarrow$	100

 $^{^1\!}Pre\text{-coded}$ response choices as printed in the question naire.

Supplementary Table 2 Averaging items to form scales (step two)

Scale	Number of items	After recording per
		Table 2, average the
		following items
Physical functioning	10	3 4 5 6 7 8 9 10 11 12
Role limitations due to	4	13 14 15 16
physical health		
Role limitations due to	3	17 18 19
emotional problems		
Energy/ Fatigue	4	23 27 29 31
Emotional wellbeing	5	24 25 26 28 30
Social functioning	2	20 32
Pain	2	21 22
General health	5	1 33 34 35 36