

**8** **Name of Journal:** *World Journal of Clinical Cases*

**Manuscript NO:** 60136

**Manuscript Type:** ORIGINAL ARTICLE

**Retrospective Study**

**Risk factors associated with acute respiratory distress syndrome in COVID-19 Patients outside Wuhan-a double-center retrospective cohort study of 197 cases in Hunan, China**

Xing-Sheng Hu, Chun-Hong Hu, Ping Zhong, Ya-Jing Wen, Xiang-Yu Chen

## Match Overview

<b>1</b>	<b>Crossref</b> 205 words Xingsheng HU, Chunhong HU, Yong Yang, Juan Chen, Ping Zhong, Yajing Wen, Xiangyu Chen. "Clinical characteristics of COVID-19 patients with pneumonia in Wuhan, China: a retrospective study." <i>Journal of Clinical Medicine</i> . 2020;9(2):257-263. doi:10.3390/jcm9020257	3%
<b>2</b>	<b>Crossref</b> 88 words Xingsheng HU, Chunhong HU, Ping Zhong, Yajing Wen, Yong Yang, Juan Chen, Xiangyu Chen. "Finding the Best Antiviral Drug for COVID-19: A Retrospective Cohort Study." <i>Journal of Clinical Medicine</i> . 2020;9(2):257-263. doi:10.3390/jcm9020257	1%
<b>3</b>	<b>Internet</b> 44 words crawled on 03-Oct-2020 <a href="http://www.researchsquare.com">www.researchsquare.com</a>	1%
<b>4</b>	<b>Internet</b> 30 words crawled on 26-May-2020 <a href="http://journals.physiology.org">journals.physiology.org</a>	<1%
<b>5</b>	<b>Internet</b> 26 words crawled on 02-Nov-2020 <a href="http://www.mdpi.com">www.mdpi.com</a>	<1%
<b>6</b>	<b>Internet</b> 25 words crawled on 20-Sep-2020	<1%

国内版

国际版

Risk Factors Associated with ARDS in Patients with COVID-19 out



ALL

IMAGES

VIDEOS

There are no results for **Risk Factors Associated with ARDS in Patients with COVID-19 outside Wuhan -a double-center retrospective cohort study of 197 cases in Hunan, China**

Check your spelling or try different keywords

Ref A: 0A82347DE06445EEA662F2085496E20E Ref B: BJ1EDGE0512 Ref C: 2020-10-22T09:13:33Z



国内版

国际版

Risk factors associated with acute respiratory distress syndrome in COVI



网页

图片

视频

学术

词典

地图

检测到您输入了英文，试试切换到国际版？搜英文结果更丰富更准确 >

415,000 条结果

时间不限 ▾

## [Risk Factors Associated With Acute Respiratory ...](#) 翻译此页

Cited by: 2451

Author: Chaomin Wu, Xiaoyan Chen, Yanping Cai,...

Publish Year: 2020

Importance: **Coronavirus** disease 2019 (**COVID-19**) is an emerging infectious disease that was first reported in **Wuhan**, China, and has subsequently spread worldwide. **Risk factors** for the clinical outcomes of **COVID-19** pneumonia have not yet been well delineated. Objective: To describe the clinical characteristics and outcomes in **patients** with **COVID-19** pneumonia who developed **acute respiratory ...**

<https://pubmed.ncbi.nlm.nih.gov/32167524>

## [Risk Factors Associated With Acute Respiratory ...](#) 翻译此页

Cited by: 2451

Author: Chaomin Wu, Xiaoyan Chen, Yanping Cai,...

Publish Year: 2020

2020-7-1 · **Patients** with severe illness may develop dyspnea and hypoxemia within 1 week after onset of the disease, which may quickly progress to **acute respiratory distress syndrome** (ARDS) or end-organ failure. 4 Certain epidemiological features and clinical characteristics of **COVID-19** have been previously reported. 3-5 However, these studies were based ...

<https://jamanetwork.com/journals/jamainternalmedicine/fullarticle/2763184> ▾

## [Risk Factors Associated With Acute Respiratory ...](#) 翻译此页

2020-3-13 · **Risk Factors Associated With Acute Respiratory Distress Syndrome** and Death in **Patients** With **Coronavirus** Disease 2019 Pneumonia in **Wuhan**, China March 2020 JAMA Internal Medicine 180(7)

[https://www.researchgate.net/publication/339918092\\_Risk\\_Factors\\_Associated\\_With\\_Acute...](https://www.researchgate.net/publication/339918092_Risk_Factors_Associated_With_Acute...)

## [Acute Respiratory Distress Syndrome \(ARDS\) in COVID ...](#) 翻译此页

2020 4 2 · **Acute respiratory distress syndrome** (ARDS) can develop in **patients** with **COVID-19** if the





ALL

IMAGES

VIDEOS

14,100 Results

Any time ▾

## Risk Factors Associated With Acute Respiratory Distress ...

<https://jamanetwork.com/journals/jamainternalmedicine/fullarticle/2763184> ▾

Jul 01, 2020 · **Patients** with severe illness may develop dyspnea and hypoxemia within 1 week after onset of the disease, which may quickly progress to **acute respiratory distress syndrome** (ARDS) or end-organ failure. 4 Certain epidemiological features and clinical characteristics of **COVID-19** have been previously reported. 3-5 However, these studies were based ...

**Cited by:** 2451

**Author:** Chaomin Wu, Xiaoyan Chen, Yanping Ca...

**Publish Year:** 2020

## Clinical characteristics and risk factors for severity of ...

<https://journals.sagepub.com/doi/full/10.1177/1753466620963035>

**Aim:**To investigate clinical characteristics and identify **risk factors** for severity of coronavirus disease 2019 (**COVID-19**) pneumonia **outside** of Wuhan, **China**. **Materials and methods:**We included 213 pat...

## Severe Covid-19 | NEJM

<https://www.nejm.org/doi/full/10.1056/NEJMcp2009575>

May 15, 2020 · **Patients** with severe **Covid-19** commonly meet the criteria for the **acute respiratory distress syndrome** (ARDS), which is defined as the **acute** onset of bilateral infiltrates, severe hypoxemia, and ...

**Cited by:** 103

**Author:** David A Berlin, Roy M Gulick, Fernando J...

**Publish Year:** 2020

## Epidemiological characteristics of patients with severe ...

<https://www.docwirenews.com/abstracts/...> ▾

2 days ago · **ABSTRACT. BACKGROUND:** The new coronavirus (**COVID-19**) rapidly resulted in a pandemic. We report the characteristics of **patients** with severe or critical severe **acute respiratory syndrome** coronavirus 2 (SARS-CoV-2) infection in Wuhan city, **China**, and the **risk factors** related to infection severity and death.