

Tick-borne encephalitis: A review of epidemiology, clinical characteristics, and management

Petra Bogovic, Franc Strle

Petra Bogovic, Franc Strle, Department of Infectious Diseases, University Medical Center Ljubljana, 1525 Ljubljana, Slovenia

Author contributions: Both authors approved the final version of the manuscript before submission.

Conflict-of-interest: The authors declare that they have no competing interests.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Correspondence to: Petra Bogovic, MD, Department of Infectious Diseases, University Medical Center Ljubljana, Japljeva 2, 1525 Ljubljana, Slovenia. petra.bogovic@kclj.si

Telephone: +386-1-5222110

Fax: +386-1-5222456

Received: July 23, 2014

Peer-review started: July 25, 2014

First decision: August 28, 2014

Revised: September 18, 2014

Accepted: March 5, 2015

Article in press: March 9, 2015

Published online: May 16, 2015

abortive form of the disease and a chronic progressive form. A post-encephalitic syndrome, causing long-lasting morbidity that often affects the quality of life develops in up to 50% of patients after acute tick-borne encephalitis. Clinical course and outcome vary by subtype of tick-borne encephalitis virus (the disease caused by the European subtype has milder course and better outcome than the disease caused by Siberian and Far-Eastern subtypes), age of patients (increasing age is associated with less favorable outcome), and host genetic factors. Since clinical features and laboratory results of blood and cerebrospinal fluid are nonspecific, the diagnosis must be confirmed by microbiologic findings. The routine laboratory confirmation of the tick-borne encephalitis virus infection is based mainly on the detection of specific IgM and IgG antibodies in serum (and cerebrospinal fluid), usually by enzyme-linked immunosorbent assay. There is no specific antiviral treatment for tick-borne encephalitis. Vaccination can effectively prevent the disease and is indicated for persons living in or visiting tick-borne encephalitis endemic areas.

Key words: Tick-borne encephalitis; Diagnosis; Epidemiology; Clinical manifestations; Treatment; Prevention/vaccination

© **The Author(s) 2015.** Published by Baishideng Publishing Group Inc. All rights reserved.

Abstract

Tick-borne encephalitis is an infection of central nervous system caused by tick-borne encephalitis virus transmitted to humans predominantly by tick bites. During the last few decades the incidence of the disease has been increasing and poses a growing health problem in almost all endemic European and Asian countries. Most cases occur during the highest period of tick activity, in Central Europe mainly from April to November. Tick-borne encephalitis is more common in adults than in children. Clinical spectrum of the disease ranges from mild meningitis to severe meningoencephalitis with or without paralysis. Rare clinical manifestations are an

Core tip: Tick-borne encephalitis (TBE) is the most common tick-borne central nervous system infection in Europe and Asia. It is caused by three subtypes of TBE virus: European, Siberian and Far-Eastern. Because of relatively severe clinical course, the absence of etiologic treatment, considerable proportion of patients with incomplete recovery after acute illness, as well as due to increasing incidence it represents a growing public health problem that could be substantially reduced with vaccination.

Bogovic P, Strle F. Tick-borne encephalitis: A review of

epidemiology, clinical characteristics, and management. *World J Clin Cases* 2015; 3(5): 430-441 Available from: URL: <http://www.wjgnet.com/2307-8960/full/v3/i5/430.htm> DOI: <http://dx.doi.org/10.12998/wjcc.v3.i5.430>

INTRODUCTION

Tick-borne encephalitis (TBE) is an important viral infection of the central nervous system in Europe and in several regions in Asia. It is caused by the Far-Eastern, Siberian and European subtype of tick-borne encephalitis virus (TBEV). Although the disease is preventable by vaccination, the incidence has been increasing during the last few decades and consequently represents a growing health problem in almost all endemic European and Asian countries. With the increase of tourism TBE is becoming a problem also outside endemic regions^[1].

Most cases occur during the highest period of tick activity, in Central Europe mainly from April to November^[2-4]. TBE caused by the European virus subtype usually presents as meningitis or meningoencephalitis and has a biphasic course in approximately two-thirds of patients. Up to 50% of patients have long-term sequelae^[4]. An occasional clinical manifestation is an abortive form of the disease^[5]. Treatment is based on the symptomatic measures. However, TBE can be successfully prevented by safe and highly effective vaccine^[4].

ETIOLOGY

The neurotropic TBEV was first described as the cause of TBE by Zilber more than 75 years ago^[6]. It is a spherical lipid-enveloped RNA virus, a member of genus *Flavivirus* within the *Flaviviridae* family. The mature virion is composed of 3 structural proteins-capsid (C), membrane (M), and envelope (E). Protein E is a major antigen which induces production of neutralizing antibodies. It can be inactivated by pasteurization^[4].

The genetic analysis shows the existence of three TBEV subtypes named as European, Siberian, and Far-Eastern subtype^[2]. They are genetically very closely related; variation in amino acids sequences between subtypes is 5%-6%^[7]. In spite of the pronounced genetic similarity of the subtypes the illness caused by individual subtype is not completely equivalent to those due to the other subtypes.

EPIDEMIOLOGY

TBE is endemic in Europe, Siberia, far-eastern Russia, northern China and Japan. During the past few decades endemic regions have expanded, and within many endemic areas the number of reported cases increased. The increase in the reported incidence rates is thought to be a result of complex interplay of social and ecological factors as well as due to increased medical

awareness and advanced diagnostics^[8,9].

The European TBEV subtype is predominantly found in Europe, but has also been identified in west Urals, and in Siberia, whereas the Siberian TBEV subtype is found in Siberia, the Baltics, and northern Finland. The Far-Eastern TBE virus subtype is endemic in far-eastern Asia and Japan, and has been identified also in central and eastern Siberia^[1,2,10,11].

In Europe and Asia between 10000 and 15000 TBE cases are reported annually^[12]. The number is very likely underestimated because in many countries notification of the disease is not mandatory and only in a subset of the countries TBE case definition is in place. TBE is endemic in 27 European countries, and is a reportable disease in only 16 countries^[1,12,13]. European countries with the highest incidence of the disease in the period 2005-2009 were Slovenia (14.1 cases per 100000 inhabitants per year), Estonia (11.1), Lithuania (10.6), and Latvia (8.8)^[1]. Pronounced yearly variations of registered TBE cases occurred. According to the latest available epidemiological data for Slovenia the incidence of the disease in 2013 was 15.0 cases per 100000 inhabitants^[14].

The primary reservoirs and hosts of TBEV in nature are small rodents; humans do not play any role in the maintenance of TBEV in nature and they are only accidental hosts. TBEV is transmitted to humans mainly by hard tick bites; in Europe the principal vector is *Ixodes ricinus* (*I. ricinus*), in parts of Eastern Europe, Russia and in far-east Asia the vector is *Ixodes persulcatus* (*I. persulcatus*) whereas in Japan Far-Eastern TBEV subtype has been demonstrated in *Ixodes ovatus* ticks. In endemic areas in Central Europe approximately 0.1% to 5.0% of ticks harbor the virus (depending on the time of the year and geographical location); in Siberia infection rates of up to 40% are reported for *I. persulcatus*^[4,15]. In Slovenia, the prevalence of ticks infected with TBEV was found to be 0.47%; 0.54% in 2005 and 0.43% in 2006^[16]. Approximately 1% of all TBEV infections in humans are probably acquired by consuming infected unpasteurized milk or milk products from infected livestock, particularly goats^[2]. This means of transmission has to be considered particularly in cases of local epidemics. Outbreaks due to oral virus transmission are more common in Eastern Europe and the Baltic states than in Central Europe^[17,18]. A few cases of laboratory-acquired TBEV infections have been documented in the literature^[19].

TBE cases usually happen in the warm months between April and November, which is also the period of the highest tick activity^[20]. In Central Europe, where the dominant tick species is *I. ricinus*, a two-peak distribution of TBE cases can be seen (first in June and July, second in September and October), whereas in the Ural region, Siberia and the Far East, where *I. persulcatus* is widespread, cases as a rule occur in May and June^[21]. In all age groups men are affected more frequently than women^[14,22-24]. On average, 10%-20%

of all reported cases of TBE occur in children^[25]. It should be pointed out that due to its unspecific clinical presentation TBE in children is often missed and is diagnosed as aseptic meningitis of unknown etiology^[26].

With increasing of tourism, TBE has become a more global problem. Therefore, it should be included in the differential diagnosis of the central nervous system infections not only for those living within an endemic region but-in case of an appropriate epidemiological history-also in patients living outside endemic areas. The risk of travel-associated TBE depends on the season of travel, degree of unprotected outdoor exposure as well as on consuming unpasteurized dairy products. TBE surveillance data available in Austria shows that an overall risk of acquiring TBE for a non-vaccinated tourist, staying in a highly endemic region for 4 wk during the TBEV transmission season, has been estimated at approximately 1 case per 10000 person-months of exposure, which is approximately equivalent to the risk of contracting typhoid fever or malaria while traveling in India^[27,28].

PATHOGENESIS AND PATHOLOGY

After an infected tick bite TBEV replication occurs locally. Dendritic skin cells (Langerhans cells) are assumed to be the first cells for viral replication and to transport the virus to local lymph nodes. From this initial site the TBEV than disseminate to extraneural tissues, especially spleen, liver and bone marrow, where further multiplication maintains viremia for several days. During the viremic phase (which clinically corresponds to the initial phase of TBE) the virus probably reaches the brain^[29,30]. The exact mechanism by which TBEV breach the blood-brain barrier is not known; four possible routes have been postulated: (1) peripheral nerves; (2) highly susceptible olfactory neurons; (3) transcytosis through vascular endothelial cells of brain capillaries; and (4) diffusion of the virus between capillary endothelial cells. The primary targets of TBEV infection in central nervous system are neurons^[31].

According to rather limited information the neuropathological findings are nonspecific. Cerebral and spinal meninges usually show diffuse infiltration with lymphocytes and sometimes neutrophils. The most extensive meningeal inflammation is in the vicinity of the cerebellum. Pathological lesions which consist of lymphocytic perivascular infiltrations, accumulation of glial cells, nerve cells necrosis, and neuronophagia are localized in the grey matter and are most often present in the medulla oblongata, pons, cerebellum, brainstem, basal ganglia, thalamus, and spinal cord. Rarely, oligodendrocytes are infected. In the motor area of the cerebral cortex degeneration and necrosis of the pyramidal cells, lymphocytic accumulation, and glial proliferation are present^[29,32].

MANIFESTATIONS OF TBEV INFECTION

The large majority of infections with TBEV are

asymptomatic; published data suggest that the ratio of asymptomatic infections is between 70% and 98%^[4,33]. However, the proportion of asymptomatic cases is hard to ascertain because patients with mild clinical signs and symptoms may remain undiagnosed.

The incubation period of TBE ranges from 2 to 28 d and is usually 7-14 d. After alimentary TBEV transmission the incubation period is as a rule shorter, usually 3 to 4 d^[20]. Investigation of a recent small outbreak of TBE after drinking raw goat milk infected with TBEV in Slovenia revealed infection with the virus in four out of four exposed individuals. Of them, three developed symptomatic infection 2 to 3 d after the milk consumption, while the one who had been vaccinated against TBE remained healthy^[18].

In about 75% of patients with TBE due to the European TBEV subtype the disease has a typical biphasic course. The majority of patients with monophasic course of the disease has central nervous system involvement (meningitis, meningoencephalitis), while a small fraction has a febrile illness with headache but no meningitis (*i.e.*, the initial phase of TBE not followed by the second, meningoencephalitic phase of the disease), named abortive form of TBE or "febrile headache"^[4,5,20,34].

The initial phase correlate with viremia and usually presents with non-specific symptoms such as moderate fever, headache, body pain (myalgia and arthralgia), fatigue, general malaise, anorexia, nausea, and others^[3]. This phase lasts for 2 to 7 d and is followed by amelioration or even an asymptomatic interval that usually lasts for about 1 wk (1-21 d). Than the second phase appears: in approximately 50% of adult patients it presents as meningitis, in about 40% as meningoencephalitis, and in around 10% as meningoencephalomyelitis^[4,34].

Meningitis, encephalitis, myelitis

Meningitis and encephalitis are the most frequent clinical forms of TBE. Meningitis typically manifests with high fever, headache, nausea and vomiting; many patients have photophobia, and some vertigo. Meningeal signs are present in most of patients. Encephalitis can be manifested by impaired consciousness ranging from somnolence to stupor and, in rare cases, coma. Other manifestations comprise personality changes, behavioral disorders, concentration and cognitive function disturbances, tongue fasciculations and tremor of extremities; very rarely focal or generalized seizures, delirium and psychosis develop. Flaccid pareses, that are a typical characteristic of meningoencephalomyelitis, usually arise during the febrile phase of the disease, and are occasionally preceded by severe pain in the affected muscle groups. The upper extremities are more often affected than the lower extremities and the proximal segments more frequently than the distal ones. Patients with pareses of respiratory muscles rather commonly require artificial ventilatory support. Involvement of the central portions of the brainstem and medulla oblongata are associated with poor prognosis. Myelitis usually occur with encephalitis, and only very rarely as the only manifestation of TBE^[4,20,35,36].

In patients with TBE the involvement of cranial nerves has been reported. Published data suggests that cranial nerve involvement is rare and mainly asymmetrical, that its occurrence varies with the severity of clinical presentation of TBE, and that in most cases it has a favorable outcome^[24,37,38]. Cranial neuritis most commonly affects ocular, facial, pharyngeal and vestibular nerves^[4,20]. Of 1218 adult patients with TBE, who were hospitalized at the Department of Infectious Diseases, University Medical Centre Ljubljana, Slovenia from 2003 to 2009, 11 (0.9%) developed peripheral facial palsy during the course of the disease (nine had unilateral and two had bilateral facial nerve involvement; no one had a central facial palsy) and in three of them TBE was associated with borrelia infection^[39]. In regions where TBE and Lyme borreliosis are endemic, concomitant infection with TBEV and *Borrelia burgdorferi* sensu lato should be considered.

Occasionally, patients with TBE have pronounced variability in heart rate or other signs of autonomic nervous system dysfunction^[40].

Abortive form of TBE

Data on this manifestation of TBE are limited. It manifests with moderate fever, headache, fatigue, and other symptoms of initial phase of the disease that are not followed by nervous system involvement. The fever typically endures for several days, and the outcome of the disease is excellent^[41,42]. In Central Europe the majority of patients with the initial phase of TBE develop the second, central nervous system phase of the disease. In 2002, Lotric-Furlan *et al*^[5] published a prospective clinical study on the etiology of febrile illnesses after a tick bite. Among 56 patients with confirmed TBE, only one (2%) had an isolated initial phase, whereas all the others (55, 98%) diagnosed at the time of the initial phase developed the second phase of the disease with pleocytosis.

In Russia this clinical manifestation is named "fever form," and is reported to represent up to 50% of all clinical presentations of TBE^[43].

TBE with normal cerebrospinal fluid cell counts

A patient with encephalitis and serologically confirmed TBEV infection but without CSF pleocytosis has been reported^[44]. In larger studies of serologically proven TBE, CSF pleocytosis was found in all patients^[3,4,24,33]. This finding however, might be due to selection bias because in these studies CSF pleocytosis was one of the inclusion criteria for the diagnosis of TBE.

Chronic progressive form of TBE

There is no agreement on the existence of chronic TBE. Cases of a chronic progressive form of TBE have been identified in Siberia and the Russian Far East. This form of TBE is believed to be caused by the Siberian TBEV subtype. Both mutation in the *TBEV NS1* gene as well as an inappropriate T-cell immune response have been implicated to be associated with chronic progressive

disease^[3]. According to information from Western Siberia 1.7% of patients with acute TBE develop a chronic progressive form of the disease^[45]. Clinical presentations include Kozshevnikov's epilepsy, lateral sclerosis, progressive neuritis, progressive muscle atrophy, and a Parkinson-like disease. Pronounced dissimilarities in the incubation period, in time to the onset of individual neurological signs/symptoms, and in the survival time after the onset of the disease have been reported^[2]. Nevertheless, progressive form of TBE is most probably not present or is extremely unusual in disease caused by European TBEV subtype. In the study carried out in Lithuania, where both, European and Siberian TBEV subtypes are present, progressive course was noted in two out of 133 consecutive patients with acute TBE^[24].

Post-encephalitic syndrome

TBE may cause long-lasting morbidity which often has an impact on patients' quality of life and, sometimes, necessitates an alteration of lifestyle. Many nonspecific neurological/neuropsychiatric symptoms and residual neurological dysfunctions have been reported in some prospective and several retrospective studies, but findings are hard to compare due to diverse study designs, distinct definitions, and variable follow-up times. Most studies also failed to comprise a control group; as a result findings are difficult to interpret because of unclear distinctions between post-encephalitic syndrome, other sequelae of TBE, and symptoms present in general population. Published data suggest that 40% to 50% of patients after acute TBE develop a post-encephalitic syndrome^[46]. The most frequently reported symptoms have been cognitive disorders, neuropsychiatric complaints (such as apathy, irritability, memory and concentration disorders, altered sleep pattern), headache, hearing loss and/or tinnitus, disturbances of vision, balance and coordination disorders, and flaccid paresis or paralysis^[24,37,38,47-49].

Lithuanian prospective clinical follow-up study showed that 46% of patients with TBE had sequelae 1 year after the onset of acute illness^[24]. In a review by Haglund and Günther four retrospective and four prospective studies on the long-term morbidity from TBE were included. Up to 46% patients suffered permanent sequelae at long-term follow-up^[47]. In 2009, Misić Majerus *et al*^[48] published a prospective study on TBE post-encephalitic syndrome. Of 124 patients aged 16-76 years, who were followed for at least 3 years, 49 (40%) developed moderate or severe post-encephalitic syndrome lasting for up to 18 mo. Permanent sequelae were seen in 14 (11.3%) patients: spinal nerve paresis in 5, hearing impairment in 6, dysarthria in 2, and severe mental disorder in 1 patient.

CLINICAL COURSE OF ACUTE DISEASE AND LONG TERM OUTCOME

Course of acute TBE and its long-term outcome depend upon the subtype of TBEV infection. The disease caused

by the European TBEV subtype usually has a biphasic course, with a severe neurologic deficit in approximately 10% of patients, and a case-fatality rate of less than 2%^[4,20]. The abortive form of TBE is rare—the initial phase most of the time move on to the second phase of the disease^[5]. Long-lasting sequelae are identified in up to 50% of adult patients^[46]. Infections with Far Eastern TBEV subtype often cause an illness with a gradual onset, more severe course, higher rates of severe neurologic sequelae, and a fatality rate of 20%–40%. Little information with a respect to the clinical course of the disease is available for Siberian TBEV subtype. The case-fatality rate is 2%–3% and some reports from Russia suggest an association with a chronic progressive form of TBE^[2,21].

Published data suggest the relationship between age of patients and severity of TBE. The disease caused by European TBEV generally has milder course and better outcome in children than in adults. Nevertheless, TBE cases with severe clinical course, permanent sequelae and even death have been described not only in adults but also in children^[23,50]. The predominant form of TBE in children and adolescents is meningitis. A summary of 8 studies on 1169 children with TBE, published from 1963 to 2005, showed that meningitis was present in 802 (69%), meningoencephalitis in 356 (30%), and meningoencephalomyelitis in 11 (1%) patients. Twenty out of 945 patients (2.1%) had long-term neurologic sequelae^[25]. Limited information is available on the relationship between age and severity of TBE in adults. Mickiene *et al.*^[24] reported about this correlation but no precise data were given. Comparison of patients over an under 60 years revealed several differences in the course and outcome of TBE and corroborated previous postulation that TBE is a more severe disease in the elderly^[22].

Some clinical studies have shown that TBE with monophasic presentation is associated with a more severe course of the acute disease^[3,51–55].

The comparison of clinical course of TBE in unvaccinated and vaccinated patients did not reveal significant differences in disease severity^[56].

The outcome of TBE is associated with clinical presentation. The risk of incomplete recovery is higher for patients who have more severe clinical illness during acute phase of TBE^[24].

In recent years, genetic factors with potential impact on the course and outcome of TBE have been of scientific interest, resulting in several investigations. For example, the assessment of the role of chemokine receptor CCR5 indicated that CCR5Δ23 allele may predispose for TBE^[57]. Barkash *et al.*^[58,59] reported on the association between CD209 gene promoter region polymorphism and predisposition to severe forms of TBE, and on possible association between 5 OAS single nucleotide polymorphisms and the outcome of TBEV infection in a Russian population. Interesting new findings on the role of host genetic factors in TBEV infections may appear in future.

DIAGNOSIS

A case of TBE is delineated by the presence of: (1) symptoms/signs indicating meningitis or meningoencephalitis; (2) an elevated cerebrospinal fluid cell count ($> 5 \times 10^6$ cells/L); and (3) microbiologic evidence of TBEV infection (the presence of specific IgM and IgG antibodies)^[60].

Blood and cerebrospinal fluid analysis

In patients with TBE blood and cerebrospinal fluid findings are nonspecific. In the first (viremic) phase of TBE leukopenia and/or thrombocytopenia is established in approximately 70% of patients, rarely abnormal liver function test results are seen^[61]. In the second phase of the disease mildly elevated leukocyte count may be present in peripheral blood (rarely $> 15 \times 10^9$ /L); erythrocyte sedimentation rate and concentration of C-reactive protein are normal in the majority of patients but may be elevated, particularly in some long-lasting severe cases. Cerebrospinal fluid examination typically reveals elevated leukocyte counts (usually lower than 500 cells/mm³), a normal glucose concentration, and a normal to slightly elevated protein concentration. Early in the course of the disease neutrophils may predominate, while later cerebrospinal fluid profile is characterized by a predominance of lymphocytes. Elevated lymphocyte counts may last for several weeks after clinical improvement^[20,62].

Microbiological investigations

At the time when neurological symptoms/signs occur TBEV has already been cleared from the blood (TBEV is present in blood in the initial but not in the meningoencephalitic phase of the disease) and is only very exceptionally present in cerebrospinal fluid. Consequently, isolation of TBEV from blood and detection of viral RNA by reverse transcriptase PCR in blood and cerebrospinal fluid of patients with TBE have a limited diagnostic yield and are as a rule not used in clinical practice^[63]. Reverse transcriptase PCR assays is mainly limited to the initial phase of the disease and could be a useful method in a diagnostic procedure of febrile illness occurring after a tick bite in areas where several tick-borne diseases are present^[64].

The routine laboratory confirmation of the TBEV infection is based mainly on the demonstration of specific antibodies in serum (and cerebrospinal fluid), usually by highly sensitive and specific enzyme-linked immunosorbent assay^[63,65]. In the majority of patients specific serum IgM and IgG antibodies are present at the beginning of the meningoencephalitic phase of the disease; rarely only IgM antibodies to TBEV are found in the first serum sample. In such cases a second serum sample has to be tested 1–2 wk later, because the demonstration of IgM antibodies alone does not suffice for the diagnosis. TBEV IgM antibodies can be detected in the serum for several months (up to 10 mo or even longer) after acute infection, whereas TBEV

IgG antibodies persist for a whole life, and mediate an immunity that prevents symptomatic reinfection^[63,66]. In cerebrospinal fluid specific IgM and IgG antibodies are detectable several days later than in serum, and in almost all cases by day 10^[63,67].

However, some limitations are necessary to take into account when using and interpreting serological testing. Specific TBE IgM antibodies may be detectable for several months after acute TBEV infection (as well as in some persons after the first two doses of primary immunization) and may lead to erroneous interpretation in case of another central nervous system disease/infection within this time period^[63,66]. Because of the close antigenic relationship between TBEV and other flaviviruses cross-reactive antibodies are induced by infections or vaccinations. This may pose a diagnostic challenge in people vaccinated against yellow fever or Japanese encephalitis and in travelers having acquired dengue, West Nile or other flavivirus infections^[68]. Such potential problems in TBE serodiagnosis can be resolved by the quantification of IgM antibodies in a single serum sample taken at the time of hospitalization. High IgM values (> 500 Arbitrary Units) are indicative of a recent infection with TBEV. Lower IgM values, however, may require the analysis of a follow-up sample and/or a specific neutralization assay to exclude the possibilities of long persisting IgM antibodies after infection or those induced by vaccination as well as cross reactive IgM antibodies^[69]. Carefulness in the interpretation of TBE serology is also needed in patients with meningoencephalitis or meningitis who had been vaccinated against TBE. In the majority of patients with vaccination breakthroughs serological response is distinct from those found in unvaccinated persons with TBE and consequently vaccination breakthrough cases may be overlooked. These cases are characterized by a delayed development of specific IgM response (during the initial week of the meningoencephalitic phase of TBE specific IgM antibodies are usually not detectable) associated with a rapid increase of specific serum IgG antibodies^[70-72]. For a reliable diagnosis of TBE in persons previously vaccinated against TBE, demonstration of intrathecal production of TBEV antibodies is required^[34].

DIFFERENTIAL DIAGNOSIS

The differential diagnosis of TBE is extensive and includes a wide variety of central nervous system infections due to other infectious agents as well as noninfectious diseases.

In the initial, viremic phase of TBE, when a patient present with fever, headache, arthralgia, myalgia, and malaise the differential diagnosis may include various viral syndromes; if nausea, vomiting, diarrhea, and anorexia are present, gastroenteritis is a possible explanation. When signs and symptoms of central nervous system involvement develop, TBE needs to be differentiated from encephalitis or aseptic meningitis due

to many other viruses. Differential diagnosis comprises also other tick-borne diseases such as Lyme borreliosis, babesiosis, human granulocytic anaplasmosis, tick-transmitted rickettsioses, and tularemia^[20].

In many endemic regions TBEV geographically overlaps with bacterial tick-borne pathogens such as *Anaplasma phagocytophilum* and *B. burgdorferi* s.l. Both diseases caused by the two agents, human granulocytic anaplasmosis and Lyme borreliosis, are treatable with antibiotics, and caution must be taken to distinguish them from TBE^[20]. Concomitant TBEV and *B. burgdorferi* s.l. infections, as well as concomitant TBEV and *A. phagocytophilum* infections have been described^[60,73-76].

The initial, viremic phase of TBE and human granulocytic anaplasmosis have a similar clinical and laboratory presentation. Both diseases are characterized by fever and headache, and both are typically associated with thrombocytopenia and leukopenia; in certain patients also mild abnormalities in liver tests are present. Nevertheless, a clinical report on chills, myalgia and arthralgia, and laboratory findings of elevated concentration of C-reactive protein and lactate dehydrogenase values point to the diagnosis of human granulocytic anaplasmosis and not in favor of the initial phase of TBE^[77].

TREATMENT

There is no specific antiviral treatment for TBE. Patients as a rule need hospitalization and supportive care based on the severity of signs/symptoms, and usually encompasses administration of antipyretics, analgesics, antiemetics, maintenance of water and electrolyte balance, and if necessary administration of anticonvulsive agents. In patients with neuromuscular paralysis leading to respiratory failure, intubation and ventilatory support are necessary. In a large prospective German study, 12% of patients were treated in intensive care unit and in 5% assisted ventilation was required^[38]. Among Slovenian patients with TBE, hospitalized at the Department of Infectious Diseases, University Medical Centre Ljubljana, in years 2000 to 2004, 6.9% were hospitalized in intensive care unit and 22.5% of them needed mechanical ventilation^[22].

Cerebral edema is a potential complication of acute viral encephalitis that aggravates the clinical picture and portends poor neurologic outcome. Patients who have significantly raised intracranial pressure are often treated with intravenous mannitol and/or steroids^[78]. Mannitol produces a shift of fluid from the edematous brain back into the intravascular space and subsequently increases circulation volume and improves cerebral perfusion pressure, and lowers intracranial pressure through cerebral autoregulation. It also acts on erythrocyte membrane fluidity and through reducing blood viscosity improves blood flow and oxygen delivery. A rebound phenomenon in intracranial hypertension occurs in about 5% of patients. To avoid complications,

it is frequently recommended that mannitol no longer be administered when the serum osmolality exceeds 320 mOsm/L^[78,79]. Despite a rather common clinical practice to treat patients who have significantly raised intracranial pressure with intravenous mannitol, there are no reliable (comparative) studies substantiating the usage of mannitol in patients with TBE.

It has been demonstrated that the use of dexamethasone results in a reduction of brain edema in acute viral encephalitis^[80]. In the Baltics and some Eastern European countries patients with TBE are quite common given corticosteroids. The use of steroids is based on the impression that they produce a good and rapid clinical response in patients with TBE, but the validation of the impression in controlled studies has been limited. In a Lithuanian study corticosteroids were used in 81 out of the 133 patients with TBE. Among patients with mild, moderate, and severe disease, 39.7%, 70.7%, and 100% received corticosteroids, respectively. Hospitalization was significantly prolonged among these patients, compared with patients who received only symptomatic treatment. Because all patients with severe disease received corticosteroids it was hard to assess their role in the outcome of TBE^[24]. In the group of Polish patients with TBE dexamethasone was used in 54.8% of patients with meningitis, in 69.6% of patients with meningoencephalitis and in 78.3% of patients with meningoencephalomyelitis. In this study the duration of hospitalization was significantly longer only in patients who received dexamethasone longer than 10 d^[49]. Perhaps corticosteroid treatment is effective in certain cases, but until the results of the randomized studies are available they cannot be recommend as a standard treatment approach^[24,49].

A case report on a patient with severe TBE who substantially improved after application of high dose intravenous immunoglobulins late in the disease course^[81], and reports on the successful treatment of encephalitis due to other arboviruses with high doses of intravenous immunoglobulins, prompted Růžek *et al.*^[82] to propose the implementation of randomized controlled treatment study on the efficacy of high dose intravenous immunoglobulins in patients with severe TBE. Such a proposal should not be interpreted as an indication for treatment of TBE with immunoglobulins.

NONSPECIFIC PREVENTIVE MEASURES

Nonspecific preventive measures comprise pasteurization of milk, reduction of tick population, and personal protective procedures.

Since milk from endemic regions may contain TBEV, pasteurization, and avoiding consumption of unpasteurized milk and dairy products, prevent infection of humans^[3,18].

Population of ticks can be reduced by impacts on the environment (*e.g.*, by regular cutting grass around the house, by usage of acaricides, and/or by control of deer population).

Nonspecific personal preventive measures include avoidance of ticks (which means avoidance of contact with vegetation, especially in deciduous and mixed forests with a rich understory and a layer of decaying vegetation on the ground that provides sufficient humidity for the development and survival of ticks), wearing light-colored clothing (light colors enable that ticks are better noticeable) with long sleeves and slacks stuck in socks or footwear (to diminish tick access to the skin), use of repellents, careful examination of the whole body for the presence of ticks, and removal of the attached ticks as soon as possible^[83]. However, TBEV is present in salivary glands of the infected tick and may be transmitted from the saliva of an infected tick within a few minutes after attachment^[3]. Although the recommended personal measures appear to be self-evident for the prevention of tick-borne diseases including TBE, the effectiveness of certain measures is limited, questionable or has not been properly assessed. Additional problem is that only a small proportion of exposed persons follow the recommended procedures in everyday life^[84,85].

VACCINATION

Historically, immunoglobulins containing gamma globulin against TBEV were used as a prophylaxis against TBE within 96 h after a tick bite in the TBE endemic regions (post-exposure prophylaxis). However, due to reports indicating a more severe disease course in children who had received the immunoglobulin^[86-88] and because protection was rather unreliable^[88], the usage of the immunoglobulins (passive immunization) in European Union has been abandoned^[89]. Nevertheless, according to recent studies in Russia, timely single administration of a specific immunoglobulin preparation in a dose of 0.05 mL/kg body weight was protective in approximately 80% of the cases^[90]. Further analysis of these findings is needed.

Active immunization is the most effective way to prevent TBE^[3,4]. Given that TBE is a zoonosis, that the source of infection is an infected animal, and that TBEV is transmitted by a tick bite and does not spread from human to human, vaccination enables only individual protection. Consequently, high immunization rate of a population in a given environment does not protect persons who are not vaccinated.

Recommendations for TBE vaccination

Because TBE incidence varies within and between different endemic areas recommendations for public vaccination strategies should be based on risk assessments conducted for individual region.

World Health Organization (WHO) recommends vaccination to people of all age groups, including children, in highly endemic areas (≥ 5 cases/100000 per year). In regions where the pre-vaccination incidence of TBE is low to moderate (5 year incidence < 5 cases/100000 per year) immunization should target

persons in the most vulnerable groups. The WHO also recommends TBE vaccination for people travelling from nonendemic areas to rural endemic areas up to altitudes of 1400 m^[91].

Central European Vaccination Awareness Group strongly recommends the introduction of universal TBE vaccination for persons > 1 year old for all countries at very high risk of TBE infections. For countries with a very low risk of TBE, recommendations for TBE vaccination should only apply to persons travelling to endemic regions^[92].

Vaccines

Two vaccines against TBE, FSME-IMMUN[®] and Encepur[®], are registered in Europe. They contain inactivated European subtype of TBEV-FSME-IMMUN is based on strain Neudorf 1, Encepur on strain K23. Both vaccines prevent not only the disease caused by the European but also those caused by the Siberian and Far-Eastern subtype of TBEV. Procedures used in the preparation of the two vaccines are similar (viruses are grown in chick embryo fibroblast cells, are inactivated by formaldehyde and are purified), and in both vaccines the adjuvant is aluminum hydroxide^[89]. Both vaccines are registered for adults and children aged 1 year and older (vaccines for children are called FSME-IMMUN 0.25 mL Junior, and Encepur Kinder, respectively).

In addition to European vaccines, two vaccines based on Far-Eastern subtype of TBEV are registered in Russia (TBE-Moscow and EnceVir); also in these vaccines viruses are grown in chick embryo cells and are inactivated with formalin. Another vaccine, which is produced and used in China, is also based on the Far-Eastern subtype of TBEV^[89].

Vaccination schedule

Several vaccination schedules exist; all of them consist of primary (basic) vaccination followed by booster doses. For a complete primary (basic) vaccination three doses, usually with an interval of 1-3 and 5-12 mo between first and second, and second and third dose, respectively, is required. Immunity is maintained with booster doses: the first booster dose is administered 3 years after completion of the primary vaccination, later on one dose is needed every 5 years. Due to the deterioration of the immune response in persons aged > 60 years (FSME IMMUN) or > 50 years (Encepur) boosters are recommended at 3 years intervals in this age group^[89].

Vaccination can begin at any time, but immunization with the first two doses is preferably carried out in the winter months to achieve protection before tick activity. Reports on individual cases of severe forms of TBE in subjects who had received only one dose of vaccine against TBE, had historically been an additional reason to start vaccination (receipt the first two doses) in winter. Subsequent information showed that incomplete vaccination does not pose an increased risk for severe

disease as compared to unvaccinated persons of the comparable age.

When it is desired to achieve protection in a short time, "fast schedule" can be used in accordance with the manufacturers' instructions^[3,4,85]. In contrast to classic approach, in "rapid vaccination" the second dose is usually administered 14 d instead of 1 to 3 mo after the first dose. Protection efficacy is comparable to that seen after classical schedule; however, scientific data on "quick schedule" is less comprehensive than for the conventional approach.

In a person who had not received the recommended doses according to the schedule but with longer intervals, vaccination does not need to be started again from the very beginning but continue with missing doses^[93]. Example: if someone had received only two doses of TBE vaccine 5 years ago and afterwards forgot to get the third dose, the vaccination should proceed with the third dose of basic vaccination and then with booster doses according to schedule. Or: a person who had received complete basic vaccination but for 15 years had not obtained any booster dose, does not need to repeat basic vaccination but just get the first booster dose and then continue immunization with booster doses according to the recommended schedule. Longer intervals between doses generally do not reduce antibody concentrations after completion of TBE vaccination, but protection in the period before the delayed dose is less reliable^[93].

Persons who had acquired TBE are esteemed to be protected against the disease and do not need vaccination.

Mode of application, dosages

TBE vaccine is administered intramuscularly into the deltoid muscle; in young children it can be given in the anterolateral thigh. The vaccine may be administered simultaneously with other vaccines (live or inactivated) but not on the same place^[89].

Doses depend upon the age of the recipient; the age limits for vaccines available in Europe slightly differ. In subjects younger than 16 years, the dose of the FSME-IMMUN vaccine is 0.25 mL, while for persons who have 16 years or are older 0.5 mL is recommended; in subjects younger than 12 years, the dose of the Encepur vaccine is 0.25 mL, while in older the dosage is 0.5 mL.

Efficacy and safety

Both European vaccines are safe and effective; particularly voluminous data exist for FSME-IMMUN vaccine of which more than 100 million doses were used.

Fourteen days after the second dose of basic vaccination protective antibodies develop in about 85% of the subjects, while after three doses more than 98% of persons with normal immunity are protected^[89].

In Austria, where the vast majority of the population is vaccinated, the incidence of TBE declined dramatically. The estimated protection after vaccination (field

effectiveness) is more than 98% for persons vaccinated according to the recommended program and more than 90% for those who received basic vaccination, but were later not vaccinated according to the planned schedule^[89].

Side effects are mild and relatively rare. They are more frequent after the first dose of vaccine than with later doses. Most commonly local pain and tenderness on pressure at the injection site take place; redness and swelling occur less often. Short-term fever after vaccination is rare in adults but relatively common in young children. Neurological complications are very rare^[89].

Storage

The vaccine must be stored in a refrigerator at a temperature between 2 °C and 8 °C. Storage at higher temperatures and freezing are not suitable^[89].

Limitations and contraindications

Vaccination is not carried out in subjects with acute febrile illness.

Vaccination is contraindicated in the case of: (1) A severe allergic reaction after previous dose of TBE vaccine; (2) Information on severe allergic reactions to vaccine components (in addition to the active ingredients, TBE vaccine also contains remnants of formaldehyde, neomycin, gentamicin and protamine sulfate); and (3) Anaphylactic hypersensitivity to eggs (TBE viruses are grown in chick embryo fibroblast cells).

Controlled clinical trials to assess the safety of TBE vaccine during pregnancy are not available; thus, pregnant women are vaccinated only after a careful individual assessment of the potential risks and benefits. There is also no sufficient data on the safety of vaccination during lactation. It is not known whether the vaccine components are excreted in human milk or not, however, mother's antibodies produced after vaccination against TBE are probably present in the milk and consequently breast-feeding baby may come in contact with them. Given that TBE vaccines are based on inactivated virus, the damage of fetus or breast-feeding child is highly unlikely.

Although there is no evidence that vaccination may trigger autoimmunity or worsen the course of autoimmune diseases, caution is needed in subjects with an autoimmune disease since information to ensure the safety of vaccination in this group of patients is limited^[89].

The manufacturer of the vaccine and some recommendations suggest that in immunocompromised persons the efficacy of vaccination is verified by serological testing approximately four weeks after the second dose, and that in case of an inadequate antibody response the second dose is repeated and followed by the third dose in accordance with the normal vaccination schedule. According to some suggestions

similar approach may apply also to all subsequent doses. Although the approach appears logical, there is no convincing clinical data to substantiate its use. As a rule the effectiveness of protection after vaccination against TBE is not verified by the detection of antibodies against TBEV in serum.

CONCLUSION

TBE is an important tick-borne central nervous system infection in Europe and Asia. Due to relatively severe clinical course combined with the absence of etiologic treatment, considerable proportion of patients with incomplete recovery after acute illness and increasing incidence, it represents a growing (public) health problem that could be substantially reduced with vaccination.

REFERENCES

- 1 Süß J. Tick-borne encephalitis 2010: epidemiology, risk areas, and virus strains in Europe and Asia-an overview. *Ticks Tick Borne Dis* 2011; **2**: 2-15 [PMID: 21771531 DOI: 10.1016/j.ttbdis.2010.10.007]
- 2 Mansfield KL, Johnson N, Phipps LP, Stephenson JR, Fooks AR, Solomon T. Tick-borne encephalitis virus-a review of an emerging zoonosis. *J Gen Virol* 2009; **90**: 1781-1794 [PMID: 19420159 DOI: 10.1099/vir.0.011437-0]
- 3 Lindquist L, Vapalahti O. Tick-borne encephalitis. *Lancet* 2008; **371**: 1861-1871 [PMID: 18514730 DOI: 10.1016/S0140-6736(08)60800-4]
- 4 Kaiser R. Tick-borne encephalitis. *Infect Dis Clin North Am* 2008; **22**: 561-575, x [PMID: 18755391 DOI: 10.1016/j.idc.2008.03.013]
- 5 Lotric-Furlan S, Avsic-Zupanc T, Strle F. An abortive form of tick-borne encephalitis (TBE)--a rare clinical manifestation of infection with TBE virus. *Wien Klin Wochenschr* 2002; **114**: 627-629 [PMID: 12422615]
- 6 Zilber LA. Spring-summer tick-borne encephalitis. *Arkhiv Biol Nauk* 1939; **56**: 255-261
- 7 Ecker M, Allison SL, Meixner T, Heinz FX. Sequence analysis and genetic classification of tick-borne encephalitis viruses from Europe and Asia. *J Gen Virol* 1999; **80** (Pt 1): 179-185 [PMID: 9934700]
- 8 Süß J. Tick-borne encephalitis in Europe and beyond--the epidemiological situation as of 2007. *Euro Surveill* 2008; **13**: 18916 [PMID: 18761916]
- 9 Randolph SE. Human activities predominate in determining changing incidence of tick-borne encephalitis in Europe. *Euro Surveill* 2010; **15**: 24-31 [PMID: 20630144]
- 10 Dobler G, Gniel D, Petermann R, Pfeffer M. Epidemiology and distribution of tick-borne encephalitis. *Wien Med Wochenschr* 2012; **162**: 230-238 [PMID: 22699708 DOI: 10.1007/s10354-012-0100-5]
- 11 Kovalev SY, Chernykh DN, Kokorev VS, Snitkovskaya TE, Romanenko VV. Origin and distribution of tick-borne encephalitis virus strains of the Siberian subtype in the Middle Urals, the north-west of Russia and the Baltic countries. *J Gen Virol* 2009; **90**: 2884-2892 [PMID: 19656959 DOI: 10.1099/vir.0.012419-0]
- 12 Dobler G. Zoonotic tick-borne flaviviruses. *Vet Microbiol* 2010; **140**: 221-228 [PMID: 19765917 DOI: 10.1016/j.vetmic.2009.08.024]
- 13 Donoso Mantke O, Escadafal C, Niedrig M, Pfeffer M, Working Group For Tick-Borne Encephalitis Virus C. Tick-borne encephalitis in Europe, 2007 to 2009. *Euro Surveill* 2011; **16**: 19976 [PMID: 21968423]
- 14 Institute of Public Health of the Republic of Slovenia. Epidemiological surveillance of infectious diseases in Slovenia in year 2013. *Tick-borne encephalitis* 2014: 57-59
- 15 Süß J. Epidemiology and ecology of TBE relevant to the production of effective vaccines. *Vaccine* 2003; **21** Suppl 1: S19-S35 [PMID: 12628811 DOI: 10.1016/S0264-410X(02)00812-5]
- 16 Durmiš E, Knap N, Saksida A, Trilar T, Duh D, Avšič-Županc T.

- Prevalence and molecular characterization of tick-borne encephalitis virus in Ixodes ricinus ticks collected in Slovenia. *Vector Borne Zoonotic Dis* 2011; **11**: 659-664 [PMID: 21028962 DOI: 10.1089/vbz.2010.0054]
- 17 **Kríz B**, Benes C, Daniel M. Alimentary transmission of tick-borne encephalitis in the Czech Republic (1997-2008). *Epidemiol Mikrobiol Immunol* 2009; **58**: 98-103 [PMID: 19526924]
 - 18 **Hudopisk N**, Korva M, Janet E, Simetinger M, Grgič-Vitek M, Gubenšek J, Natek V, Kraigher A, Strle F, Avšič-Županc T. Tick-borne encephalitis associated with consumption of raw goat milk, Slovenia, 2012. *Emerg Infect Dis* 2013; **19**: 806-808 [PMID: 23697658 DOI: 10.3201/eid1905.121442]
 - 19 **Avsic-Zupanc T**, Poljak M, Maticic M, Radsel-Medvescek A, LeDuc JW, Stiasny K, Kunz C, Heinz FX. Laboratory acquired tick-borne meningoencephalitis: characterisation of virus strains. *Clin Diagn Virol* 1995; **4**: 51-59 [PMID: 15566827 DOI: 10.1016/0928-0197(94)00062-Y]
 - 20 **Dumpis U**, Crook D, Oksi J. Tick-borne encephalitis. *Clin Infect Dis* 1999; **28**: 882-890 [PMID: 10825054 DOI: 10.1086/515195]
 - 21 **Gritsun TS**, Lashkevich VA, Gould EA. Tick-borne encephalitis. *Antiviral Res* 2003; **57**: 129-146 [PMID: 12615309 DOI: 10.1016/S0166-3542(02)00206-1]
 - 22 **Logar M**, Bogovic P, Cerar D, Avsic-Zupanc T, Strle F. Tick-borne encephalitis in Slovenia from 2000 to 2004: comparison of the course in adult and elderly patients. *Wien Klin Wochenschr* 2006; **118**: 702-707 [PMID: 17160611 DOI: 10.1007/s00508-006-0699-6]
 - 23 **Lesnicar G**, Poljak M, Seme K, Lesnicar J. Pediatric tick-borne encephalitis in 371 cases from an endemic region in Slovenia, 1959 to 2000. *Pediatr Infect Dis J* 2003; **22**: 612-617 [PMID: 12867836 DOI: 10.1097/00006454-200307000-00009]
 - 24 **Mickiene A**, Laiskonis A, Günther G, Vene S, Lundkvist A, Lindquist L. Tickborne encephalitis in an area of high endemicity in Lithuania: disease severity and long-term prognosis. *Clin Infect Dis* 2002; **35**: 650-658 [PMID: 12203160]
 - 25 **Arnez M**, Avsic-Zupanc T. Tick-borne encephalitis in children: an update on epidemiology and diagnosis. *Expert Rev Anti Infect Ther* 2009; **7**: 1251-1260 [PMID: 19968516 DOI: 10.1586/eri.09.99]
 - 26 **Hansson ME**, Orvell C, Engman ML, Wide K, Lindquist L, Lidelfelt KJ, Sundin M. Tick-borne encephalitis in childhood: rare or missed? *Pediatr Infect Dis J* 2011; **30**: 355-357 [PMID: 21412206 DOI: 10.1097/INF.0b013e3181fe3b5a]
 - 27 **Bröker M**, Gniel D. New foci of tick-borne encephalitis virus in Europe: consequences for travellers from abroad. *Travel Med Infect Dis* 2003; **1**: 181-184 [PMID: 17291912 DOI: 10.1016/j.tmaid.2003.07.001]
 - 28 **Rendi-Wagner P**. Risk and prevention of tick-borne encephalitis in travelers. *J Travel Med* 2004; **11**: 307-312 [PMID: 15544715 DOI: 10.2310/7060.2004.19107]
 - 29 **Růžek D**, Dobler G, Donoso Mantke O. Tick-borne encephalitis: pathogenesis and clinical implications. *Travel Med Infect Dis* 2010; **8**: 223-232 [PMID: 20970725 DOI: 10.1016/j.tmaid.2010.06.004]
 - 30 **Mandl CW**. Steps of the tick-borne encephalitis virus replication cycle that affect neuropathogenesis. *Virus Res* 2005; **111**: 161-174 [PMID: 15871909 DOI: 10.1016/j.virusres.2005.04.007]
 - 31 **McMinn PC**. The molecular basis of virulence of the encephalitic flaviviruses. *J Gen Virol* 1997; **78** (Pt 11): 2711-2722 [PMID: 9367356]
 - 32 **Gelpi E**, Preusser M, Laggner U, Garzuly F, Holzmann H, Heinz FX, Budka H. Inflammatory response in human tick-borne encephalitis: analysis of postmortem brain tissue. *J Neurovirol* 2006; **12**: 322-327 [PMID: 16966222 DOI: 10.1080/13550280600848746]
 - 33 **Gustafson R**, Svenungsson B, Forsgren M, Gardulf A, Granström M. Two-year survey of the incidence of Lyme borreliosis and tick-borne encephalitis in a high-risk population in Sweden. *Eur J Clin Microbiol Infect Dis* 1992; **11**: 894-900 [PMID: 1486884 DOI: 10.1007/BF01962369]
 - 34 **Bogovic P**, Lotric-Furlan S, Strle F. What tick-borne encephalitis may look like: clinical signs and symptoms. *Travel Med Infect Dis* 2010; **8**: 246-250 [PMID: 20970727 DOI: 10.1016/j.tmaid.2010.05.011]
 - 35 **Beer S**, Brune N, Kesselring J. Detection of anterior horn lesions by MRI in central European tick-borne encephalomyelitis. *J Neurol* 1999; **246**: 1169-1171 [PMID: 10653310 DOI: 10.1007/s004150050537]
 - 36 **Fausser S**, Stich O, Rauer S. Unusual case of tick borne encephalitis with isolated myeloradiculitis. *J Neurol Neurosurg Psychiatry* 2007; **78**: 909-910 [PMID: 17237139 DOI: 10.1136/jnnp.2006.110882]
 - 37 **Haglund M**, Forsgren M, Lindh G, Lindquist L. A 10-year follow-up study of tick-borne encephalitis in the Stockholm area and a review of the literature: need for a vaccination strategy. *Scand J Infect Dis* 1996; **28**: 217-224 [PMID: 8863349 DOI: 10.3109/00365549609027160]
 - 38 **Kaiser R**. The clinical and epidemiological profile of tick-borne encephalitis in southern Germany 1994-98: a prospective study of 656 patients. *Brain* 1999; **122** (Pt 11): 2067-2078 [PMID: 10545392 DOI: 10.1093/brain/122.11.2067]
 - 39 **Lotric-Furlan S**, Strle F. Peripheral facial palsy in patients with tick-borne encephalitis. *Clin Microbiol Infect* 2012; **18**: 1027-1032 [PMID: 22192120 DOI: 10.1111/j.1469-0691.2011.03719.x]
 - 40 **Kleiter I**, Steinbrecher A, Flügel D, Bogdahn U, Schulte-Mattler W. Autonomic involvement in tick-borne encephalitis (TBE): report of five cases. *Eur J Med Res* 2006; **11**: 261-265 [PMID: 16820340]
 - 41 **Misić-Majerus L**, Bujić N, Madarić V, Avsic-Zupanc T. [An abortive type of tick-borne meningoencephalitis]. *Acta Med Croatica* 2003; **57**: 111-116 [PMID: 12879690]
 - 42 **Meyer PM**, Zimmermann H, Goetschel P. Tick-borne encephalitis presenting as fever without localising signs--a case series. *Eur J Pediatr* 2010; **169**: 767-769 [PMID: 19888601 DOI: 10.1007/s00431-009-1097-7]
 - 43 **Ustinova Olu**, Volechova GM, Deviatkov MI, Gusmanova AI. [The clinico-epidemiological characteristics of tick-borne encephalitis in Perm Province]. *Zh Mikrobiol Epidemiol Immunobiol* 1997; **(3)**: 33-36 [PMID: 9304324]
 - 44 **Pöschl P**, Kleiter I, Grubwinkler S, Bumess E, Bogdahn U, Dobler G, Steinbrecher A. [Severe tick-borne encephalomyelitis with lack of cerebrospinal fluid pleocytosis]. *Fortschr Neurol Psychiatr* 2009; **77**: 591-593 [PMID: 19821222 DOI: 10.1055/s-0028-1109768]
 - 45 **Poponnikova TV**. The clinical picture of chronic tick-borne encephalitis in children. *Int J Med Microbiol* 2008; **298** Suppl 1: S351-355 [DOI: 10.1016/j.ijmm.2008.03.007]
 - 46 **Kaiser R**. Tick-borne encephalitis--still a serious disease? *Wien Med Wochenschr* 2012; **162**: 229 [PMID: 22695812 DOI: 10.1007/s10354-012-0119-7]
 - 47 **Haglund M**, Günther G. Tick-borne encephalitis--pathogenesis, clinical course and long-term follow-up. *Vaccine* 2003; **21** Suppl 1: S11-S18 [PMID: 12628810 DOI: 10.1016/S0264-410X(02)00811-3]
 - 48 **Misić Majerus L**, Daković Rode O, Ruzić Sabljic E. [Post-encephalitic syndrome in patients with tick-borne encephalitis]. *Acta Med Croatica* 2009; **63**: 269-278 [PMID: 20034327]
 - 49 **Czupryna P**, Moniuszko A, Pancewicz SA, Grygorczuk S, Kondrusik M, Zajkowska J. Tick-borne encephalitis in Poland in years 1993-2008--epidemiology and clinical presentation. A retrospective study of 687 patients. *Eur J Neurol* 2011; **18**: 673-679 [PMID: 21143706 DOI: 10.1111/j.1468-1331.2010.03278.x]
 - 50 **Logar M**, Arnez M, Kolbl J, Avsic-Zupanc T, Strle F. Comparison of the epidemiological and clinical features of tick-borne encephalitis in children and adults. *Infection* 2000; **28**: 74-77 [PMID: 10782391 DOI: 10.1007/s150100050050]
 - 51 **Wahlberg P**, Saikku P, Brummer-Korvenkontio M. Tick-borne viral encephalitis in Finland. The clinical features of Kumlunge disease during 1959-1987. *J Intern Med* 1989; **225**: 173-177 [PMID: 2703799 DOI: 10.1111/j.1365-2796.1989.tb00059.x]
 - 52 **McNair AN**, Brown JL. Tick-borne encephalitis complicated by monoplegia and sensorineural deafness. *J Infect* 1991; **22**: 81-86 [PMID: 2002236 DOI: 10.1016/0163-4453(91)91094-E]
 - 53 **Anić K**, Soldo I, Perić L, Karner I, Barac B. Tick-borne encephalitis in eastern Croatia. *Scand J Infect Dis* 1998; **30**: 509-512 [PMID: 10066055 DOI: 10.1080/00365549850161548]
 - 54 **Jereb M**, Muzlovic I, Avsic-Zupanc T, Karner P. Severe tick-borne encephalitis in Slovenia: epidemiological, clinical and laboratory

- findings. *Wien Klin Wochenschr* 2002; **114**: 623-626 [PMID: 12422614]
- 55 **Tomazic J**, Pikelj F, Schwartz B, Kunze M, Kraigher A, Matjašič M, Lesničar G, Gorišek J, Kunz C, Marth E; Slovenian TBE study group. The clinical features of tick-borne encephalitis in Slovenia. A study of 492 cases in 1994. *Antibiot Monitor* 1996; **12**: 115-120
- 56 **Kunz C**. TBE vaccination and the Austrian experience. *Vaccine* 2003; **21** Suppl 1: S50-S55 [PMID: 12628814 DOI: 10.1016/S0264-410X(02)00813-7]
- 57 **Kindberg E**, Mickiene A, Ax C, Akerlind B, Vene S, Lindquist L, Lundkvist A, Svensson L. A deletion in the chemokine receptor 5 (CCR5) gene is associated with tickborne encephalitis. *J Infect Dis* 2008; **197**: 266-269 [PMID: 18179389 DOI: 10.1086/524709]
- 58 **Barkhash AV**, Perelygin AA, Babenko VN, Brinton MA, Voevoda MI. Single nucleotide polymorphism in the promoter region of the CD209 gene is associated with human predisposition to severe forms of tick-borne encephalitis. *Antiviral Res* 2012; **93**: 64-68 [PMID: 22061615 DOI: 10.1016/j.antiviral.2011.10.017]
- 59 **Barkhash AV**, Perelygin AA, Babenko VN, Myasnikova NG, Pilipenko PI, Romaschenko AG, Voevoda MI, Brinton MA. Variability in the 2'-5'-oligoadenylate synthetase gene cluster is associated with human predisposition to tick-borne encephalitis virus-induced disease. *J Infect Dis* 2010; **202**: 1813-1818 [PMID: 21050126 DOI: 10.1086/657418]
- 60 **Lotric-Furlan S**, Petrovec M, Avsic-Zupanc T, Nicholson WL, Sumner JW, Childs JE, Strle F. Prospective assessment of the etiology of acute febrile illness after a tick bite in Slovenia. *Clin Infect Dis* 2001; **33**: 503-510 [PMID: 11462187]
- 61 **Lotric-Furlan S**, Strle F. Thrombocytopenia--a common finding in the initial phase of tick-borne encephalitis. *Infection* 1995; **23**: 203-206 [PMID: 8522376 DOI: 10.1007/BF01781197]
- 62 **Kaiser R**, Holzmann H. Laboratory findings in tick-borne encephalitis--correlation with clinical outcome. *Infection* 2000; **28**: 78-84 [PMID: 10782392 DOI: 10.1007/s150100050051]
- 63 **Holzmann H**. Diagnosis of tick-borne encephalitis. *Vaccine* 2003; **21** Suppl 1: S36-S40 [PMID: 12628812 DOI: 10.1016/S0264-410X(02)00819-8]
- 64 **Saksida A**, Duh D, Lotric-Furlan S, Strle F, Petrovec M, Avsic-Zupanc T. The importance of tick-borne encephalitis virus RNA detection for early differential diagnosis of tick-borne encephalitis. *J Clin Virol* 2005; **33**: 331-335 [PMID: 15919235 DOI: 10.1016/j.jcv.2004.07.014]
- 65 **Ludolf D**, Reinholz M, Schmitz H. Highly specific detection of antibodies to tick-borne encephalitis (TBE) virus in humans using a domain III antigen and a sensitive immune complex (IC) ELISA. *J Clin Virol* 2009; **45**: 125-128 [PMID: 19409841 DOI: 10.1016/j.jcv.2009.03.016]
- 66 **Hofmann H**, Kunz C, Heinz FX, Dippe H. Detectability of IgM antibodies against TBE virus after natural infection and after vaccination. *Infection* 1983; **11**: 164-166 [PMID: 6309676 DOI: 10.1007/BF01641297]
- 67 **Günther G**, Haglund M, Lindquist L, Sköldenberg B, Forsgren M. Intrathecal IgM, IgA and IgG antibody response in tick-borne encephalitis. Long-term follow-up related to clinical course and outcome. *Clin Diagn Virol* 1997; **8**: 17-29 [PMID: 9248655 DOI: 10.1016/S0928-0197(97)00273-0]
- 68 **Heinz FX**, Stiasny K. Flaviviruses and their antigenic structure. *J Clin Virol* 2012; **55**: 289-295 [PMID: 22999801 DOI: 10.1016/j.jcv.2012.08.024]
- 69 **Stiasny K**, Aberle JH, Chmelik V, Karrer U, Holzmann H, Heinz FX. Quantitative determination of IgM antibodies reduces the pitfalls in the serodiagnosis of tick-borne encephalitis. *J Clin Virol* 2012; **54**: 115-120 [PMID: 22421535 DOI: 10.1016/j.jcv.2012.02.016]
- 70 **Stiasny K**, Holzmann H, Heinz FX. Characteristics of antibody responses in tick-borne encephalitis vaccination breakthroughs. *Vaccine* 2009; **27**: 7021-7026 [PMID: 19789092 DOI: 10.1016/j.vaccine.2009.09.069]
- 71 **Lotric-Furlan S**, Avsic-Zupanc T, Strle F. Tick-borne encephalitis after active immunization. *Int J Med Microbiol* 2008; **298**: 309-313 [DOI: 10.1016/j.ijmm.2008.03.006]
- 72 **Andersson CR**, Vene S, Insulander M, Lindquist L, Lundkvist A, Günther G. Vaccine failures after active immunisation against tick-borne encephalitis. *Vaccine* 2010; **28**: 2827-2831 [PMID: 20167301 DOI: 10.1016/j.vaccine.2010.02.001]
- 73 **Kristoferitsch W**, Stanek G, Kunz C. [Double infection with early summer meningoencephalitis virus and *Borrelia burgdorferi*]. *Dtsch Med Wochenschr* 1986; **111**: 861-864 [PMID: 3011378 DOI: 10.1055/s-2008-1068546]
- 74 **Oksi J**, Viljanen MK, Kalimo H, Peltonen R, Marttinen R, Salomaa P, Nikoskelainen J, Budka H, Halonen P. Fatal encephalitis caused by concomitant infection with tick-borne encephalitis virus and *Borrelia burgdorferi*. *Clin Infect Dis* 1993; **16**: 392-396 [PMID: 8452951 DOI: 10.1093/clind/16.3.392]
- 75 **Cimperman J**, Maraspin V, Lotric-Furlan S, Ruzič-Sabljčić E, Avsic-Zupanc T, Picken RN, Strle F. Concomitant infection with tick-borne encephalitis virus and *Borrelia burgdorferi* sensu lato in patients with acute meningitis or meningoencephalitis. *Infection* 1998; **26**: 160-164 [PMID: 9646107 DOI: 10.1007/BF02771842]
- 76 **Lotric-Furlan S**, Petrovec M, Avsic-Zupanc T, Strle F. Concomitant tickborne encephalitis and human granulocytic ehrlichiosis. *Emerg Infect Dis* 2005; **11**: 485-488 [PMID: 15757574 DOI: 10.3201/eid1103.040776]
- 77 **Lotric-Furlan S**, Petrovec M, Avsic-Zupanc T, Logar M, Strle F. Epidemiological, clinical and laboratory distinction between human granulocytic ehrlichiosis and the initial phase of tick-borne encephalitis. *Wien Klin Wochenschr* 2002; **114**: 636-640 [PMID: 12422617]
- 78 **Kumar G**, Kalita J, Misra UK. Raised intracranial pressure in acute viral encephalitis. *Clin Neurol Neurosurg* 2009; **111**: 399-406 [PMID: 19372001 DOI: 10.1016/j.clineuro.2009.03.004]
- 79 **Diringer MN**, Zazulia AR. Osmotic therapy: fact and fiction. *Neurocrit Care* 2004; **1**: 219-233 [PMID: 16174920 DOI: 10.1385/NCC: 1: 2: 219]
- 80 **Sommer JB**, Heckmann JG, Kraus J, Neundörfer B, Erbguth FJ. [Generalized brain edema in non-purulent meningoencephalitis. The anti-edema effect of therapy with dexamethasone]. *Nervenarzt* 2000; **71**: 112-115 [PMID: 10703012 DOI: 10.1007/s001150050016]
- 81 **Kleiter I**, Jilg W, Bogdahn U, Steinbrecher A. Delayed humoral immunity in a patient with severe tick-borne encephalitis after complete active vaccination. *Infection* 2007; **35**: 26-29 [PMID: 17297586 DOI: 10.1007/s15010-006-6614-2]
- 82 **Růžek D**, Dobler G, Niller HH. May early intervention with high dose intravenous immunoglobulin pose a potentially successful treatment for severe cases of tick-borne encephalitis? *BMC Infect Dis* 2013; **13**: 306 [PMID: 23822550 DOI: 10.1186/1471-2334-13-306]
- 83 **Stanek G**, Wormser GP, Gray J, Strle F. Lyme borreliosis. *Lancet* 2012; **379**: 461-473 [PMID: 21903253 DOI: 10.1016/S0140-6736(11)60103-7]
- 84 **Vázquez M**, Muehlenbein C, Cartter M, Hayes EB, Ertel S, Shapiro ED. Effectiveness of personal protective measures to prevent Lyme disease. *Emerg Infect Dis* 2008; **14**: 210-216 [PMID: 18258112 DOI: 10.3201/eid1402.070725]
- 85 **Corapi KM**, White MI, Phillips CB, Daltroy LH, Shadick NA, Liang MH. Strategies for primary and secondary prevention of Lyme disease. *Nat Clin Pract Rheumatol* 2007; **3**: 20-25 [PMID: 17203005 DOI: 10.1038/npcrhum0374]
- 86 **Bröker M**, Kollaritsch H. After a tick bite in a tick-borne encephalitis virus endemic area: current positions about post-exposure treatment. *Vaccine* 2008; **26**: 863-868 [PMID: 18206274 DOI: 10.1016/j.vaccine.2007.11.046]
- 87 **Waldvogel K**, Bossart W, Huisman T, Boltshauser E, Nadal D. Severe tick-borne encephalitis following passive immunization. *Eur J Pediatr* 1996; **155**: 775-779 [PMID: 8874110 DOI: 10.1007/BF02002905]
- 88 **Kluger G**, Schöttler A, Waldvogel K, Nadal D, Hinrichs W, Wündisch GF, Laub MC. Tickborne encephalitis despite specific immunoglobulin prophylaxis. *Lancet* 1995; **346**: 1502 [PMID: 7491037 DOI: 10.1016/S0140-6736(95)92527-9]
- 89 **Barrett PN**, Portsmouth D, Ehrlich HJ. Tick-borne encephalitis

- virus vaccines. In: Vaccines. Plotkin SA, Orenstein W, Offit PA, editors. 6th ed. PA, USA: Elsevier, 2013: 773-788
- 90 **Pen'evskaia NA**, Rudakov NV. [Efficiency of use of immunoglobulin preparations for the postexposure prevention of tick-borne encephalitis in Russia (a review of semi-centennial experience)]. *Med Parazitol (Mosk)* 2010; **(1)**: 53-59 [PMID: 20361640]
- 91 Vaccines against tick-borne encephalitis: WHO position paper. *Wkly Epidemiol Rec* 2011; **86**: 241-256 [PMID: 21661276]
- 92 **Zavadska D**, Anca I, André F, Bakir M, Chlibek R, Cižman M, Ivaskeviciene I, Mangarov A, Mézner Z, Pokorn M, Prymula R, Richter D, Salman N, Simurka P, Tamm E, Tešović G, Urbancikova I, Usonis V. Recommendations for tick-borne encephalitis vaccination from the Central European Vaccination Awareness Group (CEVAG). *Hum Vaccin Immunother* 2013; **9**: 362-374 [PMID: 23291941 DOI: 10.4161/hv.22766]
- 93 **Schöndorf I**, Schönfeld C, Nicolay U, Zent O, Banzhoff A. Response to tick-borne encephalitis (TBE) booster vaccination after prolonged time intervals to primary immunization with the rapid schedule. *Int J Med Microbiol* 2006; **296** Suppl 40: 208-212 [PMID: 16531118 DOI: 10.1016/j.ijmm.2006.01.009]
- 94 **Heinz FX**, Stiasny K, Holzmann H, Grgic-Vitek M, Kriz B, Essl A, Kundi M. Vaccination and tick-borne encephalitis, central Europe. *Emerg Infect Dis* 2013; **19**: 69-76 [PMID: 23259984 DOI: 10.3201/eid1901.120458]

P- Reviewer: Eugenin EA, Tetsuya T, Xavier-Elsas P
S- Editor: Tian YL **L- Editor:** A **E- Editor:** Lu YJ





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

