

World Journal of *Gastroenterology*

World J Gastroenterol 2024 July 21; 30(27): 3264-3360



EDITORIAL

- 3264 Effects of excess high-normal alanine aminotransferase levels in relation to new-onset metabolic dysfunction-associated fatty liver disease: Clinical implications
McGinty G, Przemioslo R
- 3268 What aspects do we overlook in the rehabilitation of patients with inflammatory bowel disease?
Ata BN, Eyigor S
- 3273 Novel insights into autophagy in gastrointestinal pathologies, mechanisms in metabolic dysfunction-associated fatty liver disease and acute liver failure
Velikova T, Gulinac M
- 3278 Gastric cystica profunda: Another indication for minimally invasive endoscopic resection techniques?
Bedi HK, Motomura D, Shahidi N
- 3284 Pro and anti-inflammatory diets as strong epigenetic factors in inflammatory bowel disease
Rostami A, White K, Rostami K

ORIGINAL ARTICLE

Case Control Study

- 3290 Targeted metabolomics study of fatty-acid metabolism in lean metabolic-associated fatty liver disease patients
Sun PQ, Dong WM, Yuan YF, Cao Q, Chen XY, Guo LL, Jiang YY

Clinical Trials Study

- 3304 Vonoprazan-amoxicillin dual therapy for *Helicobacter pylori* eradication in Chinese population: A prospective, multicenter, randomized, two-stage study
Huang XP, Liu YJ, Lin SW, Shao YF, Qiu F, Qiu QW, Xu ZK, Chen JX, Chen LH, Lin ZQ, Dai WH, Zhang MQ, Jiang Q, Xiao ZQ, Cheng XX, Zhang XF, You WB, Chen W, Li LQ, Lin WX, Wang YF, Lai FJ, Chen LQ, Huang ZH, Zheng WQ, Wei JQ, Lin ZH

Observational Study

- 3314 Nomogram based on liver stiffness and spleen area with ultrasound for posthepatectomy liver failure: A multicenter study
Cheng GW, Fang Y, Xue LY, Zhang Y, Xie XY, Qiao XH, Li XQ, Guo J, Ding H

Randomized Controlled Trial

- 3326 Endoscopic polidocanol foam sclerobanding for the treatment of grade II-III internal hemorrhoids: A prospective, multi-center, randomized study
Qu CY, Zhang FY, Wang W, Gao FY, Lin WL, Zhang H, Chen GY, Zhang Y, Li MM, Li ZH, Cai MH, Xu LM, Shen F

Basic Study

- 3336** Distinct gut microbiomes in Thai patients with colorectal polyps

Intarajak T, Udomchaiprasertkul W, Khoiri AN, Sutheeworapong S, Kusonmano K, Kittichotirat W, Thammarongtham C, Cheevadhanarak S

LETTER TO THE EDITOR

- 3356** More on the interplay between gut microbiota, autophagy, and inflammatory bowel disease is needed

Subramanian A, Jahabardeen A, Thamaraikani T, Vellapandian C

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Shivananda Nayak, MSc, PhD, FAGE, FACB, NRCC-CC, FISBT, FABM, PGDCHC, DSc, Full Professor, Professor, Department of Preclinical Sciences, Faculty of Medical Sciences, The University of The West Indies, Mount Hope (0000), Trinidad and Tobago. shiv25@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE), MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for WJG as 4.3; Quartile: Q1. The WJG's CiteScore for 2023 is 7.8.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Xiao-Mei Zheng; Production Department Director: Xu Guo; Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski

EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF

Xian-Jun Yu (Pancreatic Oncology), Jian-Gao Fan (Chronic Liver Disease), Hou-Bao Liu

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

July 21, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

PUBLISHING PARTNER

Shanghai Pancreatic Cancer Institute and Pancreatic Cancer Institute, Fudan University
Biliary Tract Disease Institute, Fudan University

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/bpg/GerInfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

PUBLISHING PARTNER's OFFICIAL WEBSITE

<https://www.shca.org.cn>
<https://www.zs-hospital.sh.cn>



What aspects do we overlook in the rehabilitation of patients with inflammatory bowel disease?

Benil Nesli Ata, Sibel Eyigor

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade A, Grade C, Grade C

Novelty: Grade A, Grade B, Grade B

Creativity or Innovation: Grade A, Grade B, Grade C

Scientific Significance: Grade A, Grade B, Grade B

P-Reviewer: Liu W; Wang YD; Wu L

Received: March 15, 2024

Revised: June 5, 2024

Accepted: June 28, 2024

Published online: July 21, 2024

Processing time: 117 Days and 14.6 Hours



Benil Nesli Ata, Department of Physical Medicine and Rehabilitation, Izmir City Hospital, Izmir 35530, Türkiye

Sibel Eyigor, Department of Physical Medicine and Rehabilitation, Faculty of Medicine, Ege University, Izmir 35100, Türkiye

Corresponding author: Benil Nesli Ata, MD, Doctor, Department of Physical Medicine and Rehabilitation, Izmir City Hospital, Sevkett Ince Mah. 2148/11 Sk No. 1/11 Bayrakli/Izmir 35530, Türkiye. drbenilnesli@gmail.com

Abstract

In this editorial, we comment on the article by Stafie *et al.* Inflammatory bowel disease (IBD) constitutes a cluster of chronic and progressive inflammatory disorders affecting the digestive system. IBD can impede an individual's capacity to perform daily activities, hinder work productivity, limit physical capabilities, and negatively impact medical outcomes. Although physical activity and structured exercise programs are becoming increasingly important in many chronic inflammatory diseases, they are not being sufficiently implemented in IBD patients. Effective prevention of future disability and drug dependence in IBD patients requires timely diagnosis and treatment of musculoskeletal problems, including sarcopenia, as well as decreased muscle strength, aerobic capacity, and bone mineral density. To improve treatment outcomes for IBD patients, it is crucial to develop individualized rehabilitation programs tailored to their unique needs. Equally critical is the active participation of pertinent departments in this process. It is imperative to highlight the significance of creating a personalized rehabilitation program with a multidisciplinary approach in IBD management.

Key Words: Inflammatory bowel disease; Physical activity; Disability; Sarcopenia; Structured exercise; Rehabilitation

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Effectively managing clinical issues in inflammatory bowel disease (IBD) patients is critical to reducing the risk of long-term disability and facilitating optimal health outcomes. This includes addressing concerns such as decreased bone mineral density, muscle weakness, limited aerobic capacity, and sarcopenia. Customized structured exercise programs should be provided for patients in accordance with their individual needs, considering factors such as joint involvement, frailty, fatigue, and disease activity. Healthcare providers should adopt a multidisciplinary approach to provide personalized exercise recommendations, educate patients, and address misconceptions. This approach improves quality of life, minimizes complications associated with IBD, and enhances treatment success.

Citation: Ata BN, Eyigor S. What aspects do we overlook in the rehabilitation of patients with inflammatory bowel disease? *World J Gastroenterol* 2024; 30(27): 3268-3272

URL: <https://www.wjgnet.com/1007-9327/full/v30/i27/3268.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v30.i27.3268>

INTRODUCTION

Inflammatory bowel disease (IBD) is an umbrella term that encompasses two disease entities, Crohn's disease and ulcerative colitis, which are chronic, progressive inflammatory disorders of the digestive system. This disease imposes a significant burden on healthcare systems, leading to substantial annual healthcare expenditures. IBD is a condition that can lead to a variety of primary and secondary symptoms that can include diarrhea, abdominal pain, urgency, fatigue, frailty, anorexia, and depression[1]. Secondary symptoms significantly impact an individual's daily routine, work productivity, academic performance, physical abilities, medical outcomes, and overall quality of life (QOL). Young patients with IBD are seldom advised to engage in physical activity, which can result in a sedentary lifestyle due to concerns regarding potential complications[2].

OVERLOOKED ASPECTS IN INFLAMMATORY BOWEL DISEASE REHABILITATION

It is crucial to distinguish between the terms "structured exercise" and "physical activity" because they are frequently used interchangeably. Structured exercise and physical activity are two distinct types of activities that exhibit physiological differences. Structured exercise is a specialized and comprehensive program aimed at enhancing the functions of specific body systems, such as cardiovascular endurance and muscular strength. However, physical activity encompasses any form of movement that results in an increase in energy expenditure. Numerous chronic illnesses are believed to be exacerbated by inadequate physical activity[3,4]. The "BE-FIT-IBD" study found that 42.9% of IBD patients were physically inactive, despite their awareness of the importance of physical activity in managing the disease. Fear of disease flare-ups, patients' inadequate knowledge about exercise, and inadequate guidance from healthcare providers regarding physical activity are significant factors contributing to this lack of physical activity[5]. Physical inactivity has negative effects on IBD patients, including comorbidities such as cardiovascular disease, sarcopenia, osteoporosis, and mental health disorders. Physical rehabilitation and structured exercise are insufficiently implemented in the IBD population despite providing statistically significant improvements in disease activity, health outcomes, physical function, functional capacity, and overall QOL[6].

Currently, there is a lack of a comprehensive guide for healthcare practitioners or a detailed evidence-based physical rehabilitation protocol for IBD[7]. It is recommended that IBD patients engage in structured exercise under medical supervision until such time that they are deemed capable of doing so independently[8]. Inpatient rehabilitation programs can improve the physical mobility of IBD patients after they reach a state of medical stability[9]. When oxygen capacity is impaired due to anemia or dehydration, structured exercise and physical rehabilitation programs should be planned to include elements such as tempo, workload distribution, and rest intervals[10]. Employing suitable rehabilitation strategies and structured exercises can mitigate complications, such as gastrointestinal bleeding, that may develop due to excessive exercise[4]. There is evidence that shows aerobic exercise regimens can provide cardiovascular and psychosocial benefits for adult IBD patients with mild to moderate disease who are in clinical remission[11].

Despite exhibiting body compositions that are comparable to those of healthy controls, IBD patients in remission exhibit reduced muscle strength and power[12]. This may be due to impaired nutrient absorption, weight loss, decreased physical activity, and the upregulation of cytokines, which leads to muscle atrophy. Reduced muscle strength and function lead to functional deficits. Resistance exercise, when prescribed appropriately, aids muscle growth and strength by stimulating the release of biologically active myokines[13]. Populations with a higher risk of sarcopenia, malnutrition, being female, or being older have higher rates of complications. The early assessment of lower extremity and core strength can play a vital role in identifying individuals with IBD who are at risk of preclinical disability. Thus, timely diagnosis and treatment of musculoskeletal problems in these patients can effectively prevent future dependence and disability[14].

Both aerobic and resistance exercise training have been scientifically validated as effective approaches for improving cardiovascular health as well as muscle mass and strength. Numerous chronic conditions, including cardiovascular disease, diabetes, and rheumatoid arthritis, are believed to be impacted by insufficient physical activity. Interleukins (IL),

leukocytes, tumor necrosis factor, and visceral fat play a significant role in the pathogenesis and progression of IBD[15]. The potential benefits of exercise for individuals with IBD are linked to its anti-inflammatory properties. Physical activity is recognized for its ability to reduce inflammation by releasing anti-inflammatory myokines, such as IL-6, into the circulation in large quantities during exercise[16]. Furthermore, exercise may contribute to positive outcomes by reducing adipose tissue in the abdominal and intestinal regions[17]. Patients with IBD have concerns about engaging in physical exercise, unlike healthy individuals. While several studies have demonstrated the safety and effectiveness of physical activity in IBD patients, these studies primarily included patients with mild disease or those in remission[18].

The most common extra-intestinal symptom in IBD patients is articular involvement, with a prevalence of 17%-39% [19]. Although it is often characterized by involvement of axial joints, it may also be associated with peripheral arthritis such as synovitis and/or dactylitis and/or enthesopathy. IBD-related arthritis rehabilitation aims to prevent disability and deformity. Physical therapy is crucial for enhancing spinal mobility to prevent spinal deformities that could impair breathing and cause disability.

Patients with IBD have a higher risk of fractures compared to the general population. Adequate nutrition and a combination of progressive low-impact and resistance exercise training are important for preventing and treating osteoporosis[3].

IBD patients exhibit alterations in body composition due to the catabolic state caused by chronic inflammation and the use of treatments such as corticosteroids. Therefore, IBD patients frequently develop sarcopenia. The prevalence of sarcopenia is reported to be 52% and 37% in Crohn's disease and ulcerative colitis, respectively. However, accurately determining the precise prevalence is challenging[20]. Sarcopenia exerts a significant impact on QOL, duration of hospitalization, surgical outcomes, and mortality. This is an important consideration in the management of IBD, as it impacts both prognosis and treatment. Although sarcopenia is defined as age-related loss of muscle mass, it is known that it can also occur as a result of chronic illness, inactivity, and inflammation[21]. Research suggests that patients with sarcopenia may benefit from personalized resistance and aerobic exercise regimens[22].

Prehabilitation is a multimodal (nutritional, exercise, and psychological) intervention that aims to increase functional ability in the presence of impending physical stress (e.g., surgery). While most studies have concentrated on cancer patients, this strategy could have a significant impact on IBD patients with refractory disease awaiting elective surgery [23].

The epidemiology of IBD varies significantly across different regions of the globe. Although IBD is more prevalent in Western societies, which are characterized by higher socioeconomic status and improved health conditions, there has been a significant rise in the incidence and prevalence of IBD in developing countries like Türkiye[24].

In developed countries, the average rate of non-adherence to treatment is 50%, and in developing countries, this figure is even higher. Furthermore, IBD has been associated with an increase in disease activity, relapse, diminished response to medical agents, higher morbidity and mortality, and increased health expenditure[25]. To address these concerns, tele-health services may be employed, which are both feasible and acceptable in terms of patient education, evaluation of treatment compliance, and emphasizing the need for personalized rehabilitation programs. More comprehensive research is necessary to utilize these systems for improving healthcare access in developing nations[26].

CONCLUSION

In IBD patients, several factors such as reduced bone mineral density, diminished muscle strength, decreased aerobic capacity, and sarcopenia are often not adequately screened or treated during medical evaluation. A multidisciplinary approach helps address the comorbidities accompanying IBD and reduces subsequent morbidity and mortality. Furthermore, personalized rehabilitation programs can help alleviate the economic burden and frequency of disease exacerbations that are associated with the condition. Therefore, it is imperative that healthcare professionals prioritize these factors and ensure that they are addressed appropriately to improve the overall health outcomes of patients. To obtain optimal results, it is essential to collaborate with a team of specialized professionals, including a gastroenterologist, surgeon, rheumatologist, and physiatrist. This editorial aims to underline the significance of developing a specific rehabilitation program for IBD patients and assessing them from a comprehensive perspective. This is crucial in mitigating morbidity and mortality along with tackling the issue of physical inactivity among IBD patients, as was highlighted in the main article by Stafie *et al*[27].

FOOTNOTES

Author contributions: Eyigor S designed the overall concept and outline of the manuscript; Ata BN contributed to the discussion and design of the manuscript; Eyigor S and Ata BN contributed to the writing and editing of the manuscript, illustrations, and review of the literature.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: Türkiye

ORCID number: Benil Nesli Ata 0000-0003-0900-0069; Sibel Eyigor 0000-0002-9781-2712.

S-Editor: Li L

L-Editor: A

P-Editor: Zhang XD

REFERENCES

- Bilski J**, Mazur-Bialy AI, Wierdak M, Brzozowski T. The impact of physical activity and nutrition on inflammatory bowel disease: the potential role of cross talk between adipose tissue and skeletal muscle. *J Physiol Pharmacol* 2013; **64**: 143-155 [PMID: 23756389]
- Bondurri A**, Maffioli A, Danelli P. Pelvic floor dysfunction in inflammatory bowel disease. *Minerva Gastroenterol Dietol* 2015; **61**: 249-259 [PMID: 26603727]
- Elia J**, Kane S. Adult Inflammatory Bowel Disease, Physical Rehabilitation, and Structured Exercise. *Inflamm Bowel Dis* 2018; **24**: 2543-2549 [PMID: 29850914 DOI: 10.1093/ibd/izy199]
- Narula N**, Fedorak RN. Exercise and inflammatory bowel disease. *Can J Gastroenterol* 2008; **22**: 497-504 [PMID: 18478136 DOI: 10.1155/2008/785953]
- Gravina AG**, Pellegrino R, Durante T, Palladino G, D'Onofrio R, Mammone S, Arboreto G, Auletta S, Imperio G, Ventura A, Romeo M, Federico A. Inflammatory bowel diseases patients suffer from significant low levels and barriers to physical activity: The "BE-FIT-IBD" study. *World J Gastroenterol* 2023; **29**: 5668-5682 [PMID: 38077160 DOI: 10.3748/wjg.v29.i41.5668]
- Geraci A**, Tomasello G, Sabetta SP. Orthopaedic experience on inflammatory bowel disease (Lesniowski-Crohn's disease and ulcerative colitis). *Ortop Traumatol Rehabil* 2010; **12**: 430-434 [PMID: 21057150]
- Manabe I**. Chronic inflammation links cardiovascular, metabolic and renal diseases. *Circ J* 2011; **75**: 2739-2748 [PMID: 22067929 DOI: 10.1253/circj.CJ-11-1184]
- Nathan I**, Norton C, Czuber-Dochan W, Forbes A. Exercise in individuals with inflammatory bowel disease. *Gastroenterol Nurs* 2013; **36**: 437-442 [PMID: 24304528 DOI: 10.1097/SGA.0000000000000005]
- Delaney CP**, Zutshi M, Senagore AJ, Remzi FH, Hammel J, Fazio VW. Prospective, randomized, controlled trial between a pathway of controlled rehabilitation with early ambulation and diet and traditional postoperative care after laparotomy and intestinal resection. *Dis Colon Rectum* 2003; **46**: 851-859 [PMID: 12847356 DOI: 10.1007/s10350-004-6672-4]
- Dean E**. Oxygen transport deficits in systemic disease and implications for physical therapy. *Phys Ther* 1997; **77**: 187-202 [PMID: 9037219 DOI: 10.1093/ptj/77.2.187]
- Elsenbruch S**, Langhorst J, Popkirowa K, Müller T, Luedtke R, Franken U, Paul A, Spahn G, Michalsen A, Janssen OE, Schedlowski M, Dobos GJ. Effects of mind-body therapy on quality of life and neuroendocrine and cellular immune functions in patients with ulcerative colitis. *Psychother Psychosom* 2005; **74**: 277-287 [PMID: 16088265 DOI: 10.1159/000086318]
- Geerling BJ**, Badart-Smook A, Stockbrügger RW, Brummer RJ. Comprehensive nutritional status in patients with long-standing Crohn disease currently in remission. *Am J Clin Nutr* 1998; **67**: 919-926 [PMID: 9583850 DOI: 10.1093/ajcn/67.5.919]
- Costamagna D**, Costelli P, Sampaolesi M, Penna F. Role of Inflammation in Muscle Homeostasis and Myogenesis. *Mediators Inflamm* 2015; **2015**: 805172 [PMID: 26508819 DOI: 10.1155/2015/805172]
- Ward LM**, Ma J, Rauch F, Benchimol EI, Hay J, Leonard MB, Matzinger MA, Shenouda N, Lentle B, Cosgrove H, Scharke M, Konji VN, Mack DR. Musculoskeletal health in newly diagnosed children with Crohn's disease. *Osteoporos Int* 2017; **28**: 3169-3177 [PMID: 28791436 DOI: 10.1007/s00198-017-4159-0]
- van Langenberg DR**, Della Gatta P, Hill B, Zacharewicz E, Gibson PR, Russell AP. Delving into disability in Crohn's disease: dysregulation of molecular pathways may explain skeletal muscle loss in Crohn's disease. *J Crohns Colitis* 2014; **8**: 626-634 [PMID: 24332699 DOI: 10.1016/j.crohns.2013.11.024]
- Das DK**, Graham ZA, Cardozo CP. Myokines in skeletal muscle physiology and metabolism: Recent advances and future perspectives. *Acta Physiol (Oxf)* 2020; **228**: e13367 [PMID: 31442362 DOI: 10.1111/apha.13367]
- Bilski J**, Mazur-Bialy A, Brzozowski B, Magierowski M, Zahradnik-Bilska J, Wójcik D, Magierowska K, Kwiecien S, Mach T, Brzozowski T. Can exercise affect the course of inflammatory bowel disease? Experimental and clinical evidence. *Pharmacol Rep* 2016; **68**: 827-836 [PMID: 27255494 DOI: 10.1016/j.pharep.2016.04.009]
- Klare P**, Nigg J, Nold J, Haller B, Krug AB, Mair S, Thoeniger CK, Christle JW, Schmid RM, Halle M, Huber W. The impact of a ten-week physical exercise program on health-related quality of life in patients with inflammatory bowel disease: a prospective randomized controlled trial. *Digestion* 2015; **91**: 239-247 [PMID: 25823689 DOI: 10.1159/000371795]
- Peluso R**, Manguso F, Vitiello M, Iervolino S, Di Minno MN. Management of arthropathy in inflammatory bowel diseases. *Ther Adv Chronic Dis* 2015; **6**: 65-77 [PMID: 25729557 DOI: 10.1177/2040622314563929]
- Dhaliwal A**, Quinlan JL, Overthrow K, Greig C, Lord JM, Armstrong MJ, Cooper SC. Sarcopenia in Inflammatory Bowel Disease: A Narrative Overview. *Nutrients* 2021; **13** [PMID: 33671473 DOI: 10.3390/nu13020656]
- Cruz-Jentoft AJ**, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, Cooper C, Landi F, Rolland Y, Sayer AA, Schneider SM, Sieber CC, Topinkova E, Vandewoude M, Visser M, Zamboni M; Writing Group for the European Working Group on Sarcopenia in Older People 2 (EWGSOP2), and the Extended Group for EWGSOP2. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing* 2019; **48**: 601 [PMID: 31081853 DOI: 10.1093/ageing/afz046]
- Gold SL**, Raman M, Sands BE, Ungaro R, Sabino J. Review article: Putting some muscle into sarcopenia-the pathogenesis, assessment and clinical impact of muscle loss in patients with inflammatory bowel disease. *Aliment Pharmacol Ther* 2023; **57**: 1216-1230 [PMID: 37051722 DOI: 10.1111/apt.17498]
- Bak MTJ**, Ruiterkamp MFE, van Ruler O, Campmans-Kuijpers MJE, Bongers BC, van Meeteren NLU, van der Woude CJ, Stassen LPS, de Vries AC. Prehabilitation prior to intestinal resection in Crohn's disease patients: An opinion review. *World J Gastroenterol* 2022; **28**: 2403-

- 2416 [PMID: 35979261 DOI: 10.3748/wjg.v28.i22.2403]
- 24 **Can G**, Poşul E, Yılmaz B, Can H, Korkmaz U, Ermiş F, Kurt M, Dağlı Ü. Epidemiologic features of inflammatory bowel disease in Western Blacksea region of Turkey for the last 10 years: retrospective cohort study. *Korean J Intern Med* 2019; **34**: 519-529 [PMID: 29695147 DOI: 10.3904/kjim.2015.310]
- 25 **Chan W**, Chen A, Tiao D, Selinger C, Leong R. Medication adherence in inflammatory bowel disease. *Intest Res* 2017; **15**: 434-445 [PMID: 29142511 DOI: 10.5217/ir.2017.15.4.434]
- 26 **Hommel KA**, Hente E, Herzer M, Ingerski LM, Denson LA. Telehealth behavioral treatment for medication nonadherence: a pilot and feasibility study. *Eur J Gastroenterol Hepatol* 2013; **25**: 469-473 [PMID: 23325274 DOI: 10.1097/MEG.0b013e32835c2a1b]
- 27 **Stafie R**, Singeap AM, Rotaru A, Stanciu C, Trifan A. Bridging the gap: Unveiling the crisis of physical inactivity in inflammatory bowel diseases. *World J Gastroenterol* 2024; **30**: 1261-1265 [PMID: 38596487 DOI: 10.3748/wjg.v30.i10.1261]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

