

Supplementary Table 1 Sensitivity analysis of patients with available baseline and EGG variables, n (%)

Characteristics	Included, n = 76	Excluded, n = 17	P
Age in years²	52 (25.5)	51 (31)	
18–49	35 (41.2%)	7 (41.2%)	0.792
≥ 50	41 (58.8%)	10 (58.8%)	
Sex			
Male	30 (39.5%)	7 (41.2%)	1.000
Female	46 (60.5%)	10 (58.8%)	
BMI Asia-Pacific category			
Underweight	5 (6.6%)	0 (0%)	N/A ^d
Normal	11 (14.5%)	2 (11.8%)	
Overweight	15 (19.7%)	0 (0%)	
Obesity I	29 (38.2%)	1 (5.9%)	
Obesity II	16 (21.1%)	0 (0%)	
Missing	0 (0%)	14 (82.4%)	
T2DM status			
Non-T2DM	38 (50.0%)	6 (35.3%)	0.297
T2DM	38 (50.0%)	11 (64.7%)	
EGG result			
Bradygastria	47 (61.8%)	7 (41.2%)	0.247
Normogastria	18 (23.7%)	7 (41.2%)	
Tachygastria	11 (14.5%)	3 (17.6%)	

²Median (interquartile range). ^dMajority missing data on excluded group and the P value could be calculated. BMI: Body mass index; EGG: Electrogastrography; T2DM: Type 2 diabetes mellitus.