

World Journal of *Radiology*

World J Radiol 2024 November 28; 16(11): 629-707



ORIGINAL ARTICLE

Retrospective Study

- 629 Use of the vertebrae and iliac bone as references for localizing the appendix vermiformis in computed tomography
Ozturk MO, Resorlu M, Aydin S, Memis KB
- 638 Spectra of intracranial diseases in Chinese military pilots (cadets) unqualified for transfer to pilot modified high performance aircraft
Zhao Y, Gao D, Liu YB, Xue JJ, Lu X, Dong JJ, Zhang Y, Zeng J
- 644 Pancreatic volume change using three dimensional-computed tomography volumetry and its relationships with diabetes on long-term follow-up in autoimmune pancreatitis
Shimada R, Yamada Y, Okamoto K, Murakami K, Motomura M, Takaki H, Fukuzawa K, Asayama Y

Prospective Study

- 657 Right-to-left shunt detection *via* synchronized contrast transcranial Doppler combined with contrast transthoracic echocardiography: A preliminary study
Yao MJ, Zhao YY, Deng SP, Xiong HH, Wang J, Ren LJ, Cao LM
- 668 Ultra-low-dose chest computed tomography with model-based iterative reconstruction in the analysis of solid pulmonary nodules: A prospective study
O'Regan PW, Harold-Barry A, O'Mahony AT, Crowley C, Joyce S, Moore N, O'Connor OJ, Henry MT, Ryan DJ, Maher MM

CASE REPORT

- 678 Afferent loop syndrome of a patient with recurrent fever: A case report
Yuan J, Zhang YJ, Wen W, Liu XC, Chen FL, Yang Y
- 683 Successful treatment of small bowel phytobezoar using double balloon enterolithotripsy combined with sequential catharsis: A case report
Lu BY, Zeng ZY, Zhang DJ
- 689 Acute respiratory distress syndrome caused by demulsifier poisoning: A case report
Yang KY, Cui ZX

LETTER TO THE EDITOR

- 696 Carbon ion radiation therapy in prostate cancer: The importance of dosage
Treechairusame T, Taweasedt PT
- 700 Optimizing clinical decision-making for ruptured intracranial aneurysms: Current applications and future directions of computed tomography angiography
Le XY, Zhang JR, Feng JB, Li CM

- 703** Relationship between pancreatic morphological changes and diabetes in autoimmune pancreatitis: Multimodal medical imaging assessment has important potential

Zhang QB, Liu D, Feng JB, Du CQ, Li CM

ABOUT COVER

Editorial Board Member of *World Journal of Radiology*, Mustafa Zuhair Mahmoud Alhassen, BSc, MSc, PhD, Professor, Department of Radiological Sciences, College of Applied Medical Sciences, King Khalid University, Abha 61421, Saudi Arabia. malhassen@kku.edu.sa

AIMS AND SCOPE

The primary aim of *World Journal of Radiology* (*WJR*, *World J Radiol*) is to provide scholars and readers from various fields of radiology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJR mainly publishes articles reporting research results and findings obtained in the field of radiology and covering a wide range of topics including state of the art information on cardiopulmonary imaging, gastrointestinal imaging, genitourinary imaging, musculoskeletal imaging, neuroradiology/head and neck imaging, nuclear medicine and molecular imaging, pediatric imaging, vascular and interventional radiology, and women's imaging.

INDEXING/ABSTRACTING

The *WJR* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2024 Edition of Journal Citation Reports® cites the 2023 journal impact factor (JIF) for *WJR* as 1.4; JIF without journal self cites: 1.4; 5-year JIF: 1.8; JIF Rank: 133/204 in radiology, nuclear medicine and medical imaging; JIF Quartile: Q3; and 5-year JIF Quartile: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Wen-Bo Wang*; Production Department Director: *Si Zhao*; Cover Editor: *Jia-Ping Yan*.

NAME OF JOURNAL

World Journal of Radiology

ISSN

ISSN 1949-8470 (online)

LAUNCH DATE

January 31, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Thomas J Vogl

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1949-8470/editorialboard.htm>

PUBLICATION DATE

November 28, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Use of the vertebrae and iliac bone as references for localizing the appendix vermiformis in computed tomography

Muhsin Ozgun Ozturk, Mustafa Resorlu, Sonay Aydin, Kemal Bugra Memis

Specialty type: Radiology, nuclear medicine and medical imaging

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade C

Novelty: Grade A

Creativity or Innovation: Grade B

Scientific Significance: Grade B

P-Reviewer: Chen H

Received: March 15, 2024

Revised: October 1, 2024

Accepted: November 8, 2024

Published online: November 28, 2024

Processing time: 257 Days and 1.2 Hours



Muhsin Ozgun Ozturk, Sonay Aydin, Kemal Bugra Memis, Department of Radiology, Erzincan Binali Yıldırım University, Faculty of Medicine, Erzincan 24000, Türkiye

Mustafa Resorlu, Department of Radiology, Canakkale Onsekiz Mart University, Faculty of Medicine, Canakkale 17000, Türkiye

Corresponding author: Muhsin Ozgun Ozturk, MD, Researcher, Department of Radiology, Erzincan Binali Yıldırım University, Faculty of Medicine, Başbağlar, Hacı Ali Akın Cd. No. 32 24100 Erzincan Merkez/Erzincan, Erzincan 24000, Türkiye. ozturk_bul@hotmail.com

Abstract

BACKGROUND

The appendix vermiformis is a part of the gastrointestinal tract, situated in the lower right quadrant of the abdomen. Acute appendicitis, acute inflammation of the appendix vermiformis, is the most common cause of acute abdomen requiring surgical intervention. Although computed tomography (CT) offers high diagnostic efficacy in assessing the appendix across various anatomical positions, it also involves radiation exposure. Reducing exposure factors and narrowing the field of view (FOV) are ways to decrease the radiation dose to the patient. To narrow the FOV, appendix locations within the population must be defined using metric markers.

AIM

To determine the location of the appendix vermiformis on CT using the vertebrae and the right iliac bone as anatomical landmarks.

METHODS

This retrospective study examined 470 patients presenting with abdominal pain who underwent abdominal CT scans between January 01, 2015 and January 01, 2018. Forty-three patients were excluded due to various reasons. The most superior and inferior points and the origin of the appendix were measured separately in relation to the vertebrae and right iliac bone for localization. The population was divided into normal and acute appendicitis groups, and the relationship between appendix location and anthropometric parameters relationship was examined. *P* values below 0.05 were considered statistically significant.

RESULTS

The final analysis included 427 adult patients (206 females and 221 males) with a

mean age of 42.1 ± 19.5 years. An ascending appendix course was the most common (90.4%). The appendix ranged from the L2 vertebral body level to the coccygeal vertebral level relative to the vertebrae. The appendix ranged between (–) 140.5 mm and (+) 87.4 mm relative to the right iliac bone. A negative correlation was found between patient age, height, body mass index, and the highest and lowest points of the appendix in regard to the vertebrae.

CONCLUSION

The study's findings unveiled the locations of the appendix in the population in relation to the bony anatomical landmarks. These data can be used as the basis for future research aimed at reducing patient exposure to ionizing radiation.

Key Words: Appendix; Computerized tomography; Vertebrae; Ionizing radiation; Ilium

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: It is important to determine the location of the appendix vermiformis in the population using anatomical reference points. Therefore, when there is a suspicion of acute appendicitis and computed tomography is required, the field of view can be adjusted accordingly, potentially reducing the radiation dose given to the patient. Utilizing bony structures as references may offer some advantages. In particular, using the iliac bone as a reference, as opposed to the vertebrae, appears to be a more efficient approach because it is not affected by variables such as age, sex, and body mass index.

Citation: Ozturk MO, Resorlu M, Aydin S, Memis KB. Use of the vertebrae and iliac bone as references for localizing the appendix vermiformis in computed tomography. *World J Radiol* 2024; 16(11): 629-637

URL: <https://www.wjgnet.com/1949-8470/full/v16/i11/629.htm>

DOI: <https://dx.doi.org/10.4329/wjr.v16.i11.629>

INTRODUCTION

The appendix vermiformis is a part of the gastrointestinal tract and is located in the lower right quadrant of the abdomen. It has a worm-like appearance and originates from the posteromedial wall of the cecum in embryological life, around 2 cm below the terminal ileum[1,2]. The length of the appendix varies between 2 and 20 cm, with an average of 9 cm[1-3].

Atypical positions of the cecum and appendix may be encountered[4]. The relationship between the base of the appendix and the cecum is consistent in terms of location. However, the distal end of the appendix can be found in various locations, such as retrocecal, subcecal, pelvic, preileal, retroileal, right paracolic, promontoric, or subhepatic areas. These variations play an important role in the pain localization of acute appendicitis[5,6]. In Wakeley's postmortem study of 10000 cases, the frequencies of appendix localization were as follows: Retrocecal 65.3%, pelvic 31%, subcecal 2.3%, and preileal 1%[7]. However, other studies have found the most common position of the appendix to be pelvic[5,8].

Acute appendicitis, the acute inflammation of the appendix vermiformis, is the most common cause of acute abdomen requiring surgical treatment[9,10]. It is most commonly observed in adolescents and young adults and peaks in the second and third decades of life[11]. It usually starts with mild periumbilical visceral pain, which shifts to the right lower quadrant approximately 8 hours after onset. Atypical pain progression and localization can sometimes occur. Radiological imaging modalities are frequently used in the diagnosis of acute appendicitis[12].

The radiological imaging methods used in the diagnosis of acute appendicitis are ultrasonography (US), computed tomography (CT), and, rarely, magnetic resonance imaging[13]. US is usually the first modality used and may be useful in diagnosis; however, it has disadvantages, as its accuracy is highly dependent on patient, device, and operator-related factors[13,14]. In contrast, CT is independent of these factors and maintains diagnostic efficacy in patients with variable appendix positions. Furthermore, severe pain is not a limiting factor for CT scans, allowing for the evaluation of other pathologies in the differential diagnosis of acute appendicitis. The main disadvantages of CT are exposure to ionizing radiation and nephrotoxic and allergic effects associated with contrast media[15].

Obtaining diagnostically adequate images with minimal radiation exposure forms the basis of the "As Low as Reasonably Achievable (ALARA)" principle[16]. Dose modulation, kilovoltage peak adjustments, and reduction of the imaging area are some of the methods that can help reduce radiation exposure during CT scans[17]. Several studies have investigated the effectiveness of low-dose CT for diagnosing acute appendicitis by reducing exposure factors. In the study by Keyzer *et al*[18], which compared low-dose and standard-dose CT for diagnosing acute appendicitis, no difference was observed in appendix visibility or diagnostic accuracy. Similarly, Fefferman *et al*[19] found no significant difference between standard-dose and low-dose CT techniques. In another study, Sippola *et al*[20] found that a low-dose protocol showed no significant difference in diagnostic accuracy compared to the standard protocol, with a significant reduction in the mean effective radiation dose between the two techniques. These studies highlight the unnecessary dose burden associated with CT scans performed to avoid unnecessary appendectomies. In the aforementioned studies, radiation doses were lowered by adjusting exposure parameters, and the results were compared with standard-dose images[18-20]. To further minimize radiation exposure, we recommend narrowing the field of view (FOV) to prevent unnecessary

radiation exposure to the patient. This can only be accomplished by using metric markers to accurately locate the appendix in the population. Our study aimed to define the location of the appendix vermiformis using these metric markers.

MATERIALS AND METHODS

Patients who presented to the emergency department with abdominal pain and underwent non-contrast or contrast-enhanced abdominal CT between January 01, 2015 and January 01, 2018, were included in this retrospective study. This study was approved by the institutional ethics committee (No. 2011-KAEK-27/2019-E.1900162336) and carried out in compliance with the principles outlined in the Declaration of Helsinki. The requirement for informed consent was waived by the ethics committee, as all patient identification information was removed after data were collected and analyzed.

Clinical data and images of 470 randomly chosen patients who presented to the emergency department with abdominal pain and underwent intravenous contrast-enhanced or non-contrast abdominal CT scans were evaluated. The study excluded patients ($n = 43$) whose vermiform appendix could not be distinctly visualized, those with situs inversus and rotational anomalies, and individuals with a prior history of intra-abdominal surgery (including appendectomy and other intra-abdominal procedures). Localization and measurements were conducted in a single session by a single radiologist with 7 years of experience.

All abdominal CT studies were conducted using a 64-detector multidetector CT scanner. The CT acquisition parameters were as follows: Tube voltage of 120 kV, automatic tube current modulation, collimation thickness of 0.5-2 mm, tube rotation time of 0.5-1 s, slice thickness of 3 mm, and reconstruction intervals of 1.5 mm. The routine FOV of the abdominal CT scan extended from the dome of the diaphragm superiorly to the ischial tuberosities inferiorly. Iopromide 300 mg/mL was administered at a dose of 1 mL/kg to patients undergoing contrast-enhanced CT.

To objectively determine the location of the appendix in the population included in the study, we used the vertebrae and the highest point of the right iliac bone as anatomical landmarks. The superior and inferior points of the appendix were identified separately relative to the vertebrae for population-based localization. In addition, the superior point, inferior point, and origin of the appendix were measured separately in relation to the right iliac bone. In this context, the superior point of the appendix is labeled as "ApA," the inferior point is labeled "ApB," and the appendiceal origin is labeled "Ap0." For measurements using the vertebrae and the right iliac bone as references, the population was divided into two groups: Normal appendix and acute appendicitis.

To determine the location of the appendix using the vertebrae as reference points, initially, the locations of the appendix's superior and inferior points along its course were found and marked on axial CT images. Then, these points were located on thick coronal and sagittal multiplanar reformatted images to determine their respective vertebral levels. We grouped the vertebral levels for appendix localization as follows: L1 & L1-L2 intervertebral joint; L2 & L2-L3 intervertebral joint; L3 & L3-L4 intervertebral joint; L4 & L4-L5 intervertebral joint; L5 & L5-S1 intervertebral joint; and S1, S2, S3, S4, S5, and coccygeal vertebrae. The technique for determining appendix location using vertebral references is illustrated in [Figure 1](#).

We also used the iliac bone as a reference to determine the location of the appendix with a more quantitative method. For this purpose, we found and marked the lowest, and highest points, and origin of the appendix on axial CT images. We then measured the distance in millimeters between these marked appendix points on coronal reformatted images and the highest point of the right iliac bone using maximum-intensity projection images. If the marked appendix points were superior to the highest point of the right iliac bone, the measurement was expressed as a positive (+) value, and if more inferior, the measurement was expressed as a negative (-) value in millimeters. [Figure 2](#) illustrates how the appendix was localized in reference to the right iliac bone.

The course of the appendix was also investigated in the study population. Appendix vermiformis was classified as "ascending" when it ended up higher than its point of origin and "descending" when it ended up lower than its point of origin. The relationship between appendix location and some anthropometric parameters like height, weight, and body mass index (BMI) was also investigated; BMI was calculated using the formula: Weight/height^2 (kg/m^2). Age and sex data were also collected.

Data were analyzed using SPSS version 20.0 (IBM Corp., Armonk, NY, United States). The distribution of the data was examined for normality using the Kolmogorov-Smirnov test. Numerical data following a normal distribution were presented as mean \pm standard deviation, while categorical data were shown as numbers and percentages [n (%)]. The χ^2 test was used to compare percentages between the acute appendicitis and normal appendix groups. The mean values of Ap0, ApA, and ApB were compared between male and female patients, as well as between the acute appendicitis and normal appendix groups, using Student's *t*-test. Pearson correlation analysis was used to examine potential correlations between ApA and ApB values and anthropometric characteristics (weight, height, and BMI) and age. A *P* value less than 0.05 was considered statistically significant.

RESULTS

The final analysis comprised 427 patients, of which 206 (48.2%) were female and 221 (51.8%) were male. The mean patient age was 42.1 ± 19.5 (range: 18-90) years. Of the 427 CT scans, 200 (46.8%) were performed with intravenous contrast media administration, and 227 (53.2%) were performed without contrast media. The mean BMI was $25.6 \text{ kg/m}^2 \pm 6.4$ (range: 16.8-40.5) kg/m^2 .




Figure 1 Determining the localization of the highest point of the appendix using the vertebrae as a reference. A: The highest point of the appendix vermiformis is detected (orange arrow) on axial images; B: Subsequently, the highest point is verified on coronal images (orange arrow); C: The highest point (orange arrow) is marked on axial images with reference lines; D and E: The vertebral level of the marked point is determined using coronal and sagittal thick multiplanar reconstruction images, respectively, with bone window settings. Subsequently, the same assessment is conducted to identify the lowest point of the appendix. L: Lumbar vertebra; S: Sacral vertebra.

Acute appendicitis was diagnosed on CT in 68 (15.9%) patients, and the diagnosis was subsequently confirmed in surgical specimens. In 359 (84.1%) patients, the appendix appeared normal on CT, and clinical follow-up confirmed the absence of acute appendicitis. In the overwhelming majority of patients ($n = 325$) who were clinically confirmed not to have acute appendicitis, abdominal pain resolved spontaneously or with simple analgesics. In the remaining patients ($n = 34$), a combination of clinical and CT findings led to alternative diagnoses of omental infarction ($n = 2$), sigmoid diverticulitis ($n = 5$), epiploic appendagitis ($n = 7$), sigmoid volvulus ($n = 2$), mesenteric panniculitis ($n = 5$), acute cholecystitis ($n = 5$), inflammatory-infectious bowel disease ($n = 6$), and ischemic bowel disease ($n = 2$).

The appendix had an ascending course in the majority of patients, both with and without acute appendicitis (89.7% and 90.5%, respectively). There was no statistically significant difference in the course of the appendix between the normal appendix and acute appendicitis groups ($P = 0.41$). The numbers and percentages of ascending and descending appendix courses for the overall population, the appendicitis group, and the normal appendix group are displayed in Table 1.

Measurements of the appendiceal origin (Ap0), the highest point of the appendix (ApA), and the lowest point of the appendix (ApB) were made in relation to the highest point of the right iliac bone in the normal appendix group, acute appendicitis group, and the overall population ($n = 427$). The mean \pm SD, minimum, and maximum values for these measurements in each group are shown in Table 2. No statistically significant difference was found between the acute appendicitis and normal appendix groups in the ApB location relative to the right iliac bone ($P = 0.19$). However, a significant difference was found between these groups in the Ap0 and ApA locations relative to the right iliac bone ($P = 0.04$ and $P = 0.04$; respectively).

No significant statistical difference was found between male and female patients in the measurements of ApA and ApB in relation to the right iliac bone ($P = 0.08$ and $P = 0.21$, respectively). However, a significant difference was found between male and female patients in the measurements of Ap0 in relation to the right iliac bone ($P = 0.03$). The mean values of Ap0, ApA, and ApB relative to the right iliac bone for male and female patients are shown in Table 3.

In the normal appendix population, ApA ranged from the L2 vertebral body to the coccygeal vertebrae, while in the acute appendicitis population, ApA ranged from the L2 vertebra to the S3-S4 intervertebral joint. In the normal appendix population, ApB ranged from the L2 vertebral body to the coccygeal vertebrae, and in the acute appendicitis population, ApB ranged from the L4 vertebral body to the coccygeal vertebrae. ApA was most commonly found at the L5 and L5-S1 intervertebral joint level (34.4%), while ApB was most commonly found at the S1 vertebral body level (22.5%). No significant difference was found between the acute appendicitis group and the normal appendix group in the localization of ApA and ApB relative to the vertebrae ($P = 0.19$). ApA was located at or below the level of the L3 vertebra in all patients with acute appendicitis.

The numbers and percentages of ApA and ApB localizations in relation to the vertebrae are shown in Table 4.

Table 1 Courses of the appendix in the overall population, acute appendicitis group, and normal appendix group, *n* (%)

Groups	Ascending	Descending	Total
Acute appendicitis	61 (89.7)	7 (10.3)	68 (15.9)
Normal appendix	325 (90.5)	34 (9.5)	359 (84.1)
Total	386 (90.4)	41 (9.6)	427 (100)

Table 2 Minimum, maximum, and mean \pm standard deviation of appendix location in relation to the right iliac bone in the overall population, acute appendicitis group, and normal appendix group, mm

Groups	Ap0 (origin)	ApA (highest point of the appendix)	ApB (lowest point of the appendix)
Acute appendicitis group (<i>n</i> = 68)	(-) 34.1 \pm 29.1; (-) 100.8 - (+) 44.2	(-) 17.1 \pm 27.8; (-) 84.8 - (+) 48.3	(-) 60.9 \pm 27.9; (-) 121.6 - (+) 39.1
Normal appendix group (<i>n</i> = 359)	(-) 42.3 \pm 31.5; (-) 126.0 - (+) 40.1	(-) 24.2 \pm 34.1; (-) 126.0 - (+) 87.4	(-) 59.8 \pm 30.2; (-) 140.5 - (+) 44.1
Total (<i>n</i> = 427)	(-) 41.0 \pm 32.7; (-) 126.0 - (+) 44.2	(-) 23.0 \pm 30.8; (-) 126.0 - (+) 87.4	(-) 60.0 \pm 30.2; (-) 140.5 - (+) 44.1

No statistically significant difference was found between the acute appendicitis and normal appendix groups in ApB location relative to the right iliac bone ($P = 0.19$). Significant difference was found between the acute appendicitis and normal appendix groups in Ap0 and ApA locations relative to the right iliac bone ($P = 0.04$ and $P = 0.04$; respectively).

Table 3 Ap0, ApA, and ApB relative to the highest point of the right iliac bone in male and female patients, mm

Sex	Ap0 (mean \pm SD)	ApA (mean \pm SD)	ApB (mean \pm SD)
Female (<i>n</i> = 206)	(-) 48.1 \pm 30.0	(-) 26.2 \pm 30.4	(-) 60.4 \pm 31.1
Male (<i>n</i> = 221)	(-) 34.3 \pm 29.7	(-) 20.0 \pm 29.8	(-) 59.6 \pm 30.8
Total (<i>n</i> = 427)	(-) 41.0 \pm 32.7	(-) 23.0 \pm 30.8	(-) 60.0 \pm 30.2

No significant statistical difference was found between male and female patients in the measurements of ApA and ApB relative to the right iliac bone ($P = 0.08$ and $P = 0.21$; respectively). A significant difference was found between male and female patients in the measurements of Ap0 relative to the right iliac bone ($P = 0.03$).

Table 4 ApA and ApB localizations relative to the vertebrae in the overall population, *n* (%)

Vertebral levels	ApA	ApB
L1& L1-L2 intervertebral joint distance	0 (0)	0 (0)
L2& L2-L3 intervertebral joint distance	8 (1.9)	1 (0.2)
L3& L3-L4 intervertebral joint distance	29 (6.8)	2 (0.5)
L4 & L4-L5 intervertebral joint distance	88 (20.6)	20 (4.7)
L5 & L5-S1 intervertebral joint distance	147 (34.4)	81 (19.0)
S1 vertebra	95 (22.2)	96 (22.5)
S2 vertebra	34 (8.0)	81 (19.0)
S3 vertebra	19 (4.4)	66 (15.5)
S4 vertebra	4 (0.9)	56 (13.1)
S5 vertebra & coccygeal vertebrae	3 (0.7)	24 (5.6)

L: Lumbar vertebra; S: Sacral vertebra.




Figure 2 Determining the localization of the highest point of the appendix using the right iliac bone as a reference. A: The highest point of the appendix vermiformis is detected (orange arrow) on axial images; B: The highest point is verified on coronal images (orange arrow); C: The highest point (orange arrow) is marked on axial images with reference lines; D: The distance between this designated point and the highest point of the right iliac bone is measured using the thick coronal multiplanar reconstruction images in the bone window. The distance is measured from the point where the horizontal line intersects with the highest point of the right iliac bone. If the designated point is situated above the highest point of the right iliac bone, it is denoted as positive (+), and if it is positioned below, it is denoted as negative (-), with measurements in mm. In this patient, the highest point of the appendix was localized at -33 mm relative to the highest point of the right iliac bone. Subsequently, using the same technique, the origin and lowest points of the appendix vermiformis were determined. L: Lumbar vertebra; S: Sacral vertebra.

A weak negative correlation was found between patient age, height, BMI, and the vertebral levels of ApA and ApB. No correlation was found between patient age, height, BMI, and the position of ApA and ApB relative to the right iliac bone. The *P* and *R* values are shown in Table 5.

DISCUSSION

Our aim in the study was to define the location of the appendix vermiformis using metric markers. In order to determine the appendix's location, we used the vertebrae and the right iliac bone as references. By using these references, we examined the frequency and variability of appendix locations, examining associations with anthropometric parameters and the presence of acute appendicitis. We found a weak negative correlation between anthropometric parameters and the location of the uppermost (ApA) and lowermost (ApB) points of the appendix relative to the vertebrae; however, no correlation was found between anthropometric parameters and the location of the uppermost (ApA) and lowermost (ApB) points of the appendix relative to the right iliac bone. No significant difference was found between male and female patients in the localization of ApA and ApB in relation to the right iliac bone.

The need for medical imaging is growing faster than in the majority of other healthcare-related fields[21]. Today's clinicians have become increasingly dependent on imaging to confirm their clinical judgments before planning the management of acute abdominal conditions, including acute appendicitis. CT scan usage has risen sharply in the last two decades because of its reported higher accuracy compared to US for diagnosing acute appendicitis. As a result, many institutions now prefer CT as the primary imaging modality in suspected cases of acute appendicitis[22].

A consensus known as ALARA has been established in radiological practice to restrict the amount of radiation that patients are exposed to, with broad and well-founded support[23]. Reducing the FOV is one approach to decreasing radiation exposure. Therefore, it is important to know the radiological location of the appendix in the population[24].

The scientific literature contains studies that investigate the visibility rates of appendix vermiformis[25,26], various positions of the appendiceal tip (such as retrocecal or pelvic)[25,26], and uncommon locations of the appendix vermiformis[27] on CT. However, to our knowledge, only one other study in the literature has used bony structures as a reference for appendix localization on CT. In their study, Davis *et al*[28] localized the highest point of the appendix in reference to the vertebrae in a pediatric population. Our study helps to fill this gap by providing similar data for the adult

Table 5 Relationship between anthropometric measurements and the locations of ApA and ApB relative to the vertebrae and the right iliac bone

Appendix locations		Age	Height	BMI
Relative to the vertebrae	ApA	$P = 0.007/R = -0.13$	$P = 0.006/R = -0.13$	$P = 0.044/R = -0.09$
	ApB	$P = 0.028/R = -0.10$	$P = 0.007/R = -0.13$	$P = 0.01/R = -0.12$
Relative to the right iliac bone	ApA	$P = 0.33/R = -0.12$	$P = 0.19/R = -0.10$	$P = 0.12/R = -0.08$
	ApB	$P = 0.47/R = -0.09$	$P = 0.53/R = -0.13$	$P = 0.23/R = -0.14$

The relationship between anthropometric measurements and the locations of the appendix relative to the vertebrae and right iliac bone is demonstrated with P and R values. BMI: Body mass index.

population.

In our study, the appendix was found to have an ascending course most frequently in the acute appendicitis group, the normal appendix group, and the overall population (89.7%, 90.5%, and 90.4%, respectively). To the best of our knowledge, this is the first study to use radiological techniques to assess whether the appendix is ascending or descending in its course. Also, this is the first publication to examine the impact of acute appendicitis on the course of the appendix, with no significant difference found between the acute appendicitis and normal appendix groups. Understanding whether the appendix is positioned in an upward (ascending) or downward (descending) direction in the majority of the population can provide valuable guidance for surgical procedures.

To our knowledge, no previous study has determined the location of the appendix relative to the iliac bone in the population. In our study, the appendix was observed between (–) 140.5 mm and (+) 87.4 mm relative to the highest point of the right iliac bone. A significant difference was found between the acute appendicitis and normal appendix groups in the Ap0 and ApA locations relative to the right iliac bone, and the appendix was located more superiorly in the acute appendicitis group compared to the normal appendix group. This finding suggests that the appendix tends to be in a more upright position when inflamed. We also investigated the relationship between appendix localization and anthropometric parameters and found that measurements based on the right iliac bone did not vary with these parameters, thus supporting the iliac bone as a reliable reference point.

We were the first to use vertebrae as a reference for locating the appendix in the adult population. Davis *et al*[28] localized the highest point in the course of the appendix (expressed as ApA in our study) in the pediatric population, also using the vertebrae as a reference. Although their study was conducted in the pediatric population, they found ApA most frequently in the L5 vertebral level, as in our study. We found that ApA was located at or below the L3 vertebra in all patients with appendicitis, aligning with the findings of Davis *et al*[28]. In our study, the location of the appendix relative to the spine was inversely correlated with patient height, age, and BMI.

We localized the appendix relative to both the vertebrae and the right iliac bone in the population, creating a controlled trial within itself because it involves two distinct anatomical measurements. While there was variability in localization of the appendix relative to the vertebrae with age, height, and BMI, no variability was detected with these parameters according to the right iliac bone. This shows that the right iliac bone may be a more reliable reference point in determining appendix localization in the population. When localization was assessed relative to the vertebrae, no significant difference was detected between the acute appendicitis and normal appendix groups. However, when localized relative to the right iliac bone, Ap0 and ApA were significantly more superior in the acute appendicitis group compared to the normal appendix group, indicating that inflammation causes the appendix to become more erect. The vertebral body heights may conceal minute locational variations, which explains why the vertebral level does not alter much in cases of acute appendicitis.

Our study has some limitations. As with all retrospective and cross-sectional studies, an increase in sample size could alter the results. We conducted a single-center study, and all measurements were conducted by a single radiologist in a single session; therefore, intra- and interobserver variability could not be evaluated. Moreover, since the pediatric population was not included, we were unable to compare findings between adult and pediatric groups. Another limitation was the lack of intraoperative measurements to validate our results, as all measurements relied solely on CT images.

CONCLUSION

The highest point of the appendix was most frequently located at the L5 & L5-S1 intervertebral joint (34.4%), while the lowest point was most frequently found at the S1 vertebra level (22.5%). Relative to the right iliac bone, the origin, highest point, and lowest point of the appendix measured (–) 41.0 ± 32.7 mm, (–) 23.0 ± 30.8 mm, and (–) 60.0 ± 30.2 mm; respectively. While localization analyses based on vertebrae varied according to anthropometric variables, those based on the right iliac bone showed no variability. These findings provide a basis for future studies aimed at reducing ionizing radiation exposure.

FOOTNOTES

Author contributions: Ozturk MO and Resorlu M designed and performed research, analyzed the data, and added radiological images; Ozturk MO and Aydin S wrote the manuscript; Ozturk MO and Memis KB revised the manuscript.

Institutional review board statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Ethics Committee of Canakkale Onsekiz Mart University Ethics Committee of Clinical Research (2011-KAEK-27/2019-E.1900162336).

Informed consent statement: Owing to the methodology of the study, the requirement for obtaining patient consent was waived by the ethics committee.

Conflict-of-interest statement: The authors declare no conflicts of interest.

Data sharing statement: The datasets generated and/or analyzed during the current study are not publicly available due to the risk of breach of patient data privacy but are available from the corresponding author on reasonable request.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country of origin: Türkiye

ORCID number: Muhsin Ozgun Ozturk 0009-0000-3356-3481; Mustafa Resorlu 0000-0002-2941-8879; Sonay Aydin 0000-0002-3812-6333; Kemal Bugra Memis 0009-0007-6746-3906.

S-Editor: Lin C

L-Editor: Filipodia

P-Editor: Wang WB

REFERENCES

- 1 Deshmukh S, Verde F, Johnson PT, Fishman EK, Macura KJ. Anatomical variants and pathologies of the vermiform appendix. *Emerg Radiol* 2014; **21**: 543-552 [PMID: 24570122 DOI: 10.1007/s10140-014-1206-4]
- 2 Schumpelick V, Steinau G, Schlüper I, Prescher A. Surgical embryology and anatomy of the diaphragm with surgical applications. *Surg Clin North Am* 2000; **80**: 213-239, xi [PMID: 10685150 DOI: 10.1016/s0039-6109(05)70403-5]
- 3 Hodge BD, Kashyap S, Khorasani-Zadeh A. Anatomy, Abdomen and Pelvis: Appendix. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing, 2024 [PMID: 29083761]
- 4 Aminova GG. [Structure and cytoarchitectonic of the lymphoid tissue of the Appendix of man in elderly and senile ages]. *Adv Gerontol* 2018; **31**: 273-279 [PMID: 30080336]
- 5 Ghorbani A, Forouzes M, Kazemifar AM. Variation in Anatomical Position of Vermiform Appendix among Iranian Population: An Old Issue Which Has Not Lost Its Importance. *Anat Res Int* 2014; **2014**: 313575 [PMID: 25295193 DOI: 10.1155/2014/313575]
- 6 de Souza SC, da Costa SRMR, de Souza IGS. Vermiform appendix: positions and length – a study of 377 cases and literature review. *J Coloproctology* 2015; **35**: 212-216 [DOI: 10.1016/j.jcol.2015.08.003]
- 7 Wakeley CP. The Position of the Vermiform Appendix as Ascertained by an Analysis of 10,000 Cases. *J Anat* 1933; **67**: 277-283 [PMID: 17104423]
- 8 Denjalić A, Delić J, Delić-Custendil S, Muminagić S. [Variations in position and place of formation of appendix vermiformis found in the course of open appendectomy]. *Med Arh* 2009; **63**: 100-101 [PMID: 19537667]
- 9 Constantin M, Petrescu L, Mătanie C, Vrancianu CO, Niculescu AG, Andronic O, Bolocan A. The Vermiform Appendix and Its Pathologies. *Cancers (Basel)* 2023; **15** [PMID: 37568688 DOI: 10.3390/cancers15153872]
- 10 Bhangu A, Søreide K, Di Saverio S, Assarsson JH, Drake FT. Acute appendicitis: modern understanding of pathogenesis, diagnosis, and management. *Lancet* 2015; **386**: 1278-1287 [PMID: 26460662 DOI: 10.1016/S0140-6736(15)00275-5]
- 11 Hardin DM Jr. Acute appendicitis: review and update. *Am Fam Physician* 1999; **60**: 2027-2034 [PMID: 10569505]
- 12 Hwang ME. Sonography and Computed Tomography in Diagnosing Acute Appendicitis. *Radiol Technol* 2018; **89**: 224-237 [PMID: 29298941]
- 13 Di Saverio S, Podda M, De Simone B, Ceresoli M, Augustin G, Gori A, Boermeester M, Sartelli M, Coccolini F, Tarasconi A, De' Angelis N, Weber DG, Tolonen M, Birindelli A, Biffl W, Moore EE, Kelly M, Søreide K, Kashuk J, Ten Broek R, Gomes CA, Sugrue M, Davies RJ, Damaskos D, Leppäniemi A, Kirkpatrick A, Peitzman AB, Fraga GP, Maier RV, Coimbra R, Chiarugi M, Sganga G, Pisanu A, De' Angelis GL, Tan E, Van Goor H, Pata F, Di Carlo I, Chiara O, Litvin A, Campanile FC, Sakakushev B, Tomadze G, Demetashvili Z, Latifi R, Abu-Zidan F, Romeo O, Segovia-Lohse H, Baiocchi G, Costa D, Rizoli S, Balogh ZJ, Bendinelli C, Scalea T, Ivatury R, Velmahos G, Andersson R, Kluger Y, Ansaloni L, Catena F. Diagnosis and treatment of acute appendicitis: 2020 update of the WSES Jerusalem guidelines. *World J Emerg Surg* 2020; **15**: 27 [PMID: 32295644 DOI: 10.1186/s13017-020-00306-3]
- 14 Love BE, Camelo M, Nouri S, Kriger D, Ludi D, Nguyen H. Ultrasound Accuracy in Diagnosing Appendicitis in Obese Pediatric Patients. *Am Surg* 2017; **83**: 1063-1067 [DOI: 10.1177/000313481708301010]
- 15 Bahrami M, Mirgaloyebayat H, Mohajeri Z, Mohammadi H, Afshari SA, Fazeli P, Masaeli D, Nourian SMA. The diagnostic value of the

- computed tomography scan and ultrasonography in acute appendicitis. *Am J Nucl Med Mol Imaging* 2023; **13**: 11-17 [PMID: 36923598]
- 16 **Frane N**, Bitterman A. Radiation Safety and Protection. 2023 May 22. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan- [PMID: 32491431]
 - 17 **Dixon MT**, Loader RJ, Stevens GC, Rowles NP. An evaluation of organ dose modulation on a GE optima CT660-computed tomography scanner. *J Appl Clin Med Phys* 2016; **17**: 380-391 [PMID: 27167255 DOI: 10.1120/jacmp.v17i3.5724]
 - 18 **Keyzer C**, Tack D, de Maertelaer V, Bohy P, Gevenois PA, Van Gansbeke D. Acute appendicitis: comparison of low-dose and standard-dose unenhanced multi-detector row CT. *Radiology* 2004; **232**: 164-172 [PMID: 15155894 DOI: 10.1148/radiol.2321031115]
 - 19 **Fefferman NR**, Bomsztyk E, Yim AM, Rivera R, Amodio JB, Pinkney LP, Strubel NA, Noz ME, Rusinek H. Appendicitis in children: low-dose CT with a phantom-based simulation technique--initial observations. *Radiology* 2005; **237**: 641-646 [PMID: 16170015 DOI: 10.1148/radiol.2372041642]
 - 20 **Sippola S**, Virtanen J, Tammilehto V, Grönroos J, Hurme S, Niiniviita H, Lietzen E, Salminen P. The Accuracy of Low-dose Computed Tomography Protocol in Patients With Suspected Acute Appendicitis: The OPTICAP Study. *Ann Surg* 2020; **271**: 332-338 [PMID: 30048324 DOI: 10.1097/SLA.0000000000002976]
 - 21 **Bercovich E**, Javitt MC. Medical Imaging: From Roentgen to the Digital Revolution, and Beyond. *Rambam Maimonides Med J* 2018; **9** [PMID: 30309440 DOI: 10.5041/RMMJ.10355]
 - 22 **Debnath J**, George RA, Ravikumar R. Imaging in acute appendicitis: What, when, and why? *Med J Armed Forces India* 2017; **73**: 74-79 [PMID: 28123249 DOI: 10.1016/j.mjafi.2016.02.005]
 - 23 **Winder M**, Owczarek AJ, Chudek J, Pilch-Kowalczyk J, Baron J. Are We Overdoing It? Changes in Diagnostic Imaging Workload during the Years 2010-2020 including the Impact of the SARS-CoV-2 Pandemic. *Healthcare (Basel)* 2021; **9** [PMID: 34828603 DOI: 10.3390/healthcare9111557]
 - 24 **Vogiatzi T**, Menz R, Verna C, Bornstein MM, Dagassan-Berndt D. Effect of field of view (FOV) positioning and shielding on radiation dose in pediatric CBCT. *Dentomaxillofac Radiol* 2022; **51**: 20210316 [PMID: 35762346 DOI: 10.1259/dmfr.20210316]
 - 25 **Altunkas A**, Aktas F, Ozmen Z, Albayrak E, Demir O. The Normal Vermiform Appendix in Adults. *J Anat Soc India* 2022; **71**: 225-233 [DOI: 10.4103/jasi.jasi_135_19]
 - 26 **Turkoglu H**, Onur MR, Poyraz AK, Kocakoc E. Evaluation of normal appendix vermiformis in adults with multidetector computed tomography. *Clin Imaging* 2012; **36**: 758-762 [PMID: 23154006 DOI: 10.1016/j.clinimag.2012.01.028]
 - 27 **Zacharzewska-Gondek A**, Szczurowska A, Guziński M, Słasiadek M, Bładowska J. A pictorial essay of the most atypical variants of the vermiform appendix position in computed tomography with their possible clinical implications. *Pol J Radiol* 2019; **84**: e1-e8 [PMID: 31019588 DOI: 10.5114/pjr.2018.81158]
 - 28 **Davis J**, Roh AT, Petterson MB, Kopelman TR, Matz SL, Gridley DG, Connell MJ. Computed tomography localization of the appendix in the pediatric population relative to the lumbar spine. *Pediatr Radiol* 2017; **47**: 301-305 [PMID: 28091700 DOI: 10.1007/s00247-016-3773-x]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

